

DAFTAR PUSTAKA

1. Kusuma et al. Neutrofil to lymphocyte Ratio and platelet to lymphocyte Ratio as an inflammatory Biomarker in Predicting the Severity of Secondary Brain Injury: A Review Article. *Macedonian Journal of Medical Science*. 2020; 8(F):272-282.
2. Ristanto R, Indra R, Poeranto S, Setyorini I. Akurasi revised trauma score sebagai prediktor mortality Pasien cedera kepala. *Jurnal Kesehatan Hesti Wira Sakti*. 2016;4(2):76-90.
3. Yuksen et al. Clinical Faktors Predictive for Intrakranial Hemorrhage in Mild Head Injury. *Hindawi Neurology Research Internasional*.2017;20:1-5.
4. Tim Riskesdas 2018. Laporan Nasional RISKESDAS 2018. Jakarta; 2019.
5. Dewan MC, Rattani A, Gupta S, Baticulon RE, Hung YC, Punchak M, et al. Estimating the global incidence of traumatic brain injury. *J Neurosurg*. 2018.
6. Hazeldine J, Lord JM, Belli A. Traumatic brain injury and peripheral immune suppression: primer and prospectus. *Frontiers in Jurnal Sinaps*. 2018; 1(2):20-28.
7. Siahaya N, Laura B, S Huwae, Ony W. Prevalensi kasus cedera kepala berdasarkan klasifikasi derajat keparahannya pada pasien rawat inap di RSUD DR. M. Hallusy Ambon pada tahun 2018: *Mollusca Medica*. 2020;12:1-7.
8. Watanitanon A, Lyons VH, Lele AV, Krishnamoorthy V, Chandee T, et al. Clinical epidemiology of adults with moderate traumatic brain injury. *Journal critical care medicine*. 2018;46:781-787.
9. Baum J, Entezami P, Shah K, Medhkour A. Predictors of outcome in traumatic brain injury. *World Neurosurg*.2016;90:525-529.
10. Gyoneva S, Ransohoff RM. Inflammatory reaction after traumatic brain injury: Therapeutic potential of targeting cell-cell communication by chemokines. *Trends Pharmacol Sci*.2015;36(7):471–480.
11. Tangkudung G, Sampoerna JM, Khosama H. Hubungan jumlah leukosit dengan gangguan fungsi eksekutif pada cedera kepala ringan – sedang. *Neurona*.2016;33:164-170
12. Morgan E, Bankole O, Kanu O, Ojo O, Poluyi E. Comparison of the Predictive Strength of Total White Blood Cell Count within 24 hours on Outcome of Traumatic Brain Injury with Glasgow Coma Score and Pupillary Reactivity. *J Clin Case Rep Trials*. 2018;1(1):2-13.
13. Kastilong M, Subrata I, Tangkudung G, Khosama I. Neutrophyl Lymphocyte Rasio and The Outcome Of Traumatic Brain Injury. *Jurnal Sinaps*. 2018;1(2): 20-28.
14. Chen W, Yang J, Li Bingbing, Peng G, Li Tianfei. Neutrofil to lymphocyte ratio as a novel predictor of outcome in patients with severe traumatic brain injury. *J Head Trauma Rehabil*. 2017;1-7.
15. Sabouri et al. Neutrofil to lymphocyte ratio as a novel predictor of outcome in patients with severe traumatic brain injury. *World Neurosurgery*. 2020;4:185.
16. Yeatts SD, Palesch YY, Temkin N. Biostatistical Issues in TBI Clinical Trials [Internet]. Second Edi. *Handbook of Neuroemergency Clinical Trials: Second Edition*. Elsevier Inc. 2018;2:167–185.
17. Callahan BP, Rabb CH. Traumatic Brain Injury [Internet]. *Abernathy's Surgical Secrets*.

Elsevier Inc.; 2009. 99–104 p. Available from: <http://dx.doi.org/10.1016/B978-0-12-648082-5.50007-1>

18. Chen J, Qu X, Li Z, Zhang D, Hou L. Peak neutrophil-to-lymphocyte ratio correlates with clinical outcome in patients with severe traumatic brain injury. *Neurocrit Care*. 2019;30(2):334-339.
19. Muhammad Zaki, Syaiful S, Hesti L. Comparison of Lymphocyte Neutrophil Ratio in Head Injury Patients with Blood and without Intrakranial Bleeding in Dr. M. Djamil Hospital. *Biomedical Journal of Indonesia*. 2021;7(1).
20. Najem et al. Traumatic brain injury: classification, models, and markers. *Biochem. Cell Biol*. 2018;96:391-406.
21. Shaikh F, Waseem M (2021). Head Trauma. <https://www.ncbi.nlm.nih.gov/books/NBK430854/>. Diakses pada Oktober 2021
22. Advance Trauma Life Support, Student Course Manual.10 th edition. Chicago:American College of Surgeons;2018.
23. Ng SY, Lee AYW. Traumatic Brain Injuries: Pathophysiology and potential therapeutic targets. *Front Cell Neurosci*. 2019;13:528.
24. Galgano M, Toshkezi G, Qiu X, Russel T, Chin L. Traumatic brain injury: current treatment strategies and future endeavors. *Cell Transplantation* 2017;26(7): 1118-1130.
25. Crupi R, Cordaro M, Cuzzocrea S, Impellizzeri D. Management of traumatic brain injury: from present to future. *Antioxidant*. 2020; 9: 297.
26. Mark S Greenberg. *Handbook of neurosurgery*. 9 th ed. New York: Thieme; 2020.
27. National Academies of Sciences, Engineering, and Medicine. *Evaluation of the Disability Determination Process for Traumatic Brain Injury in Veterans*. Washington (DC): National Academies Press (US); 2019. <https://www.ncbi.nlm.nih.gov/books/NBK542595/>
28. Dash HH, Chavali S. Management of traumatic brain injury patients. *Korean Journal of Anesthesiology*. 2018; 71(1):12-21.
29. Carney N, Totten AM, O'Reilly C, Ulman JS, Hawryluk GWJ, Bell MJ. *Guidelines for the management of severe traumatic brain injury*, fourth edition.
30. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: A brief overview. *J Head Trauma Rehabil*. 2006;21(5):375–8.
31. Young RJ, Destian S. Imaging of traumatic intrakranial hemorrhage. *Neuroimaging Clin N Am*. 2002;12(2):189–204
32. Yuan F, Ding J, Chen H, Guo Y, Wang G, Gao WW, et al. Predicting progressive hemorrhagic injury after traumatic brain injury: Derivation and validation of a risk score based on admission characteristics. *J Neurotrauma*. 2012;29(12):2137–42.
33. Arifin M., Risdianto A. *Cedera Kepala*. Jakarta, Indonesia: Sagung Seto; 2 013. 53–58.
34. Warner C, Engelhard K. Pathophysiology of traumatic brain injury. *BJA*. 2007; 99:4-9.
35. Dash HH, Chavali S. Management of Traumatic Brain Injury Patient. *Korean J Anesthesiol*. 2018;12–21.
36. American College of Surgeons. *Best practice in the management of traumatic*. Table of

Contents. 2015; (January):1–29. Available from: [https://www.facs.org/~media/files/quality_programs/trauma/tqip/traumatic brain injury guidelines.ashx](https://www.facs.org/~media/files/quality_programs/trauma/tqip/traumatic_brain_injury_guidelines.ashx)

37. Capizzi A, Woo J, Verduzco-Gutierrez M. Traumatic Brain Injury: An Overview of Epidemiology, Pathophysiology, and Medical Management. *Med Clin North Am.*2020;104(2):213–38.
38. Liu S, Liu X, Chen S, Xiao Y, Zhuang W. Neutrophil–lymphocyte ratio predicts the outcome of intracerebral hemorrhage: A meta-analysis. 2019;98:26.
39. Al-Mufti F et al. Admission neutrophil–lymphocyte ratio predicts delayed cerebral ischemia following aneurysmal subarachnoid hemorrhage. 2019:1-7.
40. Siwicka-Gieroba et al. The Neutrophil/Lymphocyte count ratio predicts mortality in severe traumatic brain injury patients. *Journal of Clinical Medicine.* 2019;1:1-4.
41. Liu YW, Li S, Dai SS. Neutrophils in traumatic brain injury (TBI): friend or foe? *Journal of Neuroinflammation.*2018;15:146.
42. Krisanty M et al, Neutrophil-Lymphocyte rasio (NLR) as an output outcome predictor in moderate-severe head injury at Sanglah General Hospital, Denpasar, Indonesia. *Intisari sains medis.*2021;2:576-581.
43. Oliveira R., Araujo S. Glasgow outcome scale at hospital discharge as a prognostic index in patients with severe traumatic brain injury. *Arq Neuropsiquiatry.*2012;70(8):604–8.
44. Kimball R, Shachar E, Eyerly-Webb S, Patel DM, Spader H. Using the neutrophil-to-lymphocyte ratio to predict outcomes in pediatric patients with traumatic brain injury. *Clinical Neurology and Neurosurgery.* 2020.
45. Xie Q, Huang W, Shen L, Wang M, Liu K, Liu F. Combination of neutrophil-to-lymphocyte ratio and admission glasgow coma scale score is independent predictor of clinical outcome in diffuse axonal injury. *World Neurosurgery.* 2021; 152: e118-e127,
46. Lizen N, Dharmajaya R, Sastrodiningrat AG, Mahyudani. Correlation of neutrophil lymphocyte ratio to clinical outcomes based on 6 months surveillance on traumatic brain injury patients at Haji Adam Malik General Hospital (2018-2020). *Romanian Neurosurgery.* 2021; 34: 503-7.
47. Dahlan MS. Besar Sample dan Cara Pengambilan Sampel dalam Penelitian Kedokteran dan Kesehatan. 3rd ed. *Epidemiologi Indonesia.* Jakarta: Salemba Medika; 2010
48. Einarsen et al. Moderate Traumatic Brain Injury: Clinical Characteristics and a Prognostic Model of 12-Month Outcome. *World Neurosurgery.* 2018; 114: e1199-e1210.
49. D Laskowitz, G Grant. *Translational Research in Traumatic Brain Injury.* Boca Raton.NCBI Bookshelf. 2016
50. Jochems D, Rein E, Niemeijer M, Heijl M, Es MA, Nijboer T. Incidence, causes and consequences of moderate and severe traumatic brain injury as determined by Abbreviated

Injury Score in the Netherlands. *Natureportfolio*. 2021;11.

51. Ghosh A, Pal R, Mishra R, Cincu R, Moscote-Salazar LR, Agrawal A. Leukocyte neutrophil ratio (NLR) as biomarker for neuroinflammation in patients with intracerebral haemorrhage and traumatic brain injury. *Ro J Neurol*. 2021; 20(3).
52. Kodliwadmath HB, Koppad SN, Desai M, Badiger SP. Correlation of Glasgow outcome score to Glasgow coma score assessed at admission. *Int Surg J*. 2016 Nov;3(4):1959-1963.
53. Klemenc-Ketis Z, Bacovnik-Jansa U, Ogorevc M, Kersnik J. Outcome predictors of Glasgow Outcome Scale score in patients with severe traumatic brain injury. *Turkish Journal of Trauma & Emergency Surgery*. 2011;17 (6):509-515.
54. Islam MR, Khan MSK, Rahman MM, Talukder MH, Karim MR, Salam A. Comparison of Glasgow Outcome Scale (GOS) and Glasgow Coma Scale (GCS) between Surgical and Conservative Management of Spontaneous Supratentorial Intracerebral Hemorrhage Patients: A Randomized Control Trial. *Journal of Current and Advance Medical Research*. 2018; 5(2): 49-54.
55. Maiden MJ, Cameron PA, Rosenfeld JV, Cooper DJ, McLellan S, Gabbe BJ. Long-Term Outcomes Following Severe Traumatic Brain Injury in Older Adults. *AJRCCM*. 2019.
56. Dikmen S, Machamer J, Manley GT, Yuh EL, Nelson LD, Temkin NR. Functional Status Examination versus Glasgow Outcome Scale Extended as Outcome Measures in Traumatic Brain Injuries: How Do They Compare?. *Journal of Neurotrauma*. 2019; 36:2423–2429
57. Lok J, Leung W, Murphy S, Butler W, Noviski N, Lo EH. Intracranial Hemorrhage – Mechanisms of Secondary Brain Injury. *Acta Neurochir Suppl*. 2011;111: 63–69
58. Wang F, Wang L, Jiang T, Xia J, Xu F, Shen L, Kang W, et al. Neutrophil-to-Lymphocyte Ratio Is an Independent Predictor of 30-Day Mortality of Intracerebral Hemorrhage Patients: A Validation Cohort Study. *Neurotoxicity Research*. 2018; 34:347–352.