

Daftar Pustaka

- Akkurt HE, Gümüs H, Göksu H, Odaba OF, Yilmaz H. 2015. Gabapentin Treatment for Neuropathic Pain in a Child with Sciatic Nerve Injury. Hindawi Publishing Corporation:1-4.
- Amin B, Hosseinzadeh H, 2012. Evaluation of aqueous and ethanolic extracts of saffron, *Crocus sativus L.*, and its constituents, safranal and crocin in allodynia and hyperalgesia induced by chronic constriction injury model of neuropathic pain in rats. *Fitoterapia*;83(5):888-95.
- Anhui et al., 2021. The P2X7 receptor is involved in diabetic neuropathic pain hypersensitivity mediated by TRPV1 in the rat dorsal root ganglion. <https://www.frontiersin.org/articles/10.3389/fnmol.2021.663649/full>
- AustinPJ, Wu A, Moalem-Taylor G, 2012. Chronic constriction of the sciatic nerve and pain hypersensitivity testing in rats. *Journal of visualized experiments: JoVE*, (61).
- Baron R, 2006. Mechanisms of disease: neuropathic pain a clinical perspective. *Nat Clin Pract Neuro*:2(2):95-106.
- Bennett GJ, Xie YK, 1988. A Peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man. *Pain*;33:87-107.
- Bernier LP, Ase AR, Séguéla P, 2018. P2X receptor channels in chronic pain pathways. *Br. J. Pharmacol.*;175:2219–2230. doi: 10.1111/bph.13957.
- Bhardwaj HC, Muthuraman A, Hari KSL, Navis S, 2016. Antioxidative and Anti-Inflammatory Potentials of Ambroxol in Ameliorating Vincristine Induced Peripheral Neuropathic Pain in Rats. *J. Neuroinfect Dis* ;7(202):2.
- Borda AP, Sonnek FC, Fonteyne V, Papaioannou EG, 2013. Guidelines on Pain Management and Palliative Care. European Association of Urologi. P70.
- Bridges D, Thompson SWN, Rice ASC, 2001. Mechanism of Neuropathic Pain. *British Journal of Anesthesia*;87(1):12-6.
- Burnstock G, 2017. Purinergic Signalling: Therapeutic Developments. *Front Pharmacol*. Sep 25;8:661. doi: 10.3389/fphar.2017.00661. eCollection.

Calejesan AA, Kim SJ, Zhuo M, 2000. Descending facilitatory modulation of a behavioral nociceptive response by stimulation in the adult rat anterior cingulate cortex. *Eur J Pain* 2000, 4:83-96.

Carniglia et al., 2017. Neuropeptides and Microglial Activation in Inflammation, Pain, and Neurodegenerative Diseases. *Mediators Inflamm*, 5048616.

Casey KL, 1999. Forebrain mechanisms of nociception and pain: analysis through imaging. *Proc Natl Acad Sci USA*, 96:7668-7674.

Chavan SS, Pavlov VA, Tracey KJ, 2017. Mechanisms and therapeutic relevance of neuro-immune communication. *Immunity*. 46:927–42.

Childers WE, Baudy RB, 2007. N-methyl-D-aspartate antagonists and neuropathic pain: the search for relief. *J Med Chem*, 50:2557 2562.

Chessell IP, Hatcher JP, Bountra C, Michel AD, Hughes JP, Green et al., 2005. Disruption of P2X7 purinoceptor gene abolishes chronic inflammatory and neuropathic pain. *Pain*. 2005;114:386–396. doi: 10.1016/j.pain.2005.01.002.

Collo G, Neidhart S, Kawashima E, Kosco-Vilbois M, North RA, Buell G, 1997. Tissue distribution of the P2X7 receptor: *Neuropharmacology*.;36:1277–1283. doi: 10.1016/S0028-3908(97)00140-8.

Colomar A, Marty V, Médina C, Combe C, Parnet P, Amédée T, 2003. Maturation and release of interleukin-1 β by lipopolysaccharide-primed mouse Schwann cells require the stimulation of P2X7 receptors. *J Biol Chem*;278:30732–30740. doi: 10.1074/jbc.M304534200.

Costigan M, Scholz J, Woolf CJ, 2009. Neuropathic pain: a maladaptive response of the nervous system to damage. *Annual review of neuroscience*;32:1-32.

Coull JA, Boudreau D, Bachand K, Prescott SA, Nault F, Sik A et al., 2003. Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature*; 424:938-942.

Cuzzocrea S, Galen K, Lightfoot R, Masini E, Muscoli C, Mollace et al., 2006. A newly identified role for superoxide in inflammatory pain. *J Pharmacol Exp Ther*. 309:869–878.

Dahlan MS, 2009. Besar Sampel dan Cara Pengambilan Sampel Dalam Penelitian Kedokteran dan Kesehatan. Jakarta: Salemba Medika, 34.

Davis KD, Kwan CL, Crawley AP, Mikulis DJ, 1998. Functional MRI study of thalamic and cortical activations evoked by cutaneous heat, cold, and tactile stimuli. *J Neurophysiol.*;80(3):1533–1546. doi:10.1152/jn.1998.80.3.1533

DeLeo JA, Colburn RW, Nichols M, Malhotra A, 1996. Interleukin-6-mediated hyperalgesia/allodynia and increased spinal IL-6 expression in a rat mononeuropathy model. *J Interferon Cytokine Res.* 16:695–700.

Duan B, Wu LJ, Yu YQ, Ding Y, Jing L, Xu et al., 2007. Upregulation of acid-sensing ion channel ASIC1a in spinal dorsal horn neurons contributes to inflammatory pain hypersensitivity. *J Neurosci*;27:11139-11148.

Dubovy P, Brazda V, Klusakova I, Hradilova-Svizenska I, 2013. Bilateral elevation of interleukin-6 protein and mRNA in both lumbar and cervical dorsal root ganglia following unilateral chronic compression injury of the sciatic nerve. *J Neuroinflammation.* 10:55.

Forstenpointner et al., 2019. Neuroimaging Of Cold Allodynia Reveals A Central Disinhibition Mechanism Of Pain. Doverpress.doi.org :12 Pages 3055—3066, <https://doi.org/10.2147/JPR.S216508>.

Garthwaite J, 1991. Glutamate, nitric oxide and cell-cell signalling in the nervous system. *Trends Neurosci.* Feb; 14(2):60-7. [PubMed] [Ref list].

Gordon CJ, 1993. Temperature Regulation in Laboratory Animals. New York: Cambridge University Press.

González-Castro TB, Genis-Mendoza AD, Tovilla-Zárate CA, Juárez-Rojop IE, López-Narvaez ML, Pérez-Hernández et al., 2019. Association between polymorphisms of NOS1, NOS2 and NOS3 genes and suicidebehavior: A systematic review and meta-analysis. *Metab. Brain Dis.*, 34, 967–977. [CrossRef].

Hamza M, Wang XM, Wu T, Brahim JS, Rowan JS, 2010. A Dionne, R. Nitric Oxide is Negatively Correlated to Pain during Acute Inflammation. *Mol. Pain*, 6, 55. [CrossRef]

Huang LY, Gu Y, Chen Y, 2013. Communication between neuronal somata and satellite glial cells in sensory ganglia. *Glia*;61:1571–1581. doi:

10.1002/glia.22541. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

Hou D.X., Yanagita T., Uto T., Masuzaki S., Fujii M. 2005. Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: structure-activity relationship and molecular mechanism involved. *Biochem. Pharmacol.*; 70:417-25.

IASP, 2014. Epidemiology of Neuropathic Pain: How Common is Neuropathic Pain, and What Is Its Impact? International Association for the Study of Pain:1-2.

Ikeda H, Heinke B, Ruscheweyh R, Sandkuhler J, 2003. Synaptic plasticity in spinal lamina I projection neurons that mediate hyperalgesia. *Science* 299:1237-1240.

Ikeda H, Stark J, Fischer H, Wagner M, Drdla R, Jager et al., 2006. Synaptic amplifier of inflammatory pain in the spinal dorsal horn. *Science* 312:1659-1662.

Inoue K, Tsuda M, 2009. Microglia and neuropathic pain. *Glia* 57:1469-1479.

Jarvis MF, 2010. The neural-glial purinergic receptor ensemble in chronic pain states. *Trends Neurosci* 33:48–57

Ji RR, Kohno T, Moore KA, Woolf CJ, 2003. Central sensitization and LTP: do pain and memory share similar mechanisms? *Trends Neurosci* 26:696-705.

Ji RR, Xu ZZ, Gao YJ, 2014. Emerging targets in neuroinflammation-driven chronic pain. *Nat Rev drug Discov.* 13:533. 10.1038/nrd4334

Kandel ER, 2001. The molecular biology of memory storage: a dialogue between genes and synapses. *Science* 294:1030-1038.

Kang JD, Stefanovic-Racic M, McIntyre LA, Georgescu HI, Evans CH, 1997. Toward a Biochemical Understanding of Human Intervertebral Disc Degeneration and Herniation. *Spine* , 22, 1065–1073. [CrossRef] [PubMed]

Kobayashi K, Takahashi E, Miyagawa Y, Yamanaka H, Noguchi K, 2011. Induction of the P2X7 receptor in spinal microglia in a neuropathic pain

model. *Neurosci Lett.* 2011 Oct 17;504(1):57-61. doi: 10.1016/j.neulet.2011.08.058. Epub 2011 Sep 8. PMID: 21924325.

Kolios G, Brown Z, Robson RL et al., 1995. Inductible nitric oxide synthase activity and expression in a human colonic epithelial cell line, HT-29. *Br J Pharmacol*, 116: 2866–2872.

Lee KM, Jeon SM, Cho HJ, 2009. Tumor necrosis factor receptor 1 induces interleukin-6 upregulation through NF-kappaB in a rat neuropathic pain model. *Eur J Pain*, 13:794–806.

Li P, Kerchner GA, Sala C, Wei F, Huettner JE, Sheng et al., 1999. AMPA receptor-PDZ interactions in facilitation of spinal sensory synapses. *Nat Neurosci*, 2:972-977.

Li P, Wilding TJ, Kim SJ, Calejesan AA, Huettner JE, Zhuo M, 1999. Kainate-receptor-mediated sensory synaptic transmission in mammalian spinal cord. *Nature*, 397:161-164.

Li P, Zhuo M, 1998. Silent glutamatergic synapses and nociception in mammalian spinal cord. *Nature*, 393:695-698.

Luchting B, Heyn J, Woehrle T, et al., 2016. Differential expression of P2X7 receptor and IL-1 β in nociceptive and neuropathic pain. *J Neuroinflammation*, 13(1):100. Published 2016 May 4. doi:10.1186/s12974-016-0565-z.

Matricon J, Gelot A, Etienne M, Lazdunski M, Muller E, Ardid D, 2011. Spinal cord plasticity and acid-sensing ion channels involvement in a rodent model of irritable bowel syndrome. *Eur J Pain*, 15:335-343.

McCarberg BH, Billington R, 2006. Consequences of neuropathic pain: quality-of-life issues and associated costs. *Am J Manag Care*, 12:S263-268.

McMahon SB, Malcangio M, 2009. Current challenges in glia-pain biology. *Neuron*, 64:46-54.

Milligan ED, Watkins LR, 2009. Pathological and protective roles of glia in chronic pain. *Nat Rev Neurosci*, 10:23-36.

Meliala L, 2004. Terapi Rasional Nyeri. Medika Gama Press, Yogyakarta.

Mense S, 2009. "Anatomy of Nociceptors". In Basbaum, Allan, I., Bushnell, Catherine, M. (eds). Science of pain. 1st Ed. Elsevier Inc. United Kingdom. pp 11-42.

Millan MJ, 1999. The induction of pain: an integrative review. *Prog Neurobiol* 57, 1-164.

Milligan ED, Watkins LR, 2009. Pathological and protective roles of glia in chronic pain. *Nat. Rev. Neurosci*;10:23-36.

Murakami A, Kanchiku T, 2013. Anti-interleukin-6 receptor antibody reduces neuropathic pain following spinal cord injury in mice. *6(5)*: 1194–1198.

Narcisse L, Scemes E, Zhao Y, Lee SC, Brosnan CF, 2005. The cytokine IL-1 β transiently enhances P2X7 receptor expression and function in human astrocytes. *Glia*;49:245–258. doi: 10.1002/glia.20110.

Nicholson B, 2006. Differential Diagnosis: Nociceptive and Neuropathic Pain. *The American Journal of Managed Care*. P256-61.

Olson LC, Palotay JL, 1983. Epistaxis and Bullae in Cynomolgus Macaques (*Macaca fascicularis*). *Lab Anim Sci*.33:377-370. [PubMed]

Pedersen LM, Gjerstad J, 2008. Spinal cord long-term potentiation is attenuated by the NMDA-2B receptor antagonist Ro 25-6981. *Acta Physiol (Oxf)*,192:421-427.

Qu XX, Cai J, Li MJ, Chi YN, Liao FF, Liu et al., 2009. Role of the spinal cord NR2B-containing NMDA receptors in the development of neuropathic pain. *Exp Neurol*, 215:298-307.

Richner M, Jager SB, Siupka P, Vaegter CB, 2017. Hydraulic Extrusion of the Spinal Cord and Isolation of Dorsal Root Ganglia in Rodents. *J Vis Exp*. Jan 22;(119).

Sandkuhler J, 2007. Understanding LTP in pain pathways. *Mol Pain*, 3:9.

Schäfers M, Svensson CI, Sommer C, Sorkin LS, 2003. Tumor necrosis factor- α induces mechanical allodynia after spinal nerve ligation by activation of p38 MAPK in primary sensory neurons. *Journal of Neuroscience*;23(7):2517-21.

Scholz J, Woolf CJ, 2007. The neuropathic pain triad: neurons, immune cells and glia. *Nature neuroscience*;10(11):1361-8.

Sean P, Didion L, 2018. Cellular and Oxidative Mechanisms Associated with Interleukin-6 Signaling in the Vasculature.

Serizawa K, Tomizawa-Shinohara H, Magi M, Yogo K, Matsumoto Y, 2018. Anti-IL-6 receptor antibody improves pain symptoms in mice with experimental autoimmune encephalomyelitis. *J Neuroimmunol*. Jun 15;319:71-79. doi: 10.1016/j.jneuroim.2018.03.017. Epub 2018 Apr 5. PMID: 29685293.

Sim JA, Young MT, Sung HY, North RA, 2004. Surprenant A. Reanalysis of P2X7 receptor expression in rodent brain. *J Neurosci*;24:6307–6314. doi: 10.1523/JNEUROSCI.1469-04.2004.

Stefanovic-Racic M, Georgescu H, Evans C, 1994. Nitric oxide mediates suppression of cartilage proteoglycan synthesis by interleukin-1. *Biochem Biophys Res Commun*;200:142–148.

Shnayder NA, Petrova MM, Popova TE, Davidova TK, Bobrova OP, Trefilova et al., 2021. Prospects for the Personalized Multimodal Therapy Approach to Pain Management via Action on NO and NOS. <https://pubmed.ncbi.nlm.nih.gov>.

Tanaka T, Narazaki M, Ogata A, Kishimoto T, 2014. A new era for the treatment of inflammatory autoimmune diseases by interleukin-6 blockade strategy. *Semin Immunol*. 26:88.

Tang J, Ko S, Ding HK, Qiu CS, Calejesan AA, Zhuo M, 2005. Pavlovian fear memory induced by activation in the anterior cingulate cortex. *Mol Pain*, 1:6.

Tsuda, M, Shigemoto-Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S., Salter MW, et al., 2003. P2X4 receptors induced in spinal microglia tactile allodynia after nerve injury. *Nature*;424:778–783

Tsuda M, Inoue K, Salter MW, 2005. Neuropathic pain and spinal microglia: A big problem from molecules in “small” glia. *Trends Neurosci*;28:101–7.

Tsuda M, 2017. P2 receptors, microglial cytokines and chemokines, and neuropathic pain. <https://doi.org/10.1002/jnr.23816>

Vallance P, Leiper J, 2002. Blocking NO synthesis: how, where and why?. *Nat Rev Drug Discov.* 1(12):939-50

Vanderwall AG, Milligan ED, 2019. Cytokines in Pain: Harnessing Endogenous Anti-Inflammatory Signaling for Improved Pain Management. *Front Immunol.* doi: 10.3389/fimmu.2019.03009. PMID: 31921220; PMCID: PMC6935995.

Vuolteenaho K, Järvinen K, Nieminen R, Moilanen T, Knowles RG, Moilanen E, 2008. Selective iNOS inhibitor 1400W enhances anti-catabolic IL-10 and reduces destructive MMP-10 in OA cartilage. Survey of the effects of 1400W on inflammatory mediators produced by OA cartilage as detected by protein antibody array. *Clinical and Experimental Rheumatology*; 26: 275-282.

Wang S, Xu H, Zou L, Xie J, Wu H, Wu B, et al., 2016. LncRNA uc.48+ is involved in diabetic neuropathic pain mediated by the P2X3 receptor in the dorsal root ganglia. *Purinerg. Signal.* 12 139–148. 10.1007/s11302-015-9488-x.

Wei F, Li P, Zhuo M, 1999. Loss of synaptic depression in mammalian anterior cingulate cortex after amputation. *J Neurosci*; 19:9346-9354.

Wei F, Qiu CS, Kim SJ, Muglia L, Maas JW, Pineda et al., 2002. Genetic elimination of behavioral sensitization in mice lacking calmodulin-stimulated adenylyl cyclases. *Neuron*, 36:713-726.

Wei F, Vadakkan KI, Toyoda H, Wu LJ, Zhao MG, Xu et al., 2006. Calcium calmodulin-stimulated adenylyl cyclases contribute to activation of extracellular signal-regulated kinase in spinal dorsal horn neurons in adult rats and mice. *J Neurosci*; 26:851-861.

Wei F, Wang GD, Kerchner GA, Kim SJ, Xu HM, Chen et al., 2001. Genetic enhancement of inflammatory pain by forebrain NR2B overexpression. *Nat Neurosci*. 4:164-169.

Wei F, Zhuo M, 2001. Potentiation of sensory responses in the anterior cingulate cortex following digit amputation in the anaesthetised rat. *J Physiol*, 532:823-833.

Whittle BJ, 1995. Nitric oxide in physiology and pathology. *Histochem J*.27: 727–737

Widyadharma IPE, 2018. Pemberian ekstrak air umbi ubi jalar ungu (ipomoea batatas l) menyebabkan kadar malondialdehyde,PGE2,ekspresi reseptor P2X4 lebih rendah dan perilaku nyeri neuropatik lebih ringan pada tikus wistar dengan cedera saraf tepi.Udayana P: 98-99.

Woolf CJ, Salter MW, 2000. Neuronal plasticity: increasing the gain in pain. Science, 288:1765-1769.

Wu LJ, Duan B, Mei YD, Gao J, Chen JG, Zhuo et al., 2004. Characterization of acid-sensing ion channels in dorsal horn neurons of rat spinal cord. J Biol Chem,279:43716-43724.

Wu LJ, Toyoda H, Zhao MG, Lee YS, Tang J, Ko et al., 2005. Upregulation of forebrain NMDA NR2B receptors contributes to behavioral sensitization after inflammation. J Neurosci;25:11107-11116.

Wu LJ, Zhuo M, 2009. Targeting the NMDA receptor subunit NR2B for the treatment of neuropathic pain. Neurotherapeutics, 6:693-702.

Xu TL, Duan B, 2009. Calcium-permeable acid-sensing ion channel in nociceptive plasticity: a new target for pain control. Prog Neurobiol, 87:171-180.

Yoshimura M, Jessell T, 1990. Amino acid-mediated EPSPs at primary afferent synapses with substantia gelatinosa neurones in the rat spinal cord. J Physiol, 430:315-335.

Zainuddin M, 2014. Metodelogi penelitian kefarmasian dan kesehatan. Edisi 2. Airlangga University Press. Surabaya

Zhang HM, Zhou LJ, Hu XD, Hu NW, Zhang T, Liu XG, 2004. Acute nerve injury induces long-term potentiation of C-fiber evoked field potentials in spinal dorsal horn of intact rat. Sheng Li Xue Bao, 56:591-596.

Zhang WJ, Liu LY, Xu TL, 2008. Reduced potassium-chloride co-transporter expression in spinal cord dorsal horn neurons contributes to inflammatory pain hypersensitivity in rats. Neuroscience, 152:502-510.

Zarshenas MM, Petramfar P, Samani SM, Moein M, 2015. A glimpse into the pathophysiology, mechanisms, and management of neuropathic pain. Njppp;5(3):160-5.

Zhou YQ, et al., 2016. Interleukin-6: an emerging regulator of pathological pain. Journal of neuroinflammation.

Zhuo M, Gebhart GF, 1990. Spinal cholinergic and monoaminergic receptors mediate descending inhibition from the nuclei reticularis gigantocellularis and gigantocellularis pars alpha in the rat. Brain Res; 535:67-78.

Zhuo M, 2007. A synaptic model for pain: long-term potentiation in the anterior cingulate cortex. Mol Cells;23:259-271.

Zhuo M, 2007. Neuronal mechanism for neuropathic pain. Mol Pain, 3:14.

