

## DAFTAR PUSTAKA

1. Indonesia KKR. Hasil utama riskesdas 2018. Jakarta: Badan Penelitian dan Pengembangan Kesehatan, Kementerian Kesehatan Republik Indonesia. 2018;44(8):1-200.
2. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *European heart journal*. 2018;39(2):119-77.
3. Roffi M, Patrono C, Collet J-P, Mueller C, Valgimigli M, Andreotti F, et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Kardiologia Polska (Polish Heart Journal)*. 2015;73(12):1207-94.
4. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *The Lancet*. 2003;361(9351):13-20.
5. Gupta S, Gupta MM. No reflow phenomenon in percutaneous coronary interventions in ST-segment elevation myocardial infarction. *Indian heart journal*. 2016;68(4):539-51.
6. Zhu X, Shen H, Gao F, Wu S, Ma Q, Jia S, et al. Clinical Profile and Outcome in Patients with Coronary Slow Flow Phenomenon. *Cardiology research and practice*. 2019;1(1):9168153.
7. Marc MC, Iancu AC, Bălănescu ř, Dregosesc MI. Microvascular obstruction in acute myocardial infarction: an old and unsolved mystery. *Medicine and pharmacy reports*. 2019;92(3):216.
8. Niccoli G, Scalzone G, Lerman A, Crea F. Coronary microvascular obstruction in acute myocardial infarction. *European heart journal*. 2016;37(13):1024-33.
9. Vogelzang M, Vlaar PJ, Svilaas T, Amo D, Nijsten MW, Zijlstra F. Computer-assisted myocardial blush quantification after percutaneous coronary angioplasty for acute myocardial infarction: a substudy from the TAPAS trial. *European heart journal*. 2009;30(5):594-9.
10. Porto I, Hamilton-Craig C, De Maria GL, Vergallo R, Cautilli G, Galiuto L, et al. Quantitative Blush Evaluator accurately quantifies microvascular dysfunction in patients with ST-elevation myocardial infarction: Comparison with cardiovascular magnetic resonance. *American heart journal*. 2011;162(2):372-81. e2.
11. Haeck JD, Gu YL, Vogelzang M, Bilodeau L, Krucoff MW, Tijssen JG, et al. Feasibility and applicability of computer-assisted myocardial blush quantification after primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. *Catheterization and Cardiovascular Interventions*. 2010;75(5):701-6.
12. Tanboga IH, Topcu S, Aksakal E, Kalkan K, Sevimli S, Acikel M. Determinants of angiographic thrombus burden in patients with ST-segment elevation myocardial infarction. *Clinical and Applied Thrombosis/Hemostasis*. 2014;20(7):716-22.
13. Karabulut A, Uyarel H, Uzunlar B, Çakmak M. Elevated red cell distribution width level predicts worse postinterventional thrombolysis in myocardial infarction flow reflecting abnormal reperfusion in acute myocardial infarction treated with a primary coronary intervention. *Coronary artery disease*. 2012;23(1):68-72.
14. Karnad A, Poskitt TR. The automated complete blood cell count: Use of the red blood cell volume distribution width and mean platelet volume in evaluating anemia and thrombocytopenia. *Archives of internal medicine*. 1985;145(7):1270-2.

15. Sangui MB, Rödel A, Zorzo P, Borges P, Cargnin L, De Carvalho J, et al. Prognostic value of red blood cell distribution width in prediction of in-hospital mortality in patients with acute myocardial infarction. *Clinical laboratory*. 2014;60(8):1351-6.
16. Sun X-p, Chen W-m, Sun Z-j, Ding X-s, Gao X-y, Liang S-w, et al. Impact of red blood cell distribution width on long-term mortality in patients with ST-elevation myocardial infarction. *Cardiology*. 2014;128(4):343-8.
17. Osadnik T, Strzelczyk J, Hawranek M, Lekston A, Wasilewski J, Kurek A, et al. Red cell distribution width is associated with long-term prognosis in patients with stable coronary artery disease. *BMC cardiovascular disorders*. 2013;13(1):1-8.
18. Tonelli M, Sacks F, Arnold M, Moye L, Davis B, Pfeffer M. Relation between red blood cell distribution width and cardiovascular event rate in people with coronary disease. *Circulation*. 2008;117(2):163-8.
19. Felker GM, Allen LA, Pocock SJ, Shaw LK, McMurray JJ, Pfeffer MA, et al. Red cell distribution width as a novel prognostic marker in heart failure: data from the CHARM Program and the Duke Databank. *Journal of the American College of Cardiology*. 2007;50(1):40-7.
20. Ye Z, Smith C, Kullo IJ. Usefulness of red cell distribution width to predict mortality in patients with peripheral artery disease. *The American journal of cardiology*. 2011;107(8):1241-5.
21. Stiermaier T, Thiele H, Eitel I. Coronary Microvascular Obstruction: Key Factor in the Prognosis of ST-Segment-Elevation Myocardial Infarction. *2017;10(6):e006568*.
22. Iwakura K, Ito H, Kawano S, Okamura A, Tanaka K, Nishida Y, et al. Prediction of the no-reflow phenomenon with ultrasonic tissue characterization in patients with anterior wall acute myocardial infarction. *The American journal of cardiology*. 2004;93(11):1357-61.
23. Bekkers SC, Yazdani SK, Virmani R, Waltenberger J. Microvascular obstruction: underlying pathophysiology and clinical diagnosis. *Journal of the American College of Cardiology*. 2010;55(16):1649-60.
24. Constantino BT. Red cell distribution width, revisited. *Laboratory Medicine*. 2013;44(2):e2-e9.
25. Wandersee NJ, Olson SC, Holzhauer SL, Hoffmann RG, Barker JE, Hillery CA. Increased erythrocyte adhesion in mice and humans with hereditary spherocytosis and hereditary elliptocytosis. *Blood*. 2004;103(2):710-6.
26. Sergueeva A, Miasnikova G, Shah BN, Song J, Lisina E, Okhotin DJ, et al. Prospective study of thrombosis and thrombospondin-1 expression in Chuvash polycythemia. *haematologica*. 2017;102(5):e166.
27. Litvinov RI, Weisel JW. Role of red blood cells in haemostasis and thrombosis. *ISBT science series*. 2017;12(1):176-83.
28. Fogelson AL, Neeves KB. Fluid mechanics of blood clot formation. *Annual review of fluid mechanics*. 2015;47:377-403.
29. Chen H, Angerer JI, Napoleone M, Reininger AJ, Schneider SW, Wixforth A, et al. Hematocrit and flow rate regulate the adhesion of platelets to von Willebrand factor. *Biomicrofluidics*. 2013;7(6):064113.
30. Reimers RC, Sutera SP, Joist JH. Potentiation by red blood cells of shear-induced platelet aggregation: relative importance of chemical and physical mechanisms. 1984;64(6):1200-6.
31. Østerud B, Unruh D, Olsen J, Kirchhofer D, Owens III A, Bogdanov V. Procoagulant and proinflammatory effects of red blood cells on lipopolysaccharide-stimulated monocytes. *Journal of Thrombosis and Haemostasis*. 2015;13(9):1676-82.

32. Wohner N, Sótonyi P, Machovich R, Szabó L, Tenekedjiev K, Silva MM, et al. Lytic resistance of fibrin containing red blood cells. *Arteriosclerosis, thrombosis, and vascular biology*. 2011;31(10):2306-13.
33. Cines DB, Lebedeva T, Nagaswami C, Hayes V, Massefski W, Litvinov RI, et al. Clot contraction: compression of erythrocytes into tightly packed polyhedra and redistribution of platelets and fibrin. *Blood, The Journal of the American Society of Hematology*. 2014;123(10):1596-603.
34. Byrnes JR, Wolberg AS. Red blood cells in thrombosis. *Blood, The Journal of the American Society of Hematology*. 2017;130(16):1795-9.
35. Lippi G, Targher G, Montagnana M, Salvagno GL, Zoppini G, Guidi GC. Relationship between red blood cell distribution width and kidney function tests in a large cohort of unselected outpatients. *Scandinavian journal of clinical and laboratory investigation*. 2008;68(8):745-8.
36. Seyhan EC, Özgül MA, Tutar N, Ömür Im, Uysal A, Altın S. Red blood cell distribution and survival in patients with chronic obstructive pulmonary disease. COPD: *Journal of Chronic Obstructive Pulmonary Disease*. 2013;10(4):416-24.
37. Akın F, Köse N, Ayça B, Katkat F, Duran M, Uysal OK, et al. Relation between red cell distribution width and severity of coronary artery disease in patients with acute myocardial infarction. *Angiology*. 2013;64(8):592-6.
38. Lippi G, Filippozzi L, Montagnana M, Salvagno GL, Franchini M, Guidi GC, et al. Clinical usefulness of measuring red blood cell distribution width on admission in patients with acute coronary syndromes. *Clinical chemistry and laboratory medicine*. 2009;47(3):353-7.
39. Bekler A, Gazi E, Tenekecioglu E, Karaagac K, Altun B, Temiz A, et al. Assessment of the relationship between red cell distribution width and fragmented QRS in patients with non-ST elevated acute coronary syndrome. *Medical science monitor: international medical journal of experimental and clinical research*. 2014;20:413.
40. Tanboga IH, Topcu S, Nacar T, Aksakal E, Kalkan K, Kiki I, et al. Relation of coronary collateral circulation with red cell distribution width in patients with non-ST elevation myocardial infarction. *Clinical and Applied Thrombosis/Hemostasis*. 2014;20(4):411-5.
41. Osadnik T, Strzelczyk J, Bujak K, Reguła R, Wasilewski J, Fronczek M, et al. Functional polymorphism rs710218 in the gene coding GLUT1 protein is associated with in-stent restenosis. *Biomarkers in medicine*. 2015;9(8):743-50.
42. Yildiz A, Tekiner F, Karakurt A, Sirin G, Duman D. Preprocedural red blood cell distribution width predicts bare metal stent restenosis. *Coronary artery disease*. 2014;25(6):469-73.
43. Zhao K, Li Y-J, Gao S. Role of red blood cell distribution in predicting drug-eluting stent restenosis in patients with stable angina pectoris after coronary stenting. *Coronary artery disease*. 2015;26(3):220-4.
44. Fatemi O, Torguson R, Chen F, Ahmad S, Badr S, Satler LF, et al. Red cell distribution width as a bleeding predictor after percutaneous coronary intervention. *American heart journal*. 2013;166(1):104-9.
45. Salisbury AC, Amin AP, Reid KJ, Wang TY, Alexander KP, Chan PS, et al. Red blood cell indices and development of hospital-acquired anemia during acute myocardial infarction. *The American journal of cardiology*. 2012;109(8):1104-10.
46. Tziakas D, Chalikias G, Grapsa A, Gioka T, Tentes I, Konstantinides S. Red blood cell distribution width—a strong prognostic marker in cardiovascular disease—is associated with cholesterol content of erythrocyte membrane. *Clinical hemorheology and microcirculation*. 2012;51(4):243-54.

47. Lippi G, Targher G, Montagnana M, Salvagno GL, Zoppini G, Guidi GC. Relation between red blood cell distribution width and inflammatory biomarkers in a large cohort of unselected outpatients. *Archives of pathology & laboratory medicine*. 2009;133(4):628-32.
48. Veeranna V, Zalawadiya SK, Panaich SS, Ramesh K, Afonso L. The association of red cell distribution width with glycated hemoglobin among healthy adults without diabetes mellitus. *Cardiology*. 2012;122(2):129-32.
49. Lippi G, Targher G, Salvagno GL, Guidi GC. Increased red blood cell distribution width (RDW) is associated with higher glycosylated hemoglobin (HbA1c) in the elderly. *Clin Lab*. 2014;60(12):2095-8.
50. Garg N, Moorthy N, Kapoor A, Tewari S, Kumar S, Sinha A, et al. Hemoglobin A1c in nondiabetic patients: an independent predictor of coronary artery disease and its severity. *Mayo Clinic Proceedings*. 2014;89(7):908-16.
51. Kunadian V, Ford GA, Bawamia B, Qiu W, Manson JE. Vitamin D deficiency and coronary artery disease: a review of the evidence. *American heart journal*. 2014;167(3):283-91.
52. Kobayashi S, Moriya H, Aso K, Ohtake T. Vitamin E-bonded hemodialyzer improves atherosclerosis associated with a rheological improvement of circulating red blood cells. *Kidney international*. 2003;63(5):1881-7.
53. Osadnik T, Wasilewski J, Lekston A, Strzelczyk J, Kurek A, Gutowski AR, et al. Comparison of modification of diet in renal disease and chronic kidney disease epidemiology collaboration formulas in predicting long-term outcomes in patients undergoing stent implantation due to stable coronary artery disease. *Clinical Research in Cardiology*. 2014;103(7):569-76.
54. Jelkmann WE, Fandrey J, Frede S, Pagel H. Inhibition of erythropoietin production by cytokines: implications for the anemia involved in inflammatory states. *Annals of the New York Academy of Sciences*. 1994;718(1):300-11.
55. Pierce CN, Larson DF. Inflammatory cytokine inhibition of erythropoiesis in patients implanted with a mechanical circulatory assist device. *Perfusion*. 2005;20(2):83-90.
56. Yčas JW, Horrow JC, Horne BD. Persistent increase in red cell size distribution width after acute diseases: a biomarker of hypoxemia? *Clinica Chimica Acta*. 2015;448:107-17.
57. Montagnana M, Cervellin G, Meschi T, Lippi G. The role of red blood cell distribution width in cardiovascular and thrombotic disorders. *Clinical chemistry and laboratory medicine*. 2012;50(4):635-41.
58. Vogelzang M, Vlaar PJ, Svilaas T, Amo D, Nijsten MW, Zijlstra F. Computer-assisted myocardial blush quantification after percutaneous coronary angioplasty for acute myocardial infarction: a substudy from the TAPAS trial. *Eur Heart J*. 2009;30(5):594-9.
59. Nada AM. Red cell distribution width in type 2 diabetic patients. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. 2015;8:525.
60. Bilal A, Farooq JH, Kiani I, Assad S, Ghazanfar H, Ahmed I. Importance of mean red cell distribution width in hypertensive patients. *Cureus*. 2016;8(11).
61. Kurtoğlu E, Aktürk E, Korkmaz H, Sincer İ, Yılmaz M, Erdem K, et al. Elevated red blood cell distribution width in healthy smokers. *Turk Kardiyol Dern Ars*. 2013;41(3):199-206.
62. Kucera M, Balaz D, Kruzliak P, Ciccioppo R, Oravec S, Rodrigo L, et al. The effects of atorvastatin treatment on the mean platelet volume and red cell distribution width in patients with dyslipoproteinemia and comparison with plasma atherogenicity indicators—A pilot study. *Clinical biochemistry*. 2015;48(9):557-61.

63. Nagula P, Karumuri S, Otikunta AN, Yerrabandi SRV. Correlation of red blood cell distribution width with the severity of coronary artery disease—a single center study. Indian heart journal. 2017;69(6):757-61.
64. Putra BFK, Bintoro UY. Red Cell Distribution Width sebagai Prediktor Penyakit Kardiovaskular. Cermin Dunia Kedokteran. 2020;46(11):692-6.
65. Hampole CV, Mehrotra AK, Thenappan T, Gomberg-Maitland M, Shah SJ. Usefulness of red cell distribution width as a prognostic marker in pulmonary hypertension. The American journal of cardiology. 2009;104(6):868-72.
66. Xi Q, Wang Y, Liu Z, Zhao Z, Luo Q. Red cell distribution width predicts chronic thromboembolic pulmonary hypertension in patients with acute pulmonary embolism in a long-term follow-up. Clinical Chemistry and Laboratory Medicine (CCLM). 2014;52(9):e191-e5.
67. Uyarel H, Ergelen M, Cicek G, Kaya MG, Ayhan E, Turkkan C, et al. Red cell distribution width as a novel prognostic marker in patients undergoing primary angioplasty for acute myocardial infarction. Coronary artery disease. 2011;22(3):138-44.

