

## DAFTAR PUSTAKA

1. Zaenglin AL, Gruber EM, Thiboutot DM, Strauss JS. Acne variants and acneiform eruptions. In: Kang S, Amagai M, Bruckner AL, Enk AH, Margolis DJ, McMichael AJ, Orringer JS, editors. Fitzpatrick's dermatology in general medicine. 9th ed. New York: McGraw-Hill; 2019. p.1448-57.
2. Lynn DD, Umari T, Dunnick CA, Dellavalle RP. The epidemiology of acne vulgaris in late adolescence. *Adolesc Health Med Ther.* 2016;7:13–25.
3. Zaenglein AL, Thiboutot DM. Acne vulgaris. In: Bologna JL, Schaffer JV, Cerroni L, editors. *Dermatology*. 4th ed. UK: Elsevier; 2018. p. 588-603.
4. Thomas J, Kumar P, Ravi D. *The acnes*. 1st ed. Kathmandu: Jaypee Brothers Medical Publishers; 2019. p. 1-2.
5. Dinulos JGH. *Habif's clinical dermatology*. 7th ed. New York: Elsevier; 2019. p. 248.
6. Durai PCT, Nair DG. Acne vulgaris and quality of life among young adults in south India. *Indian Journal of Dermatology.* 2015;60:33-40.
7. Plewig G, Melnik B, Chen W. *Plewig and Kligman's acne and rosacea*. 4th ed. Cham. Springer Int Publishing; 2019. p. 35.
8. Tan JKL, Bhate K. A global perspective on the epidemiology of acne. *British Journal of Dermatology.* 2015;172:3-12.
9. Bagatin E, Timpano DL, Guadanhim LR, Nogueira VMA, Terzian LR, Steiner D, et al. *Acne vulgaris: Prevalence and clinical forms in adolescents from Sao Paulo, Brazil*. *An Bras Dermatol.* 2014;89(3):428–35.
10. Okoro EO, Ogunbiyi AO, George AO, Subulade MO. Association of diet with acne vulgaris among adolescents in Ibadan, southwest Nigeria. *Int J Dermatol.* 2016;55(9):982–8.
11. Sirajudin A, Tarigan S, Indria AD. Prevalensi dan gambaran epidemiologi akne vulgaris di provinsi Lampung. *JK Unila.* 2019;3(2):1.
12. Lu LY, Lai HY, Pan ZY, Wu ZX, Chen WC, Ju Q. Obese/overweight and the risk of acne vulgaris in Chinese adolescents and young adults. *HongKong J Dermatology Venereol.* 2017;25(1):5–12.

13. Alanazi T, Alajroursh W, Alharthi R, Alshalhoub M, Alshehri M. Prevalence of acne vulgaris, its contributing factors, and treatment satisfaction among the Saudi population in Riyadh, Saudi Arabia: a cross-sectional study. *J Dermatology Dermatologic Surg.* 2020;24(1):33.
14. Suppiah TSS, Sundram TKM, Tan ESS, Lee CK, Bustami NA, Tan CK. Acne vulgaris and its association with dietary intake: a Malaysian perspective. *Asia Pac J Clin Nutr.* 2018; 27(5):1141–5.
15. Kelompok studi dermatologi kosmetik Indonesia. Pedoman tata laksana akne di Indonesia. Perhimpunan Dokter Spesialis Kulit dan Kelamin Indonesia. Perpustakaan Nasional. 2015.
16. Yenny SW, Oktaviana M. Angka kepatuhan penggunaan doksisiklin pada pasien akne vulgaris derajat sedang dan berat di poliklinik kulit dan kelamin RSUP DR M Djamil Padang periode Januari 2017-Desember 2019. *MDVI. Perdoski.* 2020; 47(2):65-8.
17. Gustia R, Yenny SW, Octari S. Karakteristik penyakit kulit pada anak di poliklinik kulit dan kelamin RSUP Dr. M. Djamil Padang periode 2016-2018. *J Kedokt Syiah Kuala.* 2020;20(3):143–6.
18. Barbieri JS, Shin DB, Wang S, Margolis DJ, Takeshita J. Association of race/ethnicity and sex with differences in health care use and treatment for acne. *JAMA Dermatology.* 2020;156(3):312–9.
19. Surekha A, Sharan TR, Keerth T, Snehaseela V, Kumari JK, TLee CK, Bustami NA, Tan CK. Acne vulgaris and its association with dietary intake: a Malaysian perspective. *Asia Pac J Clin Nutr.* 2018; 27(5):1141–5.
20. Mahto A. Acne vulgaris. *Medicine.* 2017;45(6):386–9. doi:10.1016/j.mpmed.2017.03.003.
21. Bhuiyan MNI. Quality of life in patients with acne-a questionnaire study. *EAS Journal of Parasitology and Infectious Disease.* 2021;29(4):194-8.
22. Hazarika N, Rajaprabha RK. Assessment of life quality index among patients with acne vulgaris in a suburban population. *Indian J Dermatol.* 2016;61(2):163-8.
23. Bergler-Czop B, Brzezińska-Wcisło L. Pro-inflammatory cytokines in patients with various kinds of acne treated with isotretinoin. *Postep Derm*

- Alergol. 2014; 31: 21-8.
- 24. Degitz K, Ochsendorf F. Acne. JDDG. 2017;15(7):709–722.
  - 25. Szyszkowska B, Łepecka-Klusek C, Kozłowicz K, et al. The influence of selected ingredients of dietary supplements on skin condition. Postep Derm Alergol 2014; 31: 174-81.
  - 26. Mahmood SN, Bowe WP. Diet and acne update: carbohydrates emerge as the main culprit. J Drugs Dermatol. 2014;13:428-35.
  - 27. Kowalska H, Sysa-Jędrzejowska A, Woźniacka A. Role of diet in the aetiopathogenesis of acne. Przegl Dermatol. 2018;105(1):51–62.
  - 28. Youssef EMK, Youssef MKE. Diet and acne in upper Egypt. American Journal of Dermatology and Venereology. 2014;3(1): 13-22.
  - 29. Mirdamadi Y, Thielitz A, Wiede A, et al. Insulin and insulin-like growth factor-1 can modulate the phosphoinositide-3-kinase/Akt/FoxO1 pathway in SZ95 sebocytes in vitro. Mol Cell Endocrinol. 2015;415:32–44.
  - 30. Aghasi M, Golzarand M, Shab-Bidar S, Aminianfar A, Omidian M, Taheri F. Dairy intake and acne development: a meta-analysis of observational studies. Clin Nutr. 2018;38(3):1067–75.
  - 31. Kim H, Moon SY, Sohn MY, Lee WJ. Insulin-like growth factor-1 increases the expression of inflammatory biomarkers and sebum production in cultured sebocytes. Ann Dermatol. 2017;29(1):20–5.
  - 32. Melnik BC. Acne vulgaris: The metabolic syndrome of the pilosebaceous follicle. J Clin Dermatol. 2018;36(1):29–40.
  - 33. LaRosa CL, Quach KA, Koons K, et al. Consumption of dairy in teenagers with and without acne. J Am Acad Dermatol. 2016;75(2):318–322.
  - 34. Ulvestad M, Bjertness E, Dalgard F, Halvorsen JA. Acne and dairy products in adolescence: results from a Norwegian longitudinal study. J Eur Acad Dermatol Venereol. 2017;31(3):530–5.
  - 35. Juhl CR, Bergholdt HKM, Miller IM, Jemec GBE, Kanters JK, Ellervik C. Lactase persistence, milk intake, and adult acne: a mendelian randomization study of 20,416 Danish adults. Nutrients. 2018;10(8):1041.
  - 36. (NCBI) NC for BI. Review. Mesh database. Published 2014. Accessed October 2, 2021. <https://www.ncbi.nlm.nih.gov/mesh/68008892>

37. Mosby. Mosby's medical dictionary. 11st ed. St Louis: Elsevier; 2021. p. 22.
38. (NCBI) NC for BI. Review. Mesh database. Published 2015. Accessed October 2, 2021. <https://www.ncbi.nlm.nih.gov/mesh/68016454>
39. Vandenbroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al. Strengthening the reporting of observational studies in epidemiology (STROBE): Explanation and elaboration. *Int J Surg*. 2014;12(12):1500–24.
40. Burris J, Rietkerk W, Woolf K. Relationships of self reported dietary factors and perceived acne severity in a cohort of New York young adults. *J Acad Nutr Diet*. 2014;114:384-92.
41. Tsoy NO. Effect of milk and dairy products upon severity of acne for young people. *World Applied Sciences Journal*. 2016;24(3): 403-7.
42. Alghamdi H, Alhemel AM, Bukhari IA. Effect of milk and dairy products consumption on acne risk and severity in young adult patients with Acne vulgaris attending the dermatology clinics at King Fahd Hospital of the University in Alkhobar, Saudi Arabia. *Sci Med Sci*. 2017;0(4 (12)):4–8.
43. Penso L, Touvier M, Deschamps M, Szabo De Edelenyi F, Hercberg S, Ezzedine K, et al. Association between adult acne and dietary behaviors: Findings from the NutriNet-Santé Prospective Cohort Study. *JAMA Dermatology*. 2020;156(8):854–62.
44. Aalemi AK, Anwar I, Chen H. Dairy consumption and acne: a case control study in Kabul, Afghanistan. *Clin Cosmet Investig Dermatol*. 2019;12:481–7.
45. Al Suhaibawi AJK, Taher TMJ. Milk consumption among patients with acne vulgaris. *EurAsian Journal of BioSciences*. 2020;14:3935-8.
46. Say YH, Heng AHS, Reginald K, Wong YR, Teh KF, Rawanan Shah SM, et al. Modifiable and non-modifiable epidemiological risk factors for acne, acne severity and acne scarring among Malaysian Chinese: a cross-sectional study. *BMC Public Health*. 2021;21(1):1-12.