

DAFTAR PUSTAKA

1. World Health Organization. Naming the coronavirus disease (COVID-19) and the virus that causes it. 2020.
2. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497–506.
3. Kementrian Kesehatan Republik Indonesia. Pedoman pencegahan dan pengendalian corona virus disease (COVID-19) revisi ke-5. Kemenkes RI. Jakarta; 2020.
4. Menteri Kesehatan. Peta Sebaran. Covid19.go.id. 2022.
5. Cahyani C, Novida H, Soelistijo SA, Hadi U, Siagian N. Correlation between Neutrophil-to-Lymphocyte Ratio with Disease Severity in Diabetic Patients with COVID-19 at Tertiary Referral Hospital in Indonesia. *J Hunan Univ Sci*. 2021;48(9).
6. Sintoro DK, Sintoro F, Artanti D. Hubungan antara rasio neutrofil limfosit dengan derajat klinis COVID-19 pada pasien anak di RSUD Tarakan provinsi Kalimantan Utara. *Intisari Sains Medis*. 2021;12(2):449–52.
7. Pal R, Bhadada SK. COVID-19 and diabetes mellitus: An unholy interaction of two pandemics. *Diabetes Metab Syndr Clin Res Rev*. 2020;14(4):513–517.
8. International Diabetes Federation. Diabetes around the world in 2021. 2021.
9. Pahlevi R. Jumlah Penderita Diabetes Indonesia Terbesar Kelima di Dunia. *Databoks*. 2021.
10. Taher N, Huda MSB, Chowdhury TA. COVID-19 and diabetes: What have we learned so far? *Clin Med J R Coll Physicians London*. 2020;20(4):E87–90.
11. Guo W, Li M, Dong Y, Zhou H, Zhang Z, Tian C, et al. Diabetes is a risk factor

for the progression and prognosis of COVID-19. *Diabetes Metab Res Rev* [Internet]. 2020;36. Available from: <https://doi.org/10.1002/dmrr.3319>

12. Liu J, Liu Y, Xiang P, Pu L, Xiong H, Li C, et al. Neutrophil-to-lymphocyte ratio predicts critical illness patients with 2019 coronavirus disease in the early stage. *J Transl Med*. 2020;18(206).
13. Liu Y, Du X, Chen J, Jin Y, Peng L, Wang HHX, et al. Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with COVID-19. *J Infect*. 2020;81(1):e6–12.
14. Yang A, Liu J, Tao W, Li H. The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. *Int Immunopharmacol*. 2020;84.
15. Kong M, Zhang H, Cao X, Mao X, Lu Z. Higher level of Neutrophil-to-Lymphocyte is associated with severe COVID-19. *Epidemiol Infect*. 2020;148(e139).
16. Hall JE, Guyton AC. *Guyton dan Hall Buku Ajar Fisiologi Kedokteran*. 12th ed. Vol. 12, Saunders. 2011. 423–433 p.
17. Sherwood L. *Introduction to Human Physiology*. 8th ed. Cengage Learning. 2013. 419–424 p.
18. Amanda DA. Rasio Neutrofil-Limfosit pada Covid-19; Sebuah tinjauan literatur. *Wellness Heal Mag*. 2020;2(2):219–23.
19. Wiargitha IK. Hubungan Rasio Neutrofil Limfosit Dalam Memprediksi Mortalitas Pada Pasien Peritonitis Yang Dioperasi Di Rsup Sanglah Denpasar. *J Udayana Med*. 2017;
20. Imran MM, Ahmed U, Usman U, Ali M, Shaukat A, Gul N. Neutrophil / Lymphocyte Ratio – A Marker of COVID-19 Pneumonia Severity. *Int J Clin Pract*. 2021;75(4).

21. de Jager CPC, Wever PC, Gemen EFA, Kusters R, van Gageldonk-Lafeber AB, van der Poll T, et al. The Neutrophil-Lymphocyte Count Ratio in Patients with Community-Acquired Pneumonia. *PLoS One*. 2012;7(10).
22. Susilo A, Rumende CM, Pitoyo CW, Santoso WD, Yulianti M, Herikurniawan H, et al. Coronavirus Disease 2019: Tinjauan Literatur Terkini. *J Penyakit Dalam Indones*. 2020;7(1):45–67.
23. WHO. WHO coronavirus disease (COVID-19) dashboard. World Health Organization. 2022.
24. Yuliana Y. Corona virus diseases (Covid-19): Sebuah tinjauan literatur. *Wellness Heal Mag*. 2020;2(1):187–92.
25. Levani Y, Prastya AD, Mawaddatunnadila S. Coronavirus Disease 2019 (COVID-19): Patogenesis, Manifestasi Klinis dan Pilihan Terapi. *J Kedokt dan Kesehat*. 2021;17(1):44–57.
26. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727–33.
27. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. *Clin Immunol*. 2020;215.
28. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020;8:e21.
29. Li G, Fan Y, Lai Y, Han T, Li Z, Zhou P, et al. Coronavirus infections and immune responses. *J Med Virol*. 2020;92:424–32.
30. Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*. 2020;



31. Li X, Geng M, Peng Y, Meng L, Lu S. Molecular immune pathogenesis and diagnosis of COVID-19. *J Pharm Anal.* 2020;10:102–8.
32. Mardewi IGA, Yustiani NT. Gambaran hasil laboratorium pasien COVID-19 di RSUD Bali Mandara: sebuah studi pendahuluan. *Intisari Sains Medis.* 2021;12(1):374–8.
33. PDPI, PERKI, PAPDI, PERDATIN, IDAI. Pedoman Tatalaksana COVID-19 Edisi 3. 3rd ed. Pedoman Tatalaksana COVID-19. 2020. 36–37 p.
34. Kemenkes RI. Manajemen Klinis Tata Laksana Corona Virus Disease 2019 (COVID-19) di Fasilitas Pelayanan Kesehatan. 2021;6–25.
35. Mus R, Thaslifa T, Abbas M, Sunaidi Y. Studi Literatur: Tinjauan Pemeriksaan Laboratorium pada Pasien COVID-19. *J Kesehat Vokasional.* 2021;5(4):246–7.
36. Decroli E. DIABETES MELITUS TIPE 2. Kam A, Efendi YP, Decroli GP, Rahmadi A, editors. Padang: Pusat Penerbitan Bagian Ilmu Penyakit Dalam; 2019.
37. Fatimah RN. DIABETES MELITUS TIPE 2. *J Major.* 2015;4(5):93–101.
38. Soelistijo S, Suastika K, Lindarto D, Decroli E, Permana H, Sucipto K, et al. Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2021. PB. PERKENI; 2021.
39. Kementerian Kesehatan Republik Indonesia. Tetap Produktif, Cegah Dan Atasi Diabetes Mellitus. Pusat Data Dan Informasi Kementerian Kesehatan RI. 2020.
40. Utomo AA, R A, Rahmah S, Amalia R. Faktor Risiko Diabetes Mellitus Tipe 2 : A Systemic Review. *J Kaji dan Pengemb Kesehat Masy.* 2020;01(01):44–52.
41. Trisnawati SK, Setyorogo S. Faktor Risiko Kejadian Diabetes Melitus Tipe II Di Puskesmas Kecamatan Cengkareng Jakarta Barat Tahun 2012. *J Ilm Kesehat.* 2013;5(1):6–11.

42. Lou M, Luo P, Tang R, Peng Y, Yu S, Huang W, et al. Relationship between neutrophil-lymphocyte ratio and insulin resistance in newly diagnosed type 2 diabetes mellitus patients. *BMC Endocr Disord.* 2015;15(9).
43. Azab B, Camacho-Rivera M, Taioli E. Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects. *PLoS One.* 2014;9(11).
44. Damayanti R. Gambaran nilai Neutrophil-Lymphocyte Ratio (NLR) pada Pasien COVID-19 di RS Islam Siti Khadijah Palembang Tahun 2020. Repos Poltekkes Kemenkes Palembang. 2020;
45. Dyusupova A, Faizova R, Yurkovskaya O, Belyaeva T, Terekhova T, Khismetova A, et al. Clinical characteristics and risk factors for disease severity and mortality of COVID-19 patients with diabetes mellitus in Kazakhstan: A nationwide study. *Heliyon.* 2021;7(3).
46. Zhang Y, Li H, Zhang J, Cao Y, Zhao X, Yu N, et al. The clinical characteristics and outcomes of patients with diabetes and secondary hyperglycaemia with coronavirus disease 2019: A single-centre, retrospective, observational study in Wuhan. *Diabetes, Obes Metab.* 2020;22(8):1443–54.
47. Wu J, Huang J, Zhu G, Wang Q, Lv Q, Huang Y, et al. Elevation of blood glucose level predicts worse outcomes in hospitalized patients with COVID-19: A retrospective cohort study. *BMJ Open Diabetes Res Care.* 2020;8.