

## DAFTAR PUSTAKA

- Ali, M. E., Parikh, S. and Pearson, J. P. (2017) 'Total Trypsin and Active Trypsin : could they be used as a Biomarker for Duodenal Reflux ?', *Global Journal of Otolaryngology*, 3(3), pp. 1-6.
- Ali, M. S. et al. (2013) 'Bile acids in laryngopharyngeal refluxate: Will they enhance or attenuate the action of pepsin?', *Laryngoscope*, 123(2), pp. 434-439. doi: 10.1002/lary.23619.
- Andersson, O. (2009) *Laryngopharyngeal reflux development and refinement of diagnostic tools*. Göteborg: Intellecta DocuSys.
- Asaoka, D. et al. (2014) 'Current perspectives on reflux laryngitis', *Clinical journal of Gastroenterology*, 7, pp. 471-5.
- Bagian Telinga Hidung Tenggorok Bedah Kepala dan Leher Dr.RSUP. M. Djamil Padang. (no date) *Laporan Rekapitulasi Pasien Subbagian Laring Faring*. padang.
- Belafsky, P. C., Postma, G. N. and Koufman, J. A. (2001) 'The validity and reliability of the reflux finding score (RFS)', *The Laryngoscope*, 111(8), pp. 1313-1317.
- Belafsky, P. C., Postma, G. N. and Koufman, J. A. (2002) 'Validity and reliability of the reflux symptom index (RSI)', *Journal of Voice*, 16(2), pp. 274-277.
- Birtić, D. et al. (2012) 'Significance of the Pepsin from the Saliva in the Diagnosis and Treatment of Laryngopharyngeal Reflux Disease', *Collegium antropologicum*, 36(2), pp. 83-6.
- Bobin, F., Journe, F. and Lechien, J. R. (2020) 'Saliva pepsin level of laryngopharyngeal reflux patients is not correlated with reflux episodes', *Laryngoscope*, 130(5), pp. 1278-1281. doi: 10.1002/lary.28260.
- Calvo-Henríquez, C. et al. (2017) 'Is Pepsin a Reliable Marker of Laryngopharyngeal Reflux? A Systematic Review', *Otolaryngology - Head and Neck Surgery (United States)*, 157(3), pp. 385-391. doi: 10.1177/0194599817709430.
- Campagnolo, A. M. et al. (2014) 'Laryngopharyngeal Reflux: Diagnosis, Treatment, and Latest Research', *Int Arch Otorhinolaryngol*, 18, pp. 184-91.
- Campbell, R. et al. (2017) 'The Role of Helicobacter pylori in Laryngopharyngeal Reflux.', *Otolaryngology--head and neck surgery : official journal of American Academy of Otolaryngology-Head and Neck Surgery*, 156(2), pp. 255-262.
- Carrau, R. L. et al. (2004) 'The Impact of Laryngopharyngeal Reflux on Patient-Reported Quality of Life', *The Laryngoscope*, 114, pp. 670-674.
- Chiba, T. (2017) 'Laryngopharyngeal reflux disease ( LPRD ) - Review article Author';, *Medical Research Archives*, 5(2), pp. 1-14.
- De Corso, E. et al. (2021) 'Impact of bile acids on the severity of laryngo-pharyngeal reflux', *Clinical Otolaryngology*, 46(1), pp. 189-195. doi: 10.1111/coa.13643.
- Dettmar, P. W. et al. (2011) 'Review article: reflux and its consequences - the laryngeal, pulmonary and oesophageal manifestations', *Alimentary Pharmacology & Therapeutics*, 33, pp. 1-71.
- Divakaran, S. et al. (2021) 'Laryngopharyngeal reflux: Symptoms, signs, and presence of pepsin in saliva - A reliable diagnostic triad', *International Archives of Otorhinolaryngology*, 25(2), pp. E273-E278. doi: 10.1055/s-0040-1709987.
- Du, X. et al. (2017) 'The diagnostic value of pepsin detection in saliva for gastro-esophageal reflux disease: A preliminary study from China', *BMC Gastroenterology*, 17(1), pp. 1-9. doi: 10.1186/s12876-017-0667-9.
- Falk, G. L. and Vivian, S. J. (2016) 'Laryngopharyngeal reflux: diagnosis, treatment and latest research', *European Surgery - Acta Chirurgica Austriaca*, 48(2), pp. 74-91. doi: 10.1007/s10353-016-0385-5.
- Fortunato, J. E., D'Agostino Jr, R. B. and Lively, M. O. (2016) 'Pepsin in saliva as a biomarker for oropharyngeal reflux compared with 24- • hour esophageal impedance / pH monitoring in pediatric patients', *Neurogastroenterol Motil*, pp. 1-10.
- Fritz, M. A. et al. (2016) 'The Accuracy of the Laryngopharyngeal Reflux Diagnosis: Utility of the Stroboscopic Exam', *Otolaryngology - Head and Neck Surgery*, pp. 1-6.
- Galli, J. et al. (2006) 'Laryngeal carcinoma and laryngo-pharyngeal reflux disease.', *Acta otorhinolaryngologica Italica : organo ufficiale della Società italiana di*

- otorinolaringologia e chirurgia cervico-facciale*, 26(5), pp. 260–263.
- Gill, G. A. et al. (2005) 'Laryngeal Epithelial Defenses Against Laryngopharyngeal Reflux : Investigations of E-cadherin , Carbonic Anhydrase Isoenzyme III , and Pepsin', *Annals of Otology, Rhinology and Laryngology*, 114(12), pp. 913–21.
- Grabowski, M. et al. (2011) 'Pepsin and bile acids in induced sputum of chronic cough patients', *Respiratory Medicine*, 105(8), pp. 1257–1261.
- Hoppo, T. et al. (2012) 'How Much Pharyngeal Exposure Is " Normal " ? Normative Data for Laryngopharyngeal Reflux Events Using Hypopharyngeal Multichannel Intraluminal Impedance ( HMII )', *Journal of Gastrointestinal Surgery*, 16, pp. 16–25.
- Iannella, G. et al. (2015) 'International Journal of Pediatric Otorhinolaryngology Investigation of pepsin in tears of children with laryngopharyngeal reflux disease', *International Journal of Pediatric Otorhinolaryngology*, pp. 1–4.
- Johnston, N. et al. (2004) 'Pepsin and Carbonic Anhydrase Isoenzyme III as Diagnostic Markers for Laryngopharyngeal Reflux Disease', *The Laryngoscope*, 114, pp. 2129–34.
- Johnston, N. (2011) 'Review article: Uptake of pepsin at pH 7 - In non-acid reflux - Causes inflammatory, and perhaps even neoplastic, changes in the laryngopharynx', *Alimentary Pharmacology and Therapeutics*, pp. 13–20.
- Junaid, M. et al. (2020) 'Laryngopharyngeal Reflux Disease: Outcome of Patients After Treatment in Otolaryngology Clinics', *Cureus*, 12(12), pp. 1–9. doi: 10.7759/cureus.12195.
- Jung, A. R. et al. (2019) 'Association Between Pepsin in the Saliva and the Subjective Symptoms in Patients With Laryngopharyngeal Reflux', *Journal of Voice*, 33(2), pp. 150–154. doi: 10.1016/j.jvoice.2017.10.015.
- Kabir, S. (2004) 'Detection of Helicobacter pylori DNA in Feces and Saliva by Polymerase Chain Reaction: A Review', *Helicobacter*, 9(2), pp. 115–23.
- Kahrilas, P. J. (2003) 'GERD pathogenesis, pathophysiology, and clinical manifestations', *Cleveland Clinic Journal of Medicine*.
- Kamani, T. et al. (2012) 'The prevalence of laryngopharyngeal reflux in the English population', *European Archives of Oto-Rhino-Laryngology*, 269(10), pp. 2219–2225. doi: 10.1007/s00405-012-2028-1.
- Kariya, S., Okano, M. and Nishizaki, K. (2014) 'An association between Helicobacter pylori and upper respiratory tract disease: Fact or fiction?', *World Journal of Gastroenterology*, 20(6), pp. 1470–1484. doi: 10.3748/wjg.v20.i6.1470.
- Karkos, P. D. and Belafsky, P. C. (2011) 'Review article: Nonsurgical treatment of non-acid reflux', *Alimentary Pharmacology and Therapeutics*, pp. 66–70.
- Kesari, S., Chakraborty, S. and Sharma, B. (2017) 'Evaluation of Risk Factors for Laryngopharyngeal Reflux among Sikkimese Population', *Kathmandu University Medical Journal*, 15(1), pp. 29–34.
- Kilic, M. et al. (2013) 'International Journal of Pediatric Otorhinolaryngology Impact of laryngopharyngeal and gastroesophageal reflux on asthma control in children', *International Journal of Pediatric Otorhinolaryngology*, 77(3), pp. 341–345.
- Kim, J. H. et al. (2016) 'Extra-Esophageal Pepsin from Stomach Refluxate Promoted Tonsil Hypertrophy', *PLOS ONE*, pp. 1–16.
- Knight, J. et al. (2005) 'Sensitive pepsin immunoassay for detection of laryngopharyngeal reflux', *Laryngoscope*, 115(8), pp. 1473–1478.
- Krishnaswamy, R. T. et al. (2012) 'Salivary IgG assay to detect Helicobacter pylori infection in an Indian adult population', *Indian Journal of Dental Research*, 23(5), pp. 1–5.
- Kuo, C.-L. (2020) 'Acid Reflux Increases Susceptibility to Laryngopharyngeal and Oropharyngeal Cancers', *Archives of Otorhinolaryngology-Head & Neck Surgery*, 4(1), pp. 3–5. doi: 10.24983/scitemed.aohns.2020.00123.
- Kurtaran, H. et al. (2008) 'Role of Helicobacter Pylori in pathogenesis of upper respiratory system diseases', *Journal of the National Medical Association*, 100(10), pp. 1224–30.
- Li, Y. et al. (2021) 'Effects of acids, pepsin, bile acids, and trypsin on laryngopharyngeal reflux diseases: physiopathology and therapeutic targets', *European Archives of Oto-Rhino-Laryngology*, 279(6), pp. 2743–2752. doi: 10.1007/s00405-021-07201-w.
- Li, Y. W. et al. (2017) 'Relationship between salivary pepsin concentration and esophageal mucosal integrity in patients with gastroesophageal reflux disease', *Journal of*

- Neurogastroenterology and Motility*, 23(4), pp. 517–525. doi: 10.5056/jnm16178.
- Lipan, M. J., Reidenberg, J. S. and Laitman, J. T. (2006) 'Anatomy of reflux: A growing health problem affecting structures of the head and neck', *Anatomical Record - Part B New Anatomist*, pp. 261–70.
- Marshall, B., Howat, A. J. and Wright, P. A. (1999) 'Oral fluid antibody detection in the diagnosis of Helicobacter pylori infection', *J. Med. Microbiol.*, 48(11), pp. 1043–6.
- Massawe, W. A. et al. (2021) 'Laryngopharyngeal reflux disease, prevalence and clinical characteristics in ENT department of a tertiary hospital Tanzania', *World Journal of Otorhinolaryngology - Head and Neck Surgery*, 7(1), pp. 28–33. doi: 10.1016/j.wjorl.2020.04.009.
- Neeharika, M. (2017) 'Detection of Helicobacter pylori from Saliva using Enzyme-linked Immunosorbent Assay Detection of Helicobacter pylori from Saliva using Enzyme-linked Immunosorbent Assay', *Oral and Maxillofacial Pathology journal*, 8(2), pp. 85–7.
- Nunes, H. S. et al. (2016) 'Comparison between the Reflux Finding Score and the Reflux Symptom Index in the Practice of Otorhinolaryngology', *International Archives of Otorhinolaryngology*, 20(3), pp. 218–221. doi: 10.1055/s-0036-1579557.
- Patel, D. and Vaezi, M. F. (2013) 'Normal Esophageal Physiology and Laryngopharyngeal Reflux', *Otolaryngologic Clinics of NA*, 46(6), pp. 1023–41.
- Pearson, J. P. and Parikh, S. (2011a) 'Alimentary Pharmacology and Therapeutics Review article : reflux and its consequences – the laryngeal , pulmonary and oesophageal manifestations Conference held in conjunction with the 9th International Symposium Review article : nature and properties of', *Alimentary Pharmacology & Therapeutics*, 33(1), pp. 21–3.
- Pearson, J. P. and Parikh, S. (2011b) 'Review article: Nature and properties of gastro-oesophageal and extra-oesophageal refluxate', *Alimentary Pharmacology and Therapeutics*, 33(1), pp. 2–7.
- Pedersen, M. (2012) 'Otolaryngology Laryngopharyngeal Reflux – A Randomized Clinical Controlled Trial', *otolaryngology*, 1(4), pp. 1–5.
- Rahman, L. et al. (2020) 'DNA detection of Helicobacter pylori in saliva of patients with low salivary pH', *Zanco Journal of Medical Sciences*, 24(2), pp. 283–290. doi: 10.15218/zjms.2020.032.
- Rasmussen, L. T. et al. (2012) 'Detection of Helicobacter pylori in gastric biopsies, saliva and dental plaques of dyspeptic patients from Marília, São Paulo, Brazil: Presence of vacA and cagA genes', *Journal of Venomous Animals and Toxins Including Tropical Diseases*, 18(2), pp. 180–187. doi: 10.1590/S1678-91992012000200008.
- Rees, C. J. and Belafsky, P. C. (2008) 'Laryngopharyngeal reflux: Current concepts in pathophysiology, diagnosis, and treatment', *International Journal of Speech-Language Pathology*, 10(4), pp. 245–53.
- Ren, J. et al. (2017) 'PepsinA as a Marker of Laryngopharyngeal Reflux Detected in Chronic Rhinosinusitis Patients', *Laryngology and Neurolaryngology*, 156(5), pp. 893–900.
- Rimbara, E., Sasatsu, M. and Graham, D. Y. (2013) 'PCR detection of helicobacter pylori in clinical samples', *Methods in Molecular Biology*, 943, pp. 279–287. doi: 10.1007/978-1-60327-353-4\_19.
- Ryu, I. Y. et al. (2017) 'Comparison of Methods for Collecting Saliva for Pepsin Detection in Patients with Laryngopharyngeal Reflux', *Korean Journal of Otorhinolaryngology-Head Neck Surgery*, 60(11), pp. 570–4.
- Sabry, M. et al. (2022) 'Fasting salivary pepsin level as a reliable non-invasive method of screening for laryngopharyngeal reflux in Egyptian patients', *The Egyptian Journal of Internal Medicine*. doi: 10.1186/s43162-022-00100-4.
- Saruç, M. et al. (2012) 'Risk factors for laryngopharyngeal reflux', *European Archives of Oto-Rhino-Laryngology*, 269(4), pp. 1189–94.
- Sastroasmoro, S. (2011) 'Perkiraan Besar Sampel dalam Penelitian Klinis', *Dasar-dasar Metodologi Penelitian*, p. 359.
- Sereg-Bahar, M. et al. (2015a) 'Pepsin and bile acids in saliva in patients with laryngopharyngeal reflux - a prospective comparative study', *Clinical Otolaryngology*, 40(3), pp. 234–239.
- Sereg-Bahar, M. et al. (2015b) 'Pepsin and bile acids in saliva in patients with

- laryngopharyngeal reflux - a prospective comparative study', *Clinical Otolaryngology*, 40(3), pp. 234–239. doi: 10.1111/coa.12358.
- Sereg-Bahar, M., Jerin, A. and Hocevar-Boltezar, I. (2015) 'Higher levels of total pepsin and bile acids in the saliva as a possible risk factor for early laryngeal cancer', *Radiology and Oncology*, 49(1), pp. 59–64.
- Shilpa, C. et al. (2019) 'Laryngopharyngeal Reflux and GERD: Correlation Between Reflux Symptom Index and Reflux Finding Score', *Indian Journal of Otolaryngology and Head and Neck Surgery*, 71(s1), pp. 684–688. doi: 10.1007/s12070-018-1480-7.
- Silva, Á. S. (2020) 'Evaluation of the Re flux Symptom Index and the Endolaryngeal Findings Scale after Treatment in Individuals with Laryngopharyngeal Re flux'.
- Siupsinskiene, N. et al. (2017) 'Helicobacter pylori in the tonsillar tissue : a possible association with chronic tonsillitis and laryngopharyngeal reflux', *The Journal of Laryngology & Otology*, 1(1984), pp. 1–8.
- Siupsinskiene, N. et al. (2017) 'Helicobacter pylori in the tonsillar tissue: A possible association with chronic tonsillitis and laryngopharyngeal reflux', *Journal of Laryngology and Otology*, 131(6), pp. 549–556. doi: 10.1017/S0022215117000597.
- Spiridoulias, A. et al. (2015) 'Detecting laryngopharyngeal reflux in patients with upper airways symptoms: Symptoms, signs or salivary pepsin?', *Respiratory Medicine*, 109(8), pp. 963–969. doi: 10.1016/j.rmed.2015.05.019.
- Swenson, E. R. (1998) 'Carbonic anhydrase inhibitors and ventilation: A complex interplay of stimulation and suppression', *European Respiratory Journal*, 12(6), pp. 1242–1247. doi: 10.1183/09031936.98.12061242.
- Tack, J. (2006) 'Review article: the role of bile and pepsin in the pathophysiology and treatment of gastro-oesophageal reflux disease.', *Alimentary pharmacology & therapeutics*, 24(2), pp. 10–16.
- Wang, J. et al. (2018) 'Pepsin in saliva as a diagnostic biomarker in laryngopharyngeal reflux: a meta-analysis', *European Archives of Oto-Rhino-Laryngology*, 275(3), pp. 671–678. doi: 10.1007/s00405-017-4845-8.
- Wang, L. et al. (2010) 'Correlation of pepsin-measured laryngopharyngeal reflux disease with symptoms and signs', *Otolaryngology-Head and Neck Surgery*, 143(6), pp. 765–71.
- Wood, J. M. et al. (2011) 'Biomarkers and laryngopharyngeal reflux', *The Journal of Laryngology & Otology*, 125, pp. 1218–24.
- Yilmaz, T. et al. (2014) 'Laryngopharyngeal reflux and Helicobacter pylori', *World Journal of Gastroenterology*, 20(27), pp. 8964–70.