

## DAFTAR PUSTAKA

1. Gao Y, Li T, Han M, Li X, Wu D, Xu Y, *et al.* Diagnostic utility of clinical laboratory data determinations for patients with the severe COVID-19. *J Med Virol.* 2020;92(7):791-6.
2. Shiehzadegan S, Alaghemand N, Fox M, Venketaraman V. Analysis of the delta variant B.1.617.2 COVID-19. *Clin Pract.* 2021;11(4):778-784.
3. World Health Organization (2022). Weekly epidemiological update on COVID-19 - 15 March 2022. <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---15-march-2022> - Diakses 16 Maret 2022.
4. World Health Organization (2020). Indonesia coronavirus (COVID-19) statistic. <https://covid19.who.int/region/searo/country/id> - Diakses pada 16 Maret 2022.
5. Pemerintah Provinsi Sumatera Barat (2022). Data pantauan COVID-19 Provinsi Sumatera Barat. <https://corona.sumbarprov.go.id/> - Diakses pada 16 Maret 2022.
6. Dinas Komunikasi dan Informatika Pemerintah Kota Padang (2022). Situasi dan perkembangan COVID-19 Kota Padang. <http://corona.padang.go.id/> - Diakses pada 16 Maret 2022.
7. Wang Y, Wang Y, Chen Y, Qin Y. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *J Med Virol.* 2020;92(6):568-76.
8. Kementerian Kesehatan Republik Indonesia. Pedoman pencegahan dan pengendalian coronavirus disease (COVID-19). 5th ed. Jakarta: Kementerian Kesehatan RI; 2020.
9. Bohn MK, Lippi G, Horvath A, Sethi S, Koch D, Ferrari M, *et al.* Molecular, serological, and biochemical diagnosis and monitoring of COVID-19: IFCC taskforce evaluation of the latest evidence. *Clin Chem Lab. Med.* 2020;58(7):1037-52.
10. Pozdnyakova O, Connell NT, Battinelli EM, Connors JM, Fell G, Kim AS. Clinical significance of CBC and WBC morphology in the diagnosis and clinical course of COVID-19 infection. *Am J Clin Pathol.* 2021;155(3):364-375.
11. Liu J, Li S, Liu J, Liang B, Wang X, Wang H, *et al.* Longitudinal characteristics of lymphocyte responses and cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *EBioMedicine.* 2020;55:102763.
12. Chan AS, Rout A. Use of neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios in COVID-19. *J Clin Med.* 2020;12(7):448.
13. Tatum D, Taghavi S, Houghton A, Stover J, Toraih E, Duchesne J. Neutrophil-to-lymphocyte ratio and outcomes in Louisiana COVID-19 patients. *Shock (Augusta, Ga).* 2020;54(5):652-8.
14. Shang W, Dong J, Ren Y, Tian M, Li W, Hu J, *et al.* The value of clinical parameters in predicting the severity of COVID-19. *J Med Virol.* 2020;92(10):2188-92.

15. Simadibrata DM, Pandhita BAW, Ananta ME, Tango T. Platelet-to-lymphocyte ratio, a novel biomarker to predict the severity of COVID-19 patients: A systematic review and meta-analysis. 2020;23(1):20-6.
16. Qu R, Ling Y, Zhang Y, Wei L, Chen X, Li X, *et al.* Platelet-to-lymphocyte ratio is associated with prognosis in patients with coronavirus disease-19. *J Med Virol.* 2020;92(9):1533-1541.
17. Lee JH, Song S, Yoon SY, Lim CS, Song JW, Kim HS. Neutrophil to lymphocyte ratio and platelet to lymphocyte ratio as diagnostic markers for pneumonia severity. *Br J Biomed Sci.* 2016;73(3):140-2.
18. Harrison AG, Lin T, Wang P. Mechanisms of SARS-CoV-2 Transmission and Pathogenesis. *Trends Immunol.* 2020;41(12):1100-15.
19. Mina A, Besien K, Plataniias LC. Hematological manifestations of COVID-19. *Leuk Lymphoma.* 2020;61(12):2790-8.
20. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. *J Adv Res.* 2020;24:91.
21. Naqvi AAT, Fatima K, Mohammad T, Fatima U, Singh IK, Singh A, *et al.* Insights into SARS-CoV-2 genome, structure, evolution, pathogenesis and therapies: Structural genomics approach. *Biochim Biophys Acta Mol Basis Dis.* 2020;1866(10).
22. Fehr AR, Perlman S. Coronaviruses: an overview of their replication and pathogenesis. *Methods Mol Biol.* 2015;1282:1-23.
23. Kirtipal N, Bharadwaj S, Kang SG. From SARS to SARS-CoV-2, insights on structure, pathogenicity and immunity aspects of pandemic human coronaviruses. *Infect Genet Evol.* 2020;85.
24. Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): A Review. *JAMA.* 2020;324(8):782-93.
25. Bulut C, Kato Y. Epidemiology of COVID-19. *Turk J Med Sci.* 2020;50(SI-1):563-70.
26. Wang L, He W, Yu X, Hu D, Bao M, Liu H, *et al.* Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J Infect.* 2020;80(6):639.
27. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, *et al.* Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. *JAMA.* 2020;323(11):1061-1069.
28. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, *et al.* Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020;395(10223):507-13.
29. Li X, Geng M, Peng Y, Meng L, Lu S. Molecular immune pathogenesis and diagnosis of COVID-19. *J Pharm Anal.* 2020;10(2):102-8.
30. de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into emerging coronaviruses. *Nat Rev Microbiol* 2016;14(8):523-34.
31. Cevik M, Kuppalli K, Kindrachuk J, Peiris M. Virology, transmission, and pathogenesis of SARS-CoV-2. *BMJ.* 2020;371.

32. Li G, Fan Y, Lai Y, Han T, Li Z, Zhou P, *et al.* Coronavirus infections and immune responses. *J Med Virol.* 2020;92(4):424-32.
33. Wang Y, Ju M, Chen C, Yang D, Hou D, Tang X, *et al.* Neutrophil-to-lymphocyte ratio as a prognostic marker in acute respiratory distress syndrome patients: a retrospective study. *J Thorac Dis.* 2018;10(1):273-82.
34. Fan YY, Huang ZT, Li L, Wu MH, Yu T, Koup RA, *et al.* Characterization of SARS-CoV-specific memory T cells from recovered individuals 4 years after infection. *Arch Virol.* 2009;154(7):1093-9.
35. Nalbant A, Kaya T, Varim C, Yaylaci S, Tamer A, Cinemre H. Can the neutrophil/lymphocyte ratio (NLR) have a role in the diagnosis of coronavirus 2019 disease (COVID-19)? *Rev Assoc Med Bras (1992).* 2020;66(6):746-51.
36. Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med.* 2020;382(18):1708-20.
37. PDPI, PERKI, PAPDI, PERDATIN, IDAI. Pedoman tatalaksana COVID-19. 4th ed. PPDI; 2022.
38. Morfi CW, Junaidi A, Elsesmita, Asrini DN, Lestari DM, Medison I, *et al.* Kajian terkini coronavirus disease 2019 (COVID-19). *Jurnal Ilmu Kesehatan Indonesia.* 2020;1(1).
39. Yavuz S, Ünal S. Antiviral treatment of COVID-19. *Turk J Med Sci.* 2020;50(SI-1):611-9.
40. Gibson PH, Croal BL, Cuthbertson BH, Small GR, Ifezulike AI, Gibson G, *et al.* Preoperative neutrophil-lymphocyte ratio and outcome from coronary artery bypass grafting. *Am Heart J.* 2007;154(5):995-1002.
41. Yoon NB, Son C, Um SJ. Role of the neutrophil-lymphocyte count ratio in the differential diagnosis between pulmonary tuberculosis and bacterial community-acquired pneumonia. *Ann Lab Med.* 2013;33(2):105-10.
42. Liu Y, Du X, Chen J, Jin Y, Peng L, Wang HHX, *et al.* Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with COVID-19. *J Infect.* 2020;81(1):e6.
43. Forget P, Khalifa C, Defour JP, Latinne D, van Pel MC, de Kock M. What is the normal value of the neutrophil-to-lymphocyte ratio?. *BMC Res Notes.* 2017;10(1):1-4.
44. Eid MM, Al-Kaisy M, Regeia WAL, Khan HJ. The prognostic accuracy of neutrophil-lymphocyte ratio in covid-19 patients. *Front Emerg Med.* 2021;5(1):0-5.
45. Bedel C, Korkut M. Neutrophil-to-lymphocyte ratio and COVID-19. *Shock (Augusta, Ga).* 2021;56(5):874.
46. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, *et al.* Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis.* 2020;71(15):762-8.
47. Song H, Kim HJ, Park KN, Kim SH, Oh SH, Youn CS. Neutrophil to lymphocyte ratio is associated with in-hospital mortality in older adults admitted to the emergency department. *Am J Emerg Med.* 2021;40:133-7.
48. Bonow RO, Fonarow GC, O'Gara PT, Yancy CW. Association of coronavirus disease 2019 (COVID-19) with myocardial injury and mortality. *JAMA Cardiol.* 2020;5(7):751-3.

49. Gasparyan AY, Ayvazyan L, Mukanova U, Yessirkepov M, Kitas GD. The platelet-to-lymphocyte ratio as an inflammatory marker in rheumatic diseases. *Ann Lab Med.* 2019;39(4):345-57.
50. Luo H, He L, Zhang G, Yu J, Chen Y, Yin H, *et al.* Normal reference intervals of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio, and systemic immune inflammation index in healthy adults: a large multi-center study from Western China. *Clin Lab.* 2019;65(3):255-65.
51. Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LF. The trinity of COVID-19: immunity, inflammation and intervention. *Nat Rev Immunol.* 2020 Jun;20(6):363-74.
52. Henry BM, de Oliveira MHS, Benoit S, Plebani M, Lippi G. Hematologic, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): A meta-analysis. *Clin Chem Lab Med.* 2020;58(7):1021-8.
53. Sarkar S, Kannan S, Khanna P, Singh AK. Role of platelet-to-lymphocyte count ratio (PLR), as a prognostic indicator in COVID-19: A systematic review and meta-analysis. *J Med Virol.* 2022;94(1):211-21.
54. Tan L, Wang Q, Zhang D, Ding J, Huang Q, Tang YQ, *et al.* Lymphopenia predicts disease severity of COVID-19: a descriptive and predictive study. *Signal Transduct Target Ther.* 2020;5(1):1-3.
55. Akboga MK, Canpolat U, Yuksel M, Yayla C, Yilmaz S, Turak O, Ozeke O, Topaloglu S, Aras D. Platelet to lymphocyte ratio as a novel indicator of inflammation is correlated with the severity of metabolic syndrome: A single center large-scale study. *Platelets.* 2016;27(2):178-83.
56. Kartal O, Kartal AT. Value of neutrophil to lymphocyte and platelet to lymphocyte ratios in pneumonia. *Bratisl Lek Listy.* 2017;118(9):513-6.
57. Lwanga SK, Lemeshow S, World Health Organization. Sample size determination in health studies: a practical manual. World Health Organization; 1991.
58. Vermonte Philips TYW. Karakter dan persebaran covid-19 di indonesia. *CSIS Commentaries.* 2020;1:1-12.
59. Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, *et al.* Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA.*
60. Gennaro FD, Pizzol D, Marotta C, Antunes M, Racalbuto V, Veronese N, *et al.* Coronavirus Diseases (COVID-19) Current Status and Future Perspectives: A Narrative Review. *Int. J. Environ. Res. Public Health.* 2020;17(8):2690.
61. Zhang J, Dong X, Cao Y, Yuan Y, Yang Y, Yan Y, *et al.* Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy.* 2020;75(7):1730-41.
62. Schurz H, Salie M, Tromp, Goal EG, Kinnear CJ, Moller M. The X chromosome and sex-specific effects in infectious disease susceptibility. *Hum Genomics.* 2019;13(1):2.
63. Jin JM, Bai P, He W, Wu F, Liu XF, Han DM, *et al.* Gender Differences in Patients With COVID-19: Focus on Severity and Mortality. *Front Public Health.* 2020;8:152.

64. Zhao K, Li R, Wu X, Zhao Y, Wang T, Zheng Z, *et al.* Clinical features in 52 patients with COVID-19 who have increased leukocyte count: a retrospective analysis. *Eur J Clin Microbiol.* 2022;39(12):2279–87.
65. Dawood Q, Al-Hashim Z, al Hijaj BA, Jaber R, Khalaf A. Study of hematological parameters in patients with coronavirus disease 2019 in Basra. *Iraqi Journal of Hematology.* 2020;9(2):160.
66. Kong M, Zhang H, Cao X, Mao X, Lu Z. Higher level of neutrophil-to-lymphocyte is associated with severe COVID-19. *Epidemiol Infect.* 2020;148:e139.
67. Erdogan A, Can FE, Gönüllü H. Evaluation of the prognostic role of NLR, LMR, PLR, and LCR ratio in COVID-19 patients. *J Med Virol.* 2021;93(9):5555–9.
68. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, *et al.* Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol Pract.* 2020;146(1):110-8.
69. Channappanavar R, Perlman S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Springer Berlin Heidelberg.* 2017;39(5):529-39.
70. Yang AP, Liu JP, Tao WQ, Li HM. . The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. *Int Immunopharmacol.* 2020;84:106504.
71. Cho KI, Ann SH, Singh GB, Her AY, Shin ES. Combined Usefulness of the Platelet-to-Lymphocyte Ratio and the Neutrophil-to-Lymphocyte Ratio in Predicting the Long-Term Adverse Events in Patients Who Have Undergone Percutaneous Coronary Intervention with a Drug-Eluting Stent. *PLoS One.* 2015; 10(7):e0133934.
72. Balta S, Ozturk C. The platelet-lymphocyte ratio: A simple, inexpensive and rapid prognostic marker for cardiovascular events. *Platelets.* 2015;26(7):680-1.

