

## DAFTAR PUSTAKA

- Abdou HM. Dissolution, Bioavailability & Bioequivalence 1 Edition. London : Mack Publishing Company;1989
- Adachi H, Ioppolo F, Paoloni M, Santilli V. Physical characteristics, pharmacological properties and clinical efficacy of the ketoprofen patch: a new patch formulation. *Eur Rev Med Pharmacol Sci.* 2011;15(7):823-30.
- Agoes G. Sistem Penghantaran Obat Pelepasan Terkendali. Bandung: ITB Press; 2008.
- Alatas F, Sutarna TH, Pratama MR, et al. Pembentukan ko-amorf irbesartan-L-arginin dan dampaknya terhadap kelarutan dan laju disolusi irbesartan. *Riset Informasi Kesehatan.* 2020;9(2):98-106
- Allesø, N. Chieng, S. Rehder, J. Rantanen, T. Rades, J. Aaltonen, Enhanced dissolution rate and synchronized release of drugs in binary systems through formulation: Amorphous naproxen–cimetidine mixtures prepared by mechanical activation. *Journal of Controlled Release.* 2009;136:45-53.
- Altman R, Bosch B, Brune K, Patrignani P, Young C. Advances in NSAID development: evolution of diclofenac products using pharmaceutical technology. *Drugs.* 2015;75(8):859-877.
- Ansel, H. C. Pengantar Bentuk Sediaan Farmasi. (Edisi IV). Jakarta: Universitas Indonesia Press;1989
- Arakawa T, Kita Y, Koyama AH. Solubility enhancement of gluten and organic compounds by arginine. *Int J Pharm.* 2008;355(1-2):220-223.
- Arana V, Paz Y, González A, Méndez V, Méndez JD. Healing of diabetic foot ulcers in L-arginine-treated patients. *Biomed Pharmacother.* 2004;58(10):588-597.
- Babu NJ, Nangia A. Solubility advantage of amorphous drugs and pharmaceutical cocrystals. *Crystal Growth and Design.* 2011;11:2662–79
- Badawy SI, Hussain MA. Microenvironmental pH modulation in solid dosage forms. *J Pharm Sci.* 2007;96(5):948-959.
- Banakar UV. Pharmaceutical Dissolution Testing, Bioavailability & Bioequivalence : Science, Applications, and Beyond. New York : John Wiley & Sons; 1991
- Barikah, K. Z.Traditional and Novel Methods for Cocrystal Formation: A MiniReview. *Systematic Reviews in Pharmacy.* 2018; 9(1):79–82.

Berry DJ, Steed JW. Pharmaceutical cocrystals, salts and multicomponent systems; intermolecular interactions and property based design. *Adv Drug Deliv Rev.* 2017;117:3-24.

Beyer A, Radi L, Grohganz H, Löbmann K, Rades T, Leopold CS. Preparation and recrystallization behavior of spray-dried co-amorphous naproxen-indomethacin. *Eur J Pharm Biopharm.* 2016;104:72-81.

Blagden N, de Matas M, Gavan PT, York P. Crystal engineering of active pharmaceutical ingredients to improve solubility and dissolution rates. *Adv Drug Deliv Rev.* 2007;59(7):617-630.

Birmingham B & Buvanendran A. Practical management of pain 5<sup>th</sup> edition. Amsterdam : Elsevier. 2014

Bode-Böger SM, Böger RH, Alfke H, et al. L-arginine induces nitric oxide-dependent vasodilation in patients with critical limb ischemia. A randomized, controlled study. *Circulation.* 1996;93(1):85-90.

Chandra Gupta P, Sharma N, et al. Cocrystals: an Alternative Approach To Modify Physicochemical Properties of Drugs. *Ijpcbs.* 2015;5(2):427–436.

Christopher Vimalson D, Palimalakrisnan S, et al. Techniques to enhance solubility of hydrophobic drugs: An overview. *Asian Journal of Pharmaceutics.* 2016;10(2):67-75.

Collett, E. Polarized Light: Fundamentals and Applications. New York, USA: Marcel Dekker; 2009

Cruz-Cabeza, AJ. Acid-base crystalline complexes and the pKa rule. *CrystEngComm.* 2012;14(20):6362–6365.

Dengale SJ, Grohganz H, Löbmann K, et al. Recent advances in co-amorphous drug formulations. *Advanced Drug Delivery Reviews.* 2016

Departemen Kesehatan Republik Indonesia. Farmakope Indonesia Edisi V. Jakarta: Direktorat Jenderal Pengawasan Obat dan Makanan; 2014

Desiraju, G. R. (1991) “Hydration in organic crystals: Prediction from molecular structure. *Journal of the Chemical Society, Chemical Communications.* 1991;6:426–428.

Elena I, Kratochvil, B. Pharmaceutical cocrystals: Progress in Chemistry. *Elsevier Ltd.* 2010;22(5):829–836.

Gaikwad A, Tamizhrasi S, Sorti A, et al. Formulation In Vitro Characterization of Polymethacrylic Acid Nanoparticle Containing Furosemide. *Int J. Pharm Res.* 2010;2(1): 300-304.

Gardner, CR, Almarsson N, et al. Application of high throughput technologies to drug substance and drug product development: Computers and Chemical Engineering. 2004;28:943–953.

Ganong, WF. Fisiologi Saraf dan Sel Otot. Jakarta: EGC;2003

Geesala R, Bar N, Dhoke NR, et al. Porous Polymer Scaffold for On-Site Delivery of Stem Cells – Protects from Oxidative Stress and Potentiates Wound Tissue Repair. *Biomaterials*. 2015;77:1-13

Guyton, AC, Hall, JE. Buku Ajar Fisiologi Kedokteran.Jakarta: EGC;1997

Guillory, J. K. Handbook of Pharmaceutical Salts: Properties, Selection, and Use. USA: Journal of Medicinal Chemistry;2003

Hancock BC & Zografi G. Characteristics and Significance of the Amorphous State in Pharmaceutical Systems. *J Pharm Sci*. 1997;86(1):1– 12

He G, Chow PS, Tan, RH. Predicting multicomponent crystal formation: The interplay between homomeric and heteromeric interactions. *Crystal Growth and Design*. 2009;9(10):4529–4532.

Hussain K, Qamar A, Bukhari NI, et al. Impact of Particle-Size Reduction on the Solubility and Antidiabetic Activity of Extracts of Leaves of Vinca rosea. *Turk J Pharm Sci*. 2019;16(30):335-339

Hwang Sung-Joo, Park H, et al. Preparation and characteritation of glimepiride eutectic mixture with l- arginine for improvement of dissolution rate. *International Pharmaceutical Journal*. 2020;581:119288

Khankari RK., Law D, Grant DJ. Determination of water content in pharmaceutical hydrates by differential scanning calorimetry. *International Journal of Pharmaceutics*. 1992;82(1-2):117–127.

Jeffres GA, Saenger W. The role of hydrogen bonding in structure and function of the nucleic acids. In: *hydrogen bonding in biological structure*. 1996

Jensen KT, Larsen FH, Löbmann K, Rades T, Grohganz H. Influenceof variation in molar ratio on co-amorphous drug-amino acid systems. *Eur.J.Pharm. Biopharm*. 2016;107:32–39.

Jensen K, Löbmann T, Rades H, Grohganz. Improving co-amorphous drug formulations by the addition of the highly water soluble amino acid, proline. *Pharmaceutics*. 2014;6:416–435.

Lachman, L. et al. Pharmaceutical Dosage Forms : Disperse System. Florida, USA: CRCPress;1994

Lee T, Wang YW. Initial salt screening procedures for manufacturing ibuprofen. *Drug Dev. Ind. Pharm*. 2009;35:555–567.

Lenz E, Jensen KT, Blaabjerg LI, Knop, et al. Solid-state properties and dissolution behaviour of tablets containing co-amorphous indomethacin–arginine. *Eur. J. Pharm. Biopharm.* 2015;96:44–52.

Liu J, Grohganz H, Löbmann K, Rades T, & Hempel N. Co-Amorphous Drug Formulations in Numbers: Recent Advances in Co-Amorphous Drug Formulations with Focus on Co-Formability, Molar Ratio, Preparation Methods, Physical Stability, In Vitro and New Formulation Strategies. *Pharmacetics*. 2021;13(3):389

Löbmann K, Strachan C, Grohganz H, Rades T, Korhonen O, Laitinen R. Co-amorphous simvastatin and glipizide combinations show improved physical stability without evidence of intermolecular interactions. *European Journal of Pharmaceutics and Biopharmaceutics*. 2012;81:159–169.

Löbmann K, Grohganz H, Laitinen R., et al. Amino acids as co-amorphous stabilizers for poorly water soluble drugs—Part 1: preparation, stability and dissolution enhancement. *Eur. J. Pharm. Biopharm.* 2013;85:873–881.

Luangtana-anan M, Luangtana-anan M, Nunthanid, et al. Potential of different salt forming agents on the formation of chitosan nanoparticles as carriers for protein drug delivery systems. *J. Pharm. Investig.* 2009;49:37–44.

Luo S, Lutkenhaus JL, Nasrabadi H. Use of differential scanning calorimetry to study phase behavior of hydrocarbon mixtures in nano-scaleporous media. *Journal of Petroleum Science and Engineering*. 2018;163:731–738.

Maithani M, Richa R, Vertika G, Kumar D. Development and validation of a RP-HPLC method for the determination of chlorpheniramine maleate and phenylephrine in pharmaceutical dosage form. 2010

Malviya, R. et al. High performance liquid chromatography: A short review. USA: Journal of Global Pharma Technology; 2010

Martin, A., J. S. dan A. C. Farmasi Fisik Edisi 3 Jilid II. Jakarta: Universitas Indonesia Press; 1993

Meor Mohd Affandi MM, Tripathy M, Shah SA, Majeed AB. Solubility enhancement of simvastatin by arginine: thermodynamics, solute-solvent interactions, and spectral analysis. *Drug Des Devel Ther.* 2016;10:959–969.

Merimee TJ, Rabinovitz D, Riggs L, Burgess JA, Rimoin DL, Cokusick VA. Plasma growth hormone after arginine injection. *N Engl J Med.* 1967;276:434–9.

Miles L, Hall J, Jenner B, Addis R, Hutchings S. Predicting rapid analgesic onset of ibuprofen salts compared with ibu-profen acid: Tlag, Tlow, Tmed, and a novel parameter. *TCmaxRef. Curr. Med. Res. Opin.* 2018;34:1483–1490.

Moore MD dan Wildfong PLD. Aqueous solubility enhancement through engineering of binary solid composites: Pharmaceutical applications. *Journal of Pharmaceutical Innovation*. 2009;4(1):36–49.

Mura P, Bettinetti GP, Cirri M, Maestrelli F, Sorrenti M, Catenacci L. Solid state characterization and dissolution properties of naproxen–arginine–hydroxypropyl- $\beta$ -cyclodextrin ternary system. *Eur. J. Pharm. Biopharm.* 2005;59:99–106.

Mura P, Maestrelli F, Cirri M. Ternary systems of naproxen with hydroxypropyl  $\beta$ -cyclodextrin and aminoacids. *Int. J. Pharm.* 2003;260:293–302

Murakami T & Takano M. Intestinal efflux transporters and drug absorption. *Expert Opinion on Drug Metabolism & Toxicology*. 2008;4(7):923-939

Mutschler E. Dinamika Obat Buku Ajar Farmakologi dan Toksikologi. Edisi Kelima. Bandung: ITB Press;1991.

Mycek JM. Farmakologi Ulasan Bergambar. Jakarta: Widya Medika;2001. Ojarinta R., Heikkinen AT, Sievänen E, Laitinen R.. Dissolution behavior of co-amorphous amino acid-indomethacin mixtures: The ability of amino acids to stabilize the supersaturated state of indomethacin. *Eur.J. Pharm. Biopharm.* 2017;112:85–95.

Ojarinta R, Lerminiaux L, Laitinen R. Spray drying of poorly soluble drugs from aqueous arginine solution. *Int. J. Pharm.* 2017;532:89–298.

Patel D, Bertz R, Ren S, et al. A systematic review of gastric acid-reducing agent-mediated drug-drug interaction with orally administrated medications. *Clinical Pharmacokinetics*. 2020;59:447-462

Patole T dan Deshpande A. Co-Crystallization-a Technique for Solubility Enhancement. *International Journal of Pharmaceutical Sciences and Research*. 2014;5(9):3566.

Palazzo B, Walsh D, Iafisco M, et al. Amino acid synergistic effect on structure, morphology and surface properties of biomimetic apatite nanocrystals. *Acta Biomater.* 2009;5:1241–1252.

Polli J, Rekhi GS, Ausgburger LL, Shah VP. Methods to Compare Dissolution Profiles and a Rationale for Wide Dissolution Specifications for Metoprolol Tartrate Tablets. *J Pharm Sci.* 1997;86:690-700.

Pouton, CW. Formulation of poorly water-soluble drugs for oral administration: Physicochemical and physiological issues and the lipid formulation classification system. *European Journal of Pharmaceutical Sciences*. 2006;29(3-4):278–287.

- Pralhad T dan Rajendrakumar K. Study of freeze-dried quercetin- cyclodextrin binary systems by DSC, FT-IR, X-ray diffraction and SEM analysis. *Journal of Pharmaceutical and Biomedical Analysis*. 2004;34(2):333–339.
- Qi X, Zhang J, Wang W, Cao D. Solubility and stability of indomethacin in arginine-assisted solubilization system. *Pharm. Dev. Technol.* 2013;18:852–855.
- Reimer L. Scanning Electron Microscopy: Physics of Image Formation and Microanalysis, Second Edition. *Measurement Science and Technology*. 200;11(12):1826
- Rodrigues M, et al. Pharmaceutical cocrystallization techniques. Advances and challenges. *International Journal of Pharmaceutics*. 2014;547(1–2):404–420.
- Roni MA, dan Dipu MH. Dissolution Enhancement of Poorly Soluble Carbamazepine By Using. *International Journal of Pharmaceutical Sciences And Research*. 2011;2(1):49–57.
- Rukmono. Kumpulan kuliah patologi. Jakarta : Bagian patologi anatomik FK UI;2000.
- Sandri G, Bonferoni MC, Ferrari F, et al. The role of particle size in drug release and absorption. *Particulate Products*. 2014;19:323-341
- Serajuddin AT. Salt formation to improve drug solubility. *Adv. Drug Deliv.* 2007;59 :603–616.
- Shargel L, Wu-Pong S, dan Yu ABC. Applied Biopharmaceutics and Pharmacokinetics, Fifth Edition. Surabaya : Unair Press;2004
- Shinkai N, et al. Intra-articular penetration of ketoprofen and analgesic effects after topical patch application in rats. *Journal of Controlled Release*. 2007;131(2):107-112
- Sinko PJ dan Singh Y. Martin's physical pharmacy and pharmaceutical sciences: Physical chemical and biopharmaceutical principles in the pharmaceutical sciences: Sixth edition. USA: Lippincott William & Willian;2011
- SJ Dengale, H Grohganz, T Rades, K Lobmann. Recent advances in co-amorphous drug formulations. *Adv. Drug Deliv. Rev.* 2016;100:116–125
- Skineh J, Sathisaran I, Dalvi SV, Rohani S. Co-amorphous Form of Curcumin-Folic Acid Dihydrate with Increased Dissolution Rate. *Crystal Growth & Design*. 2017;17(2):101-109
- Smith B T. Remington Education: Physical Pharmacy. London: Pharmaceutical Press;2015

- Sun DD, Wen H, Taylos LS. Non-sink dissolution conditions for predicting product quality and in vivo performance of supersaturating drug delivery system. *J Pharm Sci.* 2016;105(9):2377-2388
- Swarbrick, J. Encyclopedia of Pharmaceutical Technology. London: Taylor and Francis;2013
- Sweetman SC. Martindale: The Complete Drug Reference. Thirty-sixth Edition. London: Pharmaceutical Press;2009
- Tavintharan S, Woon K, Dong X, Lim S. Niacin results in reduced monocyte adhesion in patients with type 2 diabetes mellitus. *Atherosclerosis Journal.* 2011;215(1):176-179
- Tjandrawinata RR, Hendra P, Djunarko I, et al . Anti-inflammation effect of bioactive fraction DLBS0533 containing phaleria macrocarpa and *Nigella sativa* on animal model. *Journal of Pharmacy and Pharmaceutical Science.* 2015;7(1):408-411
- Tjay TH dan Rahardja K. Obat-Obat Penting. Edisi Keenam. Jakarta: PT. Elex Media Komputindo;2007
- Trask A V, et al. Screening for crystalline salts via mechanochemistry. *Chemical Communications.* 2006;1:51–53.
- Tsume Y, et al. The Biopharmaceutics Classification System: Subclasses for in vivo predictive dissolution (IPD) methodology and IVIVC. *European Journal of Pharmaceutical Sciences.* 2014;57(1):152–163.
- Vogel G. Drug Discovery and Evaluation Pharmacological Assays Second Edition : Human & Experimental Toxicology. London: Springer;2008
- Vueba J, Carvalho V, Veiga SJ, et al. Influence of Cellulose Ether Polymers on Ketoprofen Release from Hydrophilic Matrix Tablets. *International Journal of Pharmaceutical.* 2004;58:51-59.
- Wagner H, Spahn-Langguth A, Hanafy A, Koggel P, Langguth. Intestinal drug efflux: formulation and food effects. *Advanced Drug Delivery Reviews.* 2001;13-31
- Ward RM, Kern SE, Lugo RA. Pharmacokinetics, Pharmacodynamics, and Pharmacogenetics. Amsterdam : Elsevier. 2012
- West A. Solid State Chemistry and its Applications, 2nd Edition, Student Edition. USA: JOHN WILEY & SONS;2014
- Wicaksono Y, Setyawan D, dan Siswadono. Formation of ketoprofen-malonic acid cocrystal by solvent evaporation method. *Indonesian Journal of Chemistry,* 2017;17(2):161–166.

Wu G, Morris SM. Arginine metabolism: nitric oxide and beyond.  
*Biochem J.* 1998;336:1–17

Thakur, T. S. dan Thakuria, R. Crystalline multi-component solids , an alternative for addressing the hygroscopicity issue in pharmaceutical materials  
Crystalline multi-component solids , an alternative for addressing the hygroscopicity issue in pharmaceutical materials. *CrystalGrowth & Design.* 2020;20(9):6245-6265

Tomaszewska I, Karki S, Shur J, Price R., & Fotaki N. Pharmaceutical characterisation and evaluation of cocrystals: Importance of in vitro dissolution conditions and type of coformer. *International Journal of Pharmaceutics.* 2013;53(2):380–388.

Yamamura H, Gotoh Y, Sakamoto Y. Physicochemical properties of amorphous precipitates of cimetidine–indomethacin binary system. *European Journal of Pharmaceutics and Biopharmaceutics.* 2000;49:259-265.

Zaini, E. et al. Cocrystalline phase transformation of binary mixture of trimethoprim and sulfamethoxazole by slurry technique. *Asian Journal of Pharmaceutical and Clinical Research.* 2010;3(4):26–29.

Zaini E, Fitriani L, Firdaus WA, Sidadang W, Rosaini H, Putra OD, Oyama H, Uekusa H. Improved Solubility and Dissolution Rate of Ketoprofen by the Formation of Multicomponent Crystals with Tromethamine. *Crystals.* 2022;12: 275-290

