

DAFTAR PUSTAKA

1. Mebis L, Berghe G Van Den. The hypothalamus-pituitary-thyroid axis in critical illness. *Neth J Med.* 2009;67:332–40.
2. Robertson LC, Al-Haddad M. Recognizing the critically ill patient. *Anaesth Intensive Care Med.* 2013;14:11–4.
3. Kalzén H, Larsson B, Eksborg S, Lindberg L, Edberg KE, Frostell C. Survival after PICU admission: The impact of multiple admissions and complex chronic conditions. Kou YR, editor. *PLoS One.* 2018;13:e0193294.
4. Buckley S, Nyman A, Fine-Goulden M. Two Decades of Decreasing PICU Mortality : Better care or a changing patient group? *Pediatr Crit Care Med.* 2021;22:12–3.
5. Burns JP, Sellers DE, Meyer EC, Lewis-Newby M, Truog RD. Epidemiology of Death in the PICU at Five U.S. Teaching Hospitals*. *Crit Care Med.* 2014;42:2101–8.
6. Rismala D, Fatimatuzzuhroh. Profil Pasien Sakit Kritis yang Dirawat di Pediatric Intensive Care Unit Rumah Sakit Cipto Mangunkusumo berdasar Sistem Skoring Pediatric Logistic Organ Dysfunction-2. *Sari Pediatr.* 2019;21:37–43.
7. Hajari S, Endang S, Annisa M, Hary N. Gambaran mortalitas pasien di ruang PICU RSUD Abdul Wahab Sjahranie Samarinda periode 2016-2017. *J Mutiara Mahakam.* 2019;7:41–57.
8. Rachmawati N, Ihsan I. Hubungan status vitamin D dengan mortalitas dan lama rawatan pada anak sakit kritis. *Maj Kedokt Andalas.* 2017;40:82.
9. El-Ella SSA, El-Mekkawy MS, El-Dihemey MA. Prevalence and prognostic value of non-thyroidal illness syndrome among critically ill children. *An Pediatría (English Ed.* 2019;90:237–43.
10. Boonen E, Van Den Berghe G. Endocrine responses to critical illness: Novel insights and therapeutic implications. *J Clin Endocrinol Metab.* 2014;99:1569–82.
11. Gardner DG, Shoback DM, Greenspan FS (Francis S. Greenspan's Basic & Clinical Endocrinology. McGraw-Hill Medical; 2011. 880 p.
12. Bikle DD. The free hormone hypothesis: when, why, and how to measure the free hormone levels to assess vitamin d, thyroid, sex hormone, and cortisol status. *JBMR Plus.* 2021;5.
13. Barrett KE, Heddwen B, Scott B, Susan B. Ganong's Review of Medical Physiology. 23rd ed. McGraw-Hill Medical; 2010.
14. Tanurahardja AG, Pudjiadi AH, Dwipoerwantoro PG, Pulungan A. Thyroid hormone profile and PELOD score in children with sepsis. *Paediatr Indones.* 2014;54:245.

15. Destariani CP, Lubis M, Deliana M, Yanni GN. Triiodothyronine levels and mortality in children with sepsis and septic shock. *Paediatr Indones*. 2018;58:20.
16. Sayarifard F, Yaghmaie B, Sayarifard A, Mohammadpour M, Sharifzadeh M, Abbasi F, et al. Thyroid function tests in critically ill children; any correlation with disease severity or outcome? *Iran J Pediatr*. 2018;28:1–5.
17. Plikat K, Langgartner J, Buettner R, Bollheimer LC, Woenckhaus U, Schölmerich J, et al. Frequency and outcome of patients with nonthyroidal illness syndrome in a medical intensive care unit. *Metabolism*. 2007;56:239–44.
18. Rady H. Profile of patients admitted to pediatric intensive care unit, Cairo University Hospital: 1-year study. *Ain-Shams J Anaesthesiol*. 2014;7:500.
19. Surbatovic M, Vojvodic D, Khan W. Immune response in critically ill patients. *Mediators Inflamm*. 2018;2018:10–3.
20. Pfortmueller CA, Meisel C, Fux M, Schefold JC. Assessment of immune organ dysfunction in critical illness: utility of innate immune response markers. *Intensive Care Med Exp*. 2017;5:49.
21. Kliegman R, Stanton B, St. Geme JW, Schor NF, Behrman RE, Preceded by: Nelson WE (Waldo E. Nelson Textbook of Pediatrics. 21st ed. Philadelphia: Elsevier; 2020.
22. Klein I, Danzi S. Thyroid disease and the heart. *Curr Probl Cardiol*. 2016;41:65–92.
23. Qari FA. Thyroid function status and its impact on clinical outcome in patients admitted to critical care. *Pakistan J Med Sci*. 2015;31:915–9.
24. Marks SD. Nonthyroidal illness syndrome in children. *Endocrine*. 2009;36:355–67.
25. Warner MH, Beckett GJ. Mechanisms behind the non-thyroidal illness syndrome: An update. *J Endocrinol*. 2010;205:1–13.
26. Peeters RP, Wouters PJ, Van Toor H, Kaptein E, Visser TJ, Van Den Berghe G. Serum 3,3',5'-triiodothyronine (rT3) and 3,5,3'-triiodothyronine/rT3 are prognostic markers in critically ill patients and are associated with postmortem tissue deiodinase activities. *J Clin Endocrinol Metab*. 2005;90:4559–65.
27. Nagaya T, Fujieda M, Otsuka G, Yang JP, Okamoto T, Seo H. A potential role of activated NF-κB in the pathogenesis of euthyroid sick syndrome. *J Clin Invest*. 2000;106:393–402.
28. Pappa T, Alevizaki M. Non-Thyroidal Illness. In: Vitti P, Hegedus L, editors. *Thyroid Diseases*. Springer International Publishing; 2017. p. 1–25. (Endocrinology).
29. Yu J, Koenig RJ. Induction of type 1 iodothyronine deiodinase to prevent the nonthyroidal illness syndrome in mice. *Endocrinology*. 2006;147:3580–5.

30. Jawa RS, Anillo S, Huntoon K, Baumann H, Kulaylat M. Interleukin-6 in surgery, trauma, and critical care part II: Clinical implications. *J Intensive Care Med.* 2011;26:73–87.
31. Wajner SM, Goemann IM, Bueno AL, Larsen PR, Maia AL. IL-6 promotes nonthyroidal illness syndrome by blocking thyroxine activation while promoting thyroid hormone inactivation in human cells. *J Clin Invest.* 2011;121:1834–45.
32. Tiwari L, Singhi S, Jayashree M, Baranwal A, Bansal A. Hypoalbuminemia in critically sick children. *Indian J Crit Care Med.* 2014;18:565–9.
33. Nicholson JP, Wolmarans MR, Park GR, Farman J, Care I, Hospital A, et al. The role of albumin in critical illness. *Br J Anaesth.* 2000;85:599–610.
34. Dickson PW, Howlett GJ, Schreiber G. Metabolism of prealbumin in rats and changes induced by acute inflammation. *Eur J Biochem.* 1982;129:289–93.
35. Den Brinker M, Joosten KFM, Visser TJ, Hop WCJ, De Rijke YB, Hazelzet JA, et al. Euthyroid sick syndrome in meningococcal sepsis: The impact of peripheral thyroid hormone metabolism and binding proteins. *J Clin Endocrinol Metab.* 2005;90:5613–20.
36. de Vries EM, Fliers E, Boelen A. The molecular basis of the non-thyroidal illness syndrome. *J Endocrinol.* 2015;225:R67–81.
37. Sayre NL, Lechleiter JD. Fatty acid metabolism and thyroid hormones. *Curr trends Endocrinol.* 2012;6:65–76.
38. de Vries EM, Fliers E, Boelen A. The molecular basis of the non-thyroidal illness syndrome. *J Endocrinol.* 2015;225:R67–81.
39. Haugen BR. Drugs that suppress TSH or cause central hypothyroidism. *Best Pract Res Clin Endocrinol Metab.* 2009;23:793–800.
40. Mebis L, Debaveye Y, Ellger B, Derde S, Ververs EJ, Langouche L, et al. Changes in the central component of the hypothalamus-pituitary-thyroid axis in a rabbit model of prolonged critical illness. *Crit Care.* 2009;13:1–10.
41. Fekete C, Gereben B, Doleschall M, Harney JW, Dora JM, Bianco AC, et al. Lipopolysaccharide induces type 2 iodothyronine deiodinase in the mediobasal hypothalamus: Implications for the nonthyroidal illness syndrome. *Endocrinology.* 2004;145:1649–55.
42. Bartalena L, Martino E, Brandi LS, Falcone M, Pacchiarotti A, Ricci C, et al. Lack of nocturnal serum thyrotropin surge after surgery. *J Clin Endocrinol Metab.* 1990;70:293–6.
43. Romijn JA, Wiersinga WM. Decreased nocturnal surge of thyrotropin in nonthyroidal illness. *J Clin Endocrinol Metab.* 1990;70:35–42.

44. Langouche L, Vander Perre S, Frystyk J, Flyvbjerg A, Hansen TK, Van den Berghe G. Adiponectin, retinol-binding protein 4, and leptin in protracted critical illness of pulmonary origin. *Crit Care*. 2009;13:1–6.
45. Chen K, Li F, Li J, Cai H, Strom S, Bisello A, et al. Induction of leptin resistance through direct interaction of C-reactive protein with leptin. *Nat Med*. 2006;12:425–32.
46. Koch A, Weiskirchen R, Zimmermann HW, Sanson E, Trautwein C, Tacke F. Relevance of serum leptin and leptin-receptor concentrations in critically ill patients. *Mediators Inflamm*. 2010;20:1–9.
47. Forceville X, Vitoux D, Gauzit R, Combes A, Lahilaire P, Chappuis P. Selenium, systemic immune response syndrome, sepsis, and outcome in critically ill patients. *Crit Care Med*. 1998;26:1536–44.
48. de Oliveira Iglesias SB, Leite HP, Paes AT, de Oliveira S V., Sarni ROS. Low plasma selenium concentrations in critically ill children: The interaction effect between inflammation and selenium deficiency. *Crit Care*. 2014;18:1–8.
49. Vincent JL, Forceville X. Critically elucidating the role of selenium. *Curr Opin Anaesthesiol*. 2008;21:148–54.
50. Renko K, Hofmann PJ, Stoedter M, Hollenbach B, Behrends T, Köhrle J, et al. Down-regulation of the hepatic selenoprotein biosynthesis machinery impairs selenium metabolism during the acute phase response in mice. *Faseb J*. 2009;23:1758–65.
51. Schomburg L. Selenium, selenoproteins and the thyroid gland: Interactions in health and disease. *Nat Rev Endocrinol*. 2012;8:160–71.
52. Moura Neto A, Zantut-Wittmann DE. Abnormalities of thyroid hormone metabolism during systemic illness: The low T₃ syndrome in different clinical settings. *Int J Endocrinol*. 2016;2016:1–9.
53. Economidou F, Douka E, Tzanela M, Nanas S, Kotanidou A. Thyroid function during critical illness. *Hormones*. 2011;10:117–24.
54. Brent G a, Hershman JM. Thyroxine therapy in patients with severe concentration. *J Clin Endocrinol Metab*. 2014;63:1–8.
55. Acker CG, Singh AR, Flick RP, Bernardini J, Greenberg A, Johnson JP. A trial of thyroxine in acute renal failure. *Kidney Int*. 2000;57:293–8.
56. Van Den Berghe G, Wouters P, Weekers F, Mohan S, Baxter RC, Veldhuis JD, et al. Reactivation of pituitary hormone release and metabolic improvement by infusion of growth hormone-releasing peptide and thyrotropin-releasing hormone in patients with protracted critical illness. *J Clin Endocrinol Metab*. 1999;84:1311–23.
57. Portman MA, Slee A, Olson AK, Cohen G, Karl T, Tong E, et al. Triiodothyronine supplementation in infants and children undergoing cardiopulmonary bypass (TRICC): A multicenter placebo-controlled randomized trial: Age analysis. *Circulation*.

2010;122:224–33.

58. Marwali EM, Lopolisa A, Sani AA, Rayhan M, Roebiono PS, Fakhri D, et al. Indonesian study: Triiodothyronine for infants less than 5 months undergoing cardiopulmonary bypass. *Pediatr Cardiol*. 2022;43:726–34.
59. Watson RS, Crow SS, Hartman ME, Lacroix J, Odetola FO. Epidemiology and outcomes of pediatric multiple organ dysfunction syndrome. *Pediatr Crit Care Med*. 2017;18:S4–16.
60. Leteurtre S, Martinot A, Duhamel A, Gauvin F, Grandbastien B, Thi Vu Nam, et al. Development of a pediatric multiple organ dysfunction score. *Med Decis Mak*. 1999;19:399–410.
61. Leteurtre S, Duhamel A, Salleron J, Grandbastien B, Lacroix J, Leclerc F. PELOD-2. *Crit Care Med*. 2013;41:1761–73.
62. El-Nawawy A, Mohsen AA, Abdel-Malik M, Taman SO. Performance of the pediatric logistic organ dysfunction (PELOD) and (PELOD-2) scores in a pediatric intensive care unit of a developing country. *Eur J Pediatr*. 2017;176:849–55.
63. Yanni GN, Destariani CP, Lubis AN, Deliana M. Thyroid hormone profile in children with sepsis: Does euthyroid sick syndrome exist? *Open Access Maced J Med Sci*. 2019;7:1110–3.
64. Simanjuntak YR, Saputra I, Triratna S, Bakri A, Iriani Y. Validation of PELOD-2 score as a predictor of life-threatening organ dysfunction in pediatric sepsis. *Paediatr Indones*. 2020;60:227–32.
65. Winiarti E, Kosim MS, Supriatna M. Predicting PICU patient survival prognosis: Pediatric Logistic Organ Dysfunction vs Pediatric Index of Mortality scores. *Paediatr Indones*. 2012;52:165.
66. Colucci P, Yue CS, Ducharme M, Benvenega S. A review of the pharmacokinetics of levothyroxine for the treatment of hypothyroidism. *Eur Endocrinol*. 2013;9:40–7.
67. Strich D, Edri S, Gillis D. Current normal values for TSH and FT3 in children are too low: evidence from over 11,000 samples. *J Pediatr Endocrinol Metab*. 2012;25:245–8.
68. K. M, Sharma N. Clinical profile and outcome of patients admitted to pediatric intensive care unit in a tertiary care teaching hospital of Puducherry, India. *Int J Contemp Pediatr*. 2020;7:1280.
69. Muruli B., Bhavya G, K.M C. A study to evaluate serum free triiodothyronine levels as a predictive indicator of outcome in critically ill children. *Int J Contemp Pediatr*. 2018;5:883.
70. Djoko K, Donny KM. Asupan Iodium Anak Usia Sekolah di Indonesia. *Gizi Indones*. 2010;33:8–19.

71. Hernandez A. Type 3 deiodinase is critical for the maturation and function of the thyroid axis. *J Clin Invest.* 2006;116:476–84.
72. Wajner SM, Maia AL. New insights toward the acute non-thyroidal illness syndrome. *Front Endocrinol (Lausanne).* 2012;3:1–7.
73. Calcaterra V, Biganzoli G, Dilillo D, Mannarino S, Fiori L, Pelizzo G, et al. Non-thyroidal illness syndrome and SARS-CoV-2-associated multisystem inflammatory syndrome in children. *J Endocrinol Invest.* 2022;45:199–208.
74. Shao P, Guo S, Li G, Qin D, Li S, Luan Y. Determinants of deranged thyroid function parameters in children admitted for management of diabetic ketoacidosis/diabetic ketosis. *BMC Endocr Disord.* 2020;20:135.
75. Marks SD, Haines C, Rebeyka IM, Couch RM. Hypothalamic-pituitary-thyroid axis changes in children after cardiac surgery. *J Clin Endocrinol Metab.* 2009;94:2781–6.
76. Zhang L, Wu Y, Huang H, Liu C, Cheng Y, Xu L, et al. Performance of PRISM III, PELOD-2, and P-MODS scores in two pediatric intensive care units in China. *Front Pediatr.* 2021;9.
77. Passos L, Barbosa AM. Nonthyroidal illness syndrome as marker of severity in pediatric high dependency units: A single-center prospective cross-sectional observation study from Brazil. *J Pediatr Crit Care.* 2022;9:5.

