

DAFTAR PUSTAKA

- Abdou, H.M. (1989). *Dissolution, bioavailability and bioequivalence*. Pennsylvania: Mark Publishing Company Easton.
- Ankit, M., Manish, Y., & Dinesh, C. (2014). Solid dispersion: A technique to improve solubility of poorly water soluble drug. *Indo Am J Pharm Res*, 4(6), 2855-66.
- Ansel, Howard C. (2008). *Pengantar Bentuk Sediaan Farmasi*. Jakarta: Universitas Indonesia Press.
- Banakar, U.V. (1991). *Pharmaceutical Dissolution Testing*. New York: Marcel Dekker, Inc.
- Barzegar-Jalali, M., Valizadeh, H., Shadbad, M. R. S., Adibkia, K., Mohammadi, G., Farahani, A., ... & Nokhodchi, A. (2010). Cogrounding as an approach to enhance dissolution rate of a poorly water-soluble drug (gliclazide). *Powder Technology*, 197(3), 150-158.
- Borba, P. A. A., Pinotti, M., Andrade, G. R. S., da Costa, N. B., Olchanheski, L. R., Fernandes, D., ... & Stulzer, H. K. (2015). The effect of mechanical grinding on the formation, crystalline changes and dissolution behaviour of the inclusion complex of telmisartan and β -cyclodextrins. *Carbohydrate polymers*, 133, 373-383.
- British Pharmacopoeia Commission. (2009). *British Pharmacopoeia 2009*. London: The Pharmaceutical Press.
- Brittain, H. G. (2009). *Profiles of Drug Substances, Excipients and Related Methodology Volume 34*. Burlington: Academic Press.
- Chiou, W. L., & Riegelman, S. (1971). Pharmaceutical applications of solid dispersion systems. *Journal of pharmaceutical sciences*, 60(9), 1281-1302.
- Chivate, N., Anuradha C., Shubhangi S., Jagdish S. (2012). Improvement of Bioavailability and Solubility of Telmisartan by Solid Dispersion Technique using Various Carriers. *Int. J. Pharm. Sci. Rev. Res.*, 19(2), 36-41.
- Chono, S., Takeda, E., Seki, T., & Morimoto, K. (2008). Enhancement of the dissolution rate and gastrointestinal absorption of pranlukast as a model poorly water-soluble drug by grinding with gelatin. *International journal of pharmaceutics*, 347(1), 71-78.
- Costa, P., & Lobo, J. M. S. (2001). Modeling and comparison of dissolution profiles. *European journal of pharmaceutical sciences*, 13(2), 123-133.

- Dachriyanus. (2004). *Analisis struktur Senyawa organik secara spektroskopi*. Padang: Universitas Andalas.
- Dammann, H. G., Müller, P., & Simon, B. (1983). 24 Hour intragastric acidity and single night-time dose of three H₂-blockers. *The Lancet*, 322(8358), 1078.
- David, S. E., Timmins, P., & Conway, B. R. (2012). Impact of the counterion on the solubility and physicochemical properties of salts of carboxylic acid drugs. *Drug development and industrial pharmacy*, 38(1), 93-103.
- Departemen Kesehatan RI. (1979). *Farmakope Indonesia Edisi III*. Jakarta: Departemen Kesehatan Republik Indonesia.
- Departemen Kesehatan RI. (2014). *Farmakope Indonesia Edisi V*. Jakarta: Departemen Kesehatan Republik Indonesia.
- DrugBank. Famotidine, diakses pada tanggal 9 Februari 2016 dari <http://www.drugbank.ca/drugs/DB00927>.
- Fahmy, R. H., & Kassem, M. A. (2008). Enhancement of famotidine dissolution rate through liquisolid tablets formulation: in vitro and in vivo evaluation. *European Journal of Pharmaceutics and Biopharmaceutics*, 69(3), 993-1003.
- Fernandes, N. C., Jagdale, S. C., Chabukswar, A. R., & Kuchekar, B. S. (2009). Superdisintegrants effect on three model drugs from different BCS classes. *Research J. Pharm. and Tech*, 2(2), 335-337.
- Fillah, U. A. (2014). *Pembentukan Kokristal Ibuprofen-Famotidin dan Evaluasi Laju Disolusinya. (Skripsi)*. Padang: Universitas Andalas.
- Geneidi, A. S., ElShamy, A. H., Awad, G. A. S., Hassan, A. O. K., & Eldemerdash, E. (2004). Solid dispersion of famotidine: factorially designedcapsule formulation and in vivo evaluation of antiulcer activity. *Saudi Pharm. J*, 12, 119-129.
- Ibrahim, M. A., & El-Badry, M. (2014). Formulation of immediate release pellets containing famotidine solid dispersions. *Saudi Pharmaceutical Journal*, 22(2), 149-156.
- Kalia, A., & Poddar, M. (2011). Solid dispersions: an approach towards enhancing dissolution rate. *Int J Pharm Pharm Sci*, 3(4), 9-19.
- Katzung, B. G., Masters, S. B., Trevor, A. J. (2012). *Basic and Clinical Pharmacology 12th Edition*. San Francisco: McGraw Hill Companies.

- Kawabata, Y., Wada, K., Nakatani, M., Yamada, S., & Onoue, S. (2011). Formulation design for poorly water-soluble drugs based on biopharmaceutics classification system: basic approaches and practical applications. *International Journal of Pharmaceutics*, 420(1), 1-10.
- Komalasari, P. A. (2013). Karakterisasi Dispersi Padat Famotidin menggunakan PEG 6000 dan Sorbitol. (*Skripsi*). Padang: Universitas Andalas.
- Krishnamoorthy, V., & Prasad, V. P. R. (2012). Physicochemical characterization and in vitro dissolution behavior of olanzapine-mannitol solid dispersions. *Brazilian Journal of Pharmaceutical Sciences*, 48(2), 243-255.
- Li, D. X., Han, M. J., Balakrishnan, P., Yan, Y. D., Oh, D. H., Joe, J. H., ... & Choi, H. G. (2010). Enhanced oral bioavailability of flurbiprofen by combined use of micelle solution and inclusion compound. *Archives of pharmacol research*, 33(1), 95-101.
- Martin, A., J. Swarbrick., & A Cammart. (1993). *Farmasi Fisika: Dasar-dasar Farmasi Fisika dalam Ilmu Farmasetik*. Penerjemah: Yoshita. Jakarta: UI Press.
- McEvoy, G. K. (2011). *AHFS: Drug Information Essentials*. Bethesda, MD: American Society of Health-System Pharmacist.
- Meinisasti, R., Halim, A., & Zaini, E. (2015). Karakterisasi Fisikokimia Sistem Biner Siprofloksasin HCl-PEG 4000. *Jurnal Sains Farmasi & Klinis*, 2(1), 30-35.
- Nikghalb, L. A., Singh, G., Singh, G., & Kahkeshan, K. F. (2012). Solid dispersion: methods and polymers to increase the solubility of poorly soluble drugs. *Journal of Applied Pharmaceutical Science*, 2(10), 170-175.
- Patil, J.S., Kadam, D.V., Marapur, S.C., & Kamalapur, M.V. (2010). Inclusion Complex System: A Novel Technique to Improve Solubility and Bioavailability of Poorly Soluble Drugs: A Review. *International Journal of Pharmaceutical Sciences Review and Research*. 2(2): 29-34.
- Pubchem. Famotidine, diakses pada tanggal 8 Februari 2016 dari <https://pubchem.ncbi.nlm.nih.gov/compound/famotidine#section=Chemical-and-Physical-Properties>.
- Punitha, S., Vedha Hari, B. N., & Karthikeyan, D. (2010). Enhancement of celecoxib solubility by solid dispersion using mannitol. *Int J Pharm Pharm Sci*, 2(4), 109-111.

- Reddy, V. V., Aswini, M., Harini, P., Prathyusha, B., & Sreevani, M. S. (2013). Design and Characterization of Microcrystals for Enhanced Dissolution Rate of Telmisartan. *International Journal of Innovative Pharmaceutical Research*, 4(1), 263-268.
- Rowe, R.C., Sheskey, P.J., & Quinn, M.E. (2009). *Handbook of pharmaceutical excipient*, (6nd Ed). London: The Pharmaceutical Press.
- Runa, F., Sarkar, M. R., Sultana, R., Jahan, K., & Labu, Z. K. (2013). Study on dissolution improvement of allopurinol by co-grinding and fusion method using solid dispersion technique. *Journal of Biomedical and Pharmaceutical Research*, 2(3), 1-7.
- Sahoo, S., Chakraborti, C. K., Mishra, S. C., & Nanda, U. N. (2011). Scanning Electron Microscopy as an analytical tool for particle size distribution and aspect ratio analysis of Ciprofloxacin Mucoadhesive Polymeric suspension. *Int JRRAS*;6:94-100.
- Shargel, L., Wu-Pong, S., & Yu, A.B.C. (2012). *Biofarmasetika dan Farmakokinetika Terapan*. Terjemahan oleh Budi Suprpti, Edisi Kelima. Surabaya: Airlangga University Press.
- Singh, S., Kasture, S. B., Mohanty, P. K., Jaliwala, Y., Karchuli, M. S., Agarwal, A., & Yadav, Y. (2011). A Review on Solid Sispersion. *Int. J. of Pharm. & Life Sci.(IJPLS)*, 2(9), 1035-1040.
- Someshwar, K., Rama, G., Harikiran, L., Krishna, K., & Srinivas, A. (2011). Dissolution Enhancement of a poorly water soluble drug using water soluble carriers. *Journal of Advanced Pharmaceutical Science*, 1(1), 99-110.
- Spence, J. K., Bhattachar, S. N., Wesley, J. A., Martin, P. J., & Babu, S. R. (2005). Increased dissolution rate and bioavailability through comicronization with microcrystalline cellulose. *Pharmaceutical development and technology*, 10(4), 451-460.
- Sweetman, S.C. (2009). *Martindale Ed 36*. London: Pharmaceutical Press.
- Troy, D. (2005). *Remington: The Science and Practice of Pharmacy 21st Edition*. Philadelphia: Lippincot William and Wilkins.
- Zaini, E., Yeyet, C.S., Soewandhi, S.N., & Halim, A. (2010). Identifikasi Interaksi Fisika antara Trimetoprim dan Sulfametoksazol dengan Metode Kontak Kofler dan Reaksi Kristalisasi. *Majalah Farmasi Indonesia*, 21(1), 32-39.

- Zaini, E., Witarsah, A. S., & Agustin, R. (2014). Enhancement of Dissolution Rate of Meloxicam by Co-grinding Technique using Hydroxypropyl methylcellulose. *Journal of Chemical and Pharmaceutical Research*, 6(11), 263-267.
- Zhong, L., Zhu, X., Luo, X., & Su, W. (2013). Dissolution properties and physical characterization of telmisartan–chitosan solid dispersions prepared by mechanochemical activation. *AAPS PharmSciTech*, 14(2), 541-550.

