

ABSTRAK

KARAKTERISTIK KASUS PLASENTA PREVIA DI RSUP DR. M. DJAMIL JANUARI 2014 – DESEMBER 2015

Oleh:

Telu Rahmat Chamtio Meyanto


Kematian ibu masih menjadi masalah kesehatan di Indonesia. Penyebab kematian ibu terbanyak di Indonesia adalah perdarahan obstetri dan plasenta previa merupakan salah satu perdarahan obstetri yang sering terjadi. Penyebab plasenta previa belum diketahui dengan pasti, namun terdapat beberapa faktor yang meningkatkan insiden plasenta previa, seperti usia risiko tinggi, paritas tinggi, riwayat seksio sesarea, dan riwayat kuretase.

Penelitian telah dilakukan di RSUP Dr. M. Djamil Padang terhadap data rekam medis semua ibu hamil yang didiagnosis dengan plasenta previa periode Januari 2014 – Desember 2015 untuk mengetahui karakteristik kasus plasenta previa. Penelitian ini menggunakan metode deskriptif dengan desain observasional. Data yang terkumpul diolah, dan disajikan dalam bentuk tabel.

Penelitian ini mendapatkan jumlah plasenta previa di RSUP Dr. M. Djamil periode Januari 2014 – Desember 2015 sebanyak 38 kasus. Persentase kasus terbanyak ditemukan pada kelompok usia faktor risiko (52,6%), kelompok multipara (63,2%), kelompok yang memiliki riwayat seksio sesarea (60,5%), dan kelompok yang tidak memiliki riwayat kuretase (55,3%).

Berdasarkan hasil penelitian dapat disimpulkan bahwa persentase kasus plasenta previa terbanyak pada usia risiko tinggi (<20 tahun dan >35 tahun), paritas terbanyak pada kelompok multiparitas, kelompok ada riwayat seksio sesarea dan kelompok tidak ada riwayat kuretase.

Kata kunci : Plasenta previa, usia, paritas, seksio sesarea, kuretase



ABSTRACT

CHARACTERISTIC OF PLACENTA PREVIA CASES IN M. DJAMIL HOSPITAL ON JANUARY 2014 – DECEMBER 2015

By

Telu Rahmat Chamtio Meyanto

Maternal mortality is still a health problem in Indonesia. The mayor causes of maternal death in Indonesia is obstetrical bleeding and placenta previa is one of obstetrical bleeding that often occurs. The etiology of placenta previa is still unknown, but there are some factors that increase the incidence of placenta previa, like high risk age, multiparity, history of cesarean section, and history of currettage.

The research in Dr. M. Djamil Hospital had been done to medical record of all gravid patients that diagnosed with placenta previa on January 2014 – December 2015 to know the characteristics of plasenta previa cases. This research used descriptive methode with observational design. All data that collected is processed and presented in tabular form.

This research found 38 cases of placenta previa at Dr. M. Djamil Hospital on January 2014 – December 2015. The high percentration of case was found in high risk age group (52,6%), multiparity group (63,2%), the group that have history of cesarean section (60,5%), and the group that don't have history of currettage (55,3%).

Based on the result of this research, we can conclude that the highest percentration of placenta previa was found in high risk age group (<20 years and >35 years), multiparity group, the group that have history of cesarean section, and the group that don't have history of currettage.

Keywords : Placenta previa, age, parity, cesarean section, currettage

