

DAFTAR PUSTAKA

- Alladin A, Harrison M. Preeclampsia : Systemic endothelial damage leading to increased activation of the blood coagulation cascade. 2012. Journal of Biotech Research vol.4 p.26-43. USA.
- Amraoui F, Spijkers L, Lahsinoui HH, Vogt L, Post JVD, Peters S et al. sFlt-1 elevates blood pressure by augmenting endothelin-1-mediated vasoconstriction in mice. 2014. Plos One vol. 9 issue.3. Netherland.
- Arriaga PI, Herraiz I, Jimenez EAL, Montes EG, Denk B, Galindo A. Uterine artery doppler and sFlt-1/PIGF ratio : usefulness in diagnosis of preeclampsia. 2013. Ultrasound Obstet Gynecology vol.41 p.530-537. ISUOG. Germany : John Wiley & Sons.
- Bell JM, et al. A historical overview of preeclampsia/eclampsia. 2010. Journal of Obstet Gynecol Neonatal Nurs vol.39 no.5 p.510-528. Pittsburgh : NIH Public Access
- Bilano LV, Ota E, Ganchimeg T, Mori R, Souza JP. Risk factor of preeclampsia/eclampsia and its adverse outcome in low and middle income countries : a WHO secondary analysis. 2014. Plos One vol. 9 issue.3. Switzerland.
- Cunningham FG, Leveno JK, Bloom SL, Spong CY, Dashe JS, Hoffman BL et al. *Williams Obstetrics*. 24nd ed. Chapter 10 : Hypertensive Disorder. 2014. New York : Williams and Wilkins.
- Dharma R, Wibowo N, Raranta HPT. Disfungsi endotel pada preeklampsia. 2005. Makara Kesehatan vol. 9 no.2 p.63-69. Jakarta : Universitas Indonesia
- Donato V, Lacquaniti A, Cernaro V, Buemi A, Lupica R, Lucisano S, et al. Before and after preeclampsia : what biomarker are there. 2012. Current woman's health review vol.8 p.269-275. Italy : Bentham Science Publisher
- Hagman H, Thadhani R, Benzing T, Karumanchi SA, Stepan H. The promises of angiogenic markers for the early diagnosis and prediction of preeclampsia. 2012. Clinical Chemistry vol.58. Germany : The American Association for Clinical Chemistry
- Hagmann H, Bosung V, Belaidi AA, Fridman A, Karumanchi SA, Thadhani R et al. Low molecular weight heparin increase circulating sFlt-1 levels and enhances urinary elimination. 2014. Plos One vol.9 issue.1. USA
- Haram K, Svendsen E, Abildgaard U. The HELLP syndrome : clinical issues and management. A review. 2009. BMC Pregnancy and Childbirth. Norway : Biomed Central
- Hassan MF, Rund NMA, Salama AH. An elevated maternal sFlt-1 to PIGF ratio at midtrimester is a useful predictor for preeclampsia. 2013. Hindawi Publishing Corporations Obstetrics and Gynecology International.

- Hirokoshi K, Maeshima Y, Kobayashi K, Matsuura E, Sugiyama H, Yamasaki Y et al. Elevated serum sFlt-1/Ang-2 ratio in woman with preeclampsia. 2007. *Nephron Clinical Practices* vol.106 p.43-50. Japan : S.Karger AG
- Hladunewich M, Karumanchi SA, Lafayette R. Pathophysiology of the clinical manifestations of preeclampsia. 2007. *Clinical Journal of American Social Nephrology* vol.2 p.543-549. Canada. American Society of Nephrology
- Kemenkes. Kualitas Pelayanan Kesehatan Ibu di Indonesia. 2014. Kemenkes RI
- Kita N, Mitshushita J. A possible placental factor for preeclampsia : sFlt-1. 2008. *Current Medical Chemistry* vol.15 p.711-715. Japan : Bentham Science Publisher.
- Levine RJ, Maynard SE, Qian C, Lim KH, England LJ, et al. Circulating angiogenic factors and the risk of preeclampsia. 2004. *The New England Journal of Medicine*. Boston
- Levine RJ. Preeclampsia, a soluble fms-tyrosine kinase 1, and the risk of reduced thyroid function. 2009. *BMJ*. USA
- Li Q, Et al. The use of artemisinin compounds as angiogenesis inhibitor. 2013. Division of experimental therapeutics, Walter Reed Army Institute of Research. Silver Spring USA
- Lukito SJ, Dewi P. Gambaran Histopatologi Arteri Spiralis Alas Plasenta pada Preeklampsia/Eklampsia dan Kehamilan Normotensive. *Majalah Kedokteran Nusantara*. volume 40. No.3. p173-9. 2007. Universitas Sumatera Utara
- Makris A, Thornton C, Thompson J, Thompson S, Martin R, Ogle R et al. Uteroplacental ischemia result in proteinuric hypertension and elevated sFlt-1. *Kidney International*, p.977-984. 2007. Australia. International Society of Nephrology.
- Nevo O, Lee DK, Caniggia I. Attenuation of VEGFR-2 expression by sFlt-1 and low oxygen in human placenta. 2013. *Plos One* vol.8 issue.11. Canada
- Ohkuchi A, Hirashima C, Suzuki H, Takahashi K, Yoshida M, Matsubara S et al. Evaluation of a new and automated electrochemiluminescent immunoassay for plasma sFlt-1 and PIGF levels in woman with preeclampsia. 2010. *Hypertension research* vol.33 p.422-427. Japan : The Japanese Society of Hypertension.
- Palmer K, Saglam B, Whitehead C, Stock O, Lappas M, Tong S. Severe early onset preeclampsia is not associated with COMT expression. 2011. *The American Journal of Pathology* vol.178 no.6. Australia : elsevier inc.
- Powe EC, Levine RJ, Karumanchi SA. Preeclampsia, a disease of the maternal endothelium The Role of Antiangiogenic Factors and Implications for Later Cardiovascular Disease. 2011. Boston : Harvard medical school. American Heart Association Inc
- Putra R J. Hubungan Faktor Biologik Terhadap Kadar sFlt1 PAda Wanita Hamil Dengan Preeklampsia Berat. 2010. Tesis.Universitas Andalas.

Rekam Medik, 2011. Bagian Obstetri dan Ginekologi RSUP. Dr. M. Djamil Padang periode 1 Januari sampai 31 Desember 2011.

_____, 2012. Bagian Obstetri dan Ginekologi RSUP. Dr. M. Djamil Padang periode 1 Januari sampai 31 Desember 2012.

_____, 2013. Bagian Obstetri dan Ginekologi RSUP. Dr. M. Djamil Padang periode 1 Januari sampai 31 Desember 2013.

Schaarschmidt W. The Course Of Angiogenic Factors in Early Vs Late onset Preeclampsia. 2013. Journal of Perinatologi Medicine Vol. 41 no 5 P511-516. Germany.

Schoofs K, Grittner U, Engels T, Pape J, Denk B et al. The importance of repeated measurements of the sFlt-1/PIGF ratio for the prediction of preeclampsia and intrauterine growth restriction. 2014. Journal of Perinatologi Medicine vol.42 no.1 p.61-68. Germany : Charite University Medicine

Taylor RN, Cunningham FG, *Chesley's Hypertensive Disorder in Pregnancy*. 4th ed. 2015. London : Elsevier Inc.

Thein TT, Myint T, Lwin S, Oo WM, Kyaw AK, Myint MK et al. Promoting antenatal care service for early detection of preeclampsia. 2012. WHO South-East Asia Journal of Public Health vol.1 no.3 p.290-298. Myanmar

Vivo DA, Baviera G, Giordano G, Todarello G, Corrado F et al. Endoglin, PIGF and sFlt-1 as markers for predicting preeclampsia. Acta Obstetrica et Gynecologica vol.87 p.837-842. Italy : Universitas of messina

World Health Organisation. Who recommendation for prevention and treatment of preeclampsia and eclampsia. 2011. Switzerland : WHO.

Xiong Y, Lieberman DA, Tront JS, Huang Y, Hoffman B et al. Gadd45a stress signaling regulates sFlt-1 expression in preeclampsia. 2009. Journal of cellular Physiology vol.220 p.632-639. Pennsylvania : Wiley-Liss Inc

