

DAFTAR PUSTAKA

- Allaire AD, Ballenger KA, Wells SR, McMahon MJ, Lessey BA. Placental apoptosis in preeclampsia. *Obstet Gynecol* 2000;96:271 e6
- Aly AS, Khandelwal M, Zhao J, Mehmet AH, Sammel MD, Parry S. Neutrophils are stimulated by syncytiotrophoblast microvillous membranes to generate superoxide radicals in women with preeclampsia. *Am.J.Obstet Gynecol* 2004; 190:252-8.
- Black S, Kadyrov M, Kaufmann P, Ugele B, Emans N, Huppertz B. Syncytial fusion of human trophoblast depends on caspase 8. *Cell Death Differ* 2004;11:90 e8.
- Barden A, Graham D, Beilin LJ, Ritchie J, Baker R, Walters BN, et al. Neutrophil CD11B expression and neutrophil activation in preeclampsia. *Clin Sci (Lond)* 1997;92:37-44.
- Baker PN, Cunningham FG. (2009). Platelet and coagulation abnormalities. In R. J. Lindheimer MD, *Chesley's Hypertensive Disorders* (2nd Edition ed., pp. 349–375). Connecticut: Elsevier.
- Berthold H 2008. Placental Origins of Preeclampsia. *In : Challenging the Current Hypothesis. American Heart Association.*
- Chen Q, Chen L, Liu B, Vialli C, Stone P, Ching LM, Chamley L. (2010). The role of autocrine TGFbeta1 in endothelial cell activation induced by phagocytosis of necrotic trophoblasts: a possible role in the pathogenesis of pre-eclampsia. *J Pathol* , 87-95.
- Côté, Brown, Lam, Dadelszen, Firoz, Liston. (2008). Diagnostic accuracy of urinary spot protein:creatinine ratio for proteinuria in hypertensive pregnant women: systematic review. *BMJ* .
- Cunningham, F. (2014). Hypertensive Disorders in Pregnancy. In *Williams Obstetrics, 24th eds* (pp. 1515-1516). New York: The McGraw-Hill Companies.
- Dekker GA and Sibai BM. The Immunology of preeclampsia. In *Seminars in Perinatology*. Philadelphia. WB Saunders Company 1999.23;1.p:24-33
- Duley L., Meher S., Abalos E. (2006). Management of pre-eclampsia. *BMJ* , 463–468.

- Duckitt K., Harrington D. (2005). Risk factors for preeclampsia at antenatal booking: systematic review. *BMJ* , 565.
- Goswami D , D. S. Tannetta , L. A. Magee, A. Fuchisawa, Redman CW, Sargent IL and P. von Dadelszen, Excess Syncytiotrophoblast Microparticle Shedding is a Feature of Early-onset Pre-eclampsia, but not Normotensive Intrauterine Growth Restriction, *Placenta* 2006; 27, 56-61
- Formigli L, Papucci L, Tani A, Schiavone N, Tempestini A, Orlandini GE, et al. Aponecrosis: morphological and biochemical exploration of a syncretic process of cell death sharing apoptosis and necrosis. *J Cell Physiol* 2000;182:41 e9.
- Guller S, Role of the syncytium in placenta-mediated complications of preeclampsia, *Thrombosis Research*. 2009; 124: 389–392
- Group.Kumar P, Sharma JB. (2010). Hypertensive Disorders in Pregnancy. *JIMSA* , 261-266.
- Gupta AK, Holzgreve W, Huppertz B, Malek A, Schneider H, Hahn S. Detection of Fetal DNA and RNA in Placenta-Derived Syncytiotrophoblast Microparticles Generated in Vitro. *Clin.Chem*. 2004; 50:2187-90.
- Hahn.S , Huppertz B, Holzgreve W. Fetal Cells and Cell Free Fetal Nucleic Acids in Maternal Blood: New Tools to Study Abnormal Placentation. *Placenta* .2005; 26:515-526
- Hahn S, Rusterholz C, Hosli I, Lapaire O. Cell-free nucleic acids as potential markers for preeclampsia. *Placenta* 2011;32(Suppl.):S17e20.
- Halicka HD, Bedner E, Darzynkiewicz Z. Segregation of RNA and separate packaging of DNA and RNA in apoptotic bodies during apoptosis. *Exp Cell Res* 2000;260:248 e56.
- Hung TH, Skepper JN, Charnock-Jones DS, Burton GJ. Hypoxiareoxygenation: a potent inducer of apoptotic changes in the human placenta and possible etiological factor in preeclampsia. *Circ Res* 2002;90:1274 e81.
- Huppertz B, Frank HG, Kingdom JC, Reister F, Kaufmann P. Villous cytotrophoblast regulation of the syncytial apoptotic cascade in the human) placenta. *Histochem.Cell Biol*. 1998; 110:495-508
- Huppertz B, Tews DS, Kaufmann P. Apoptosis and syncytial fusion in human placental trophoblast and skeletal muscle. *Int Rev Cytol* 2001; 205:215 e53.

Indumati V., Kodliwadmath MV., Sheela MK. (2011). The Role of serum Electrolytes in Pregnancy induced hypertension. *Journal of Clinical and Diagnostic Research* , 66-69.

Ishihara N, Matsuo H, Murakoshi H, Laoag-Fernandez JB, Samoto T, Maruo T. Increased apoptosis in the syncytiotrophoblast in human term placentas complicated by either preeclampsia or intrauterine growth retardation. *Am.J.Obstet Gynecol* 2002; 186:158-66.

Jenkins MB . Preeclampsia. HYPERLINK "<http://www.emedicine.htm>"
<http://www.emedicine.htm> .

Johansen M, Redman CW, Wilkins T, Sargent IL. Trophoblast deportation in human pregnancy--its relevance for pre-eclampsia. *Placenta* 1999; 20:531-9.

Kadyrov M, Schmitz C, Black S, Kaufmann P, Huppertz B. Pre-eclampsia and maternal anaemia display reduced apoptosis and opposite invasive phenotypes of extravillous trophoblast. *Placenta* 2003; 24:540 e8

Knight M, Redman CW, Linton EA, Sargent IL. Shedding of syncytiotrophoblast microvilli into the maternal circulation in preeclamptic pregnancies. *BJOG* 1998;105:632-40.

Kumar, S., Lo, D.Y.M., Smarason, A.K., Chan, L.Y.S., Linton, E.A., Redman, C.W.G.,Sargent, I.L. Pre-eclampsia is associated with increased levels of circulatingapoptotic microparticles and fetal cell-free DNA. *J. Soc. Gynecol. Invest.* 2000; 7 (Supl.): 181a.

Kumar P., Sharma JB. (2010). Hypertensive Disorders in Pregnancy. *JIMSA* , 261-266.

Lam C, Lim KH, Karumanchi A. (2005). Circulating Angiogenic Factors in the Pathogenesis and Prediction of Preeclampsia. *Hypertension* , 1077-1085.

Leung DN, Smith SC, To KF, Sahota DS, Baker PN. Increased placental apoptosis in pregnancies complicated by preeclampsia. *Am.J.Obstet.Gynecol.* 2001; 184:1249-50.

Leung TN, Zhang J, Lau TK, Chan LY, Lo YM. Increased maternal plasma fetal DNA concentrations in women who eventually develop preeclampsia.*Clin Chem* 2001 Jan;47(1):137e9.

- Levine RJ, Hauth JC, Curet LB, Sibai BM, Catalano PM, Morris CD, et al. Trial of calcium to prevent preeclampsia. *N Engl J Med* 1997 Jul 10;337(2):69e76.
- Levine RJ, Qian C, Leshane ES, Yu KF, England LJ, Schisterman EF, et al. Twostage elevation of cell-free fetal DNA in maternal sera before onset of preeclampsia. *Am J Obstet Gynecol* 2004 Mar;190(3):707e13
- Lim KH., Zhou Y., Janatpour M. (2006). Human cytotrophoblast differentiation / invasion is abnormal in pre-eclampsia. *Int J Gynaecol Obstet* , 23-27.
- Lindheimer MD, Conrad KP, Karumanchi SA. (2008). Renal physiology and disease in pregnancy. In S. a. Giebisch, & H. S. Alpern RJ (Ed.), *The Kidney : Physiology and Pathophysiology* (4th Edition ed., pp. 2339 –2398). San Diego: Elsevier.
- Lo YM, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CW, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997;350:485 e7.
- Lo YM, Tein MS, Lau TK, Haines CJ, Leung TN, Poon PM, et al. Quantitative analysis of fetal DNA in maternal plasma and serum: implications for noninvasive prenatal diagnosis. *Am J Hum Genet* 1998; 62:768 e75
- Lo YM, Lau TK, Zhang J, Leung TN, Chang AM, Hjelm NM, et al. Increased fetal DNA concentrations in the plasma of pregnant women carrying fetuses with trisomy 21. *Clin Chem* 1999;45:1747 e51.
- Louise, Michael, Lucilla, Rennae, Jenny, Philip, et al. (2014). Early Pregnancy Prediction of Preeclampsia in Nulliparous Women, Combining Clinical Risk and Biomarkers. *Hypertension* , 644-652.
- Madi J, sulin D. Angka kematian pasien Preeklampsia dan eklampsia di Perjan RS Dr. M. Djamil padang tahun 1998-2002, bagian Obstetri dan Ginekologi FK Unand/ RS Dr. M. Djamil Padang. Disampaikan pada Kongres POGI XII; Yogyakarta 4-9 Juli 2003
- Manyonda IT. The immunology of preeclampsia. In *The Immunology of human reproduction*. London. Taylor & Francis. 2006.p:79-94
- Moore, AG. (2012). Angiogenic biomarkers for prediction of maternal and neonatal complications in suspected preeclampsia. *J Matern Fetal Neonatal Med* (25), 2651-2657.

- Olav L., Andrew S., Holger S. (2010). The preeclampsia biomarkers soluble fms-like tyrosine kinase-1 and placental growth factor: current knowledge, clinical implications and future application. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 122–129.
- Prawirohardjo, S. (2011). *Ilmu Kebidanan*. Jakarta: PT. Bina Pustaka Sarwono Prawirohardjo.
- Redman CW, Sargent IL. Placental debris, oxidative stress and preeclampsia. *Placenta* 2000;21:597 e602.
- Redman CWG, Sargent IL. (2001). The pathogenesis of Pre-eclampsia. *Gynecol Obstet Fertil*, 29, 518-522.
- Redman CW, Tannetta DS, Dragovic RA, et al. Review: does size matter? Placental debris and the pathophysiology of pre-eclampsia. *Placenta* 2012;33 (Supl.):S48–54
- Roberts, JM., Cooper, DW. (2001). Pathogenesis and genetics of preeclampsia. *Lancet*, 53–56.
- Roberts, JM., Gammil, HS. (2005). Preeclampsia Recent Insight. *Hypertension* (46), 1243-1249.
- Robson SC. Hypertension and renal disease in pregnancy. Dewhurst's Textbook of obstetric and Gynecology for Postgraduates, sixth edition, edited by Edmonds DK. London, Blackwell and Science Ltd. 1999: 166-185
- Reddy A, Zhong XY, Rusterholz C, et al. The effect of labour and placental separation on the shedding of syncytiotrophoblast microparticles, cell-free DNA and mRNA in normal pregnancy and pre-eclampsia. *Placenta* 2008;29(11): 942–949
- Roeshadi RH. Hipertensi dalam kehamilan. Dalam : Ilmu Kedokteran Fetomaternal, edisi perdana. Himpunan Kedokteran Fetomaternal. Perkumpulan Obstetri dan Ginekologi. Surabaya. 2004: 494-99
- Saftlas, AF., Olson, DR., Franks, AL., Atrash, HK., Pokras, R.. (1990). Epidemiology of preeclampsia and eclampsia in the United States 1979-1986. *Am J Obstet Gynecol*, 460-465.
- Saputra, NP. (2014). *Analisa Faktor Risiko Preeklampsia*. Padang: PPDS Obstetri dan Ginekologi FK UNAND.
- Sargent IL, Germain SJ, Sacks GP, Kumar S, Redman CW. Trophoblast deportation and the maternal inflammatory response in pre-eclampsia. *J.Reprod.Immunol*. 2003; 59:153-60.

- Sheehan HL, Lynch JP. (1973). *The pathology of toxemia*. Baltimore: Wilkins and Wilkins.
- Shennan AH, Waugh J. (2003). The measurement of blood pressure and proteinuria in pregnancy. (H. Critchly, A. MacLean, L. Poston, & J. Walker, Eds.) *Preeclampsia* , pp. 305-324.
- Sibai, B. M. (2003). Diagnosis and Management of Gestational Hypertension and Preeclampsia. *The American College of Obstetricians and Gynecologists* , 102 (1).
- Sifakis S, Zaravinos A, Maiz N, Spandidos DA, Nicolaidis KH. First-trimester maternal plasma cell-free fetal DNA and preeclampsia. *Am J Obstet Gynecol* 2009 Nov;201(5)(472):e1e7
- Sofowan S. Preeklampsia – Ekalampsia di beberapa rumah sakit di Indonesia, patogenesis dan kemungkinan pencegahannya. *Maj Obstet Ginekol Indones*, Juli. 2003:27(3)
- Soleymanlou N, Jurisica I, Nevo O, Ietta F, Zhang X, Zamudio S, et al. Molecular evidence of placental hypoxia in preeclampsia. *J Clin Endocrinol Metab* 2005 Jul;90(7): 4299e308
- Steegers EAP, Dadelszen P, Duvekot JJ, Pijnenborg R . (2010). Preeclampsia in Seminar. 376 , 631-644.
- Swinkels DW, de Kok JB, Hendriks JC, Wiegerinck E, Zusterzeel PL, Steegers EA. Hemolysis, elevated liver enzymes, and low platelet count (HELLP) syndrome as a complication of preeclampsia in pregnant women increases the amount of cell-free fetal and maternal DNA in maternal plasma and serum. *Clin Chem* 2002;48(4):650e3.
- Tjoa ML, Cindrova-Davies T, Spasic-Boskovic O, Bianchi DW, Burton GJ. Trophoblastic oxidative stress and the release of cell-free fetal-placental DNA. *Am J Pathol* 2006 Aug;169(2):400e4.
- Wagner, L. (2004). Diagnosis and Management of Preeclampsia. *Am Fam Physician* , 2317-2324.
- Watanaga T, Metznerbauer M, Peter I, Johnson KL, Bianchi DW. Placental volume, as measured by 3-dimensional sonography and levels of maternal plasma cell-free fetal DNA. *Am J Obstet Gynecol* 2005;193(2):496e500.
- Winkstrom 2007. Biochemical And Epidemiological Studies Of Early-Onset Dan Late-Onset Pre Eclampsia. In Digital Comprehensive Summaries From The Faculty Of Medicine

Yu Chen, Huang Yajuan, Rongzhen Jiang, Teng Yincheng. Syncytiotrophoblast-derived microparticle shedding in early-onset and late-onset severe pre-eclampsia *International Journal of Gynecology and Obstetrics* 2012; 119:234–238

Zhou Y, Damsky CH, Fisher SJ. (1997). Preeclampsia is associated with failure of human cytotrophoblasts to mimic a vascular adhesion phenotype. One cause of defective endovascular invasion in this syndrome? *J Clin Invest* , 2152-2164.

Zhong XY, Holzgreve W, Hahn S. The levels of circulatory cell free fetal DNA in maternal plasma are elevated prior to the onset of preeclampsia. *Hypertens Pregnancy* 2002;21:77e83.

