

DAFTAR PUSTAKA

1. Nile S, Nile A, Qiu J, Li L, Jia X, Kai G. COVID-19: Pathogenesis, cytokine storm and therapeutic potential of interferons. *Cytokine Growth Factor Rev.* 2020;53:66.
2. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in wuhan, china, of novel coronavirus–infected pneumonia. *N Engl J Med.* 2020;382(13):1.
3. Samuel I, Wreksoatmodjo BR. Anosmia pada COVID-19. *Cermin Dunia Kedokt.* 2021;48:25–9.
4. Chen Y, Liu Q, Guo D. Emerging coronaviruses: genome structure, replication, and pathogenesis. *J Med Virol.* 2020;92:418–20.
5. WHO (2022). Weekly epidemiological update on COVID-19 - 20 April 2022. <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---12-april-2022> - diakses 23 Juni 2022.
6. Kementerian Kesehatan RI (2022). Situasi terkini perkembangan coronavirus disease (COVID-19) 17 April 2022. <https://infeksiemerging.kemkes.go.id/situasi-infeksi-emerging/situasi-terkini-perkembangan-coronavirus-disease-covid-19-17-april-2022> - diakses 18 April 2022.
7. Pemerintah Provinsi Sumatera Barat (2022). Data pantuan COVID-19 Provinsi Sumatera Barat. <https://corona.sumbarprov.go.id/web> - diakses 18 April 2022.
8. Susilo A, Rumende CM, Pitoyo CW, Santoso WD, Yulianti M, Herikurniawan H, et al. Coronavirus disease 2019: Tinjauan literatur terkini. *J Penyakit Dalam Indones.* 2020 Apr 1;7(1):45–67.
9. Ulfah S, Candi C, Darmawan ES. Risk factors of covid-19 transmission between hospital employees. In: neliti. Masters Program in Public Health, Sebelas Maret University; 2020. p. 30–5.
10. Kementerian Kesehatan Republik Indonesia. Pedoman pencegahan dan pengendalian Coronavirus Disease (COVID-19). 4th ed. Kementerian Kesehatan RI; 2020.
11. Ge H, Wang X, Yuan X, Xiao G, Wang C, Deng T, et al. The epidemiology and clinical information about COVID-19. *Eur J Clin Microbiol Infect Dis.* 2020;39(6):1014–5.
12. Moein ST, Hashemian SMR, Mansourafshar B, Khorram-Tousi A, Tabarsi P, Doty RL. Smell dysfunction: a biomarker for COVID-19. *Int Forum Allergy Rhinol.* 2020;10(8):944–50.

13. WHO (2020). Coronavirus disease (COVID-19). https://www.who.int/health-topics/coronavirus#tab=tab_3 - diakses 12 Juni 2022.
14. Lechien JR, Chiesa-Estomba CM, De Santi DR, Horoi M, Le Bon SD, Rodriguez A, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Oto-Rhino-Laryngology*. 2020 Aug 1;277:2252–9.
15. Schäfer L, Schriever VA, Croy I. Human olfactory dysfunction: causes and consequences. *Cell Tissue Res*. 2021;383(1):569.
16. Mutiawati E, Fahriani M, Mamada SS, Fajar JK, Frediansyah A, Maliga HA, et al. Anosmia and dysgeusia in SARS-CoV-2 infection: Incidence and effects on COVID-19 severity and mortality, and the possible pathobiology mechanisms - a systematic review and meta-analysis. *F1000Research*. 2021;13:5.
17. Poerbonegoro NL, Reksodiputro MH, Sari DP, Mufida T, Rahman MA, Reksodiputro LA, et al. Cross-sectional study on the proportion of smell and taste disturbances in hospitalized COVID-19 patients. *Ann Med Surg*. 2021;71:1–4.
18. Kaye R, Chang CWD, Kazahaya K, Brereton J, Denneny JC. COVID-19 anosmia reporting tool: Initial findings. *Otolaryngol - Head Neck Surg (United States)*. 2020;163(1):133–4.
19. Gautier JF, Ravussin Y. A new symptom of COVID-19: Loss of taste and smell. *Obesity*. 2020;28(5):848.
20. Eliezer M, Hautefort C, Hamel AL, Verillaud B, Herman P, Houdart E, et al. Sudden and complete olfactory loss of function as a possible symptom of COVID-19. *JAMA Otolaryngol - Head Neck Surg*. 2020;146(7):674–5.
21. Vaira LA, Salzano G, Deiana G, De Riu G. Anosmia and ageusia: Common findings in COVID-19 patients. *Laryngoscope*. 2020;130:1787.
22. Klopfenstein T, Zahra H, Kadiane-oussou NJ. New loss of smell and taste : Uncommon symptoms in COVID-19 patients in Nord Franche-Comte cluster, France. *Int J Infect Dis*. 2020;117:22.
23. Bax F, Tascini C, Valente M, Marini A, Surcinelli A, Pellitteri G, et al. Hyposmia and dysgeusia in covid-19. *Neurol Clin Pract*. 2021;11(2):92–6.
24. Mullol J, Alobid I, Mariño-Sánchez F, Izquierdo-Domínguez A, Marin C, Klimek L, et al. The loss of smell and taste in the COVID-19 outbreak: A tale of many countries. *Curr Allergy Asthma Rep*. 2020;20:1–5.
25. Otoum MM, Obeidat MB, Yamin HY, Al husban WA kareem, Alkhlaifat AM. The prevalence of anosmia in COVID-19 patients and the associated factors. *Sch J Appl Med Sci*. 2022;10(2):245–7.

26. Hasan MM, Tamanna NA, Jamal MN, Uddin MJ. The prevalence of olfactory dysfunction and its associated factors in patients with COVID-19 infection. medRxiv. 2021;3–9.
27. Friske MM, Giannone F, Senger M, Seitz R, Hansson AC, Spanagel R. Chronic alcohol intake regulates expression of SARS-CoV2 infection-relevant genes in an organ-specific manner. bioRxiv. 2022;3–22.
28. Zhao Y, Liu Y, Yi F, Zhang J, Xu Z, Liu Y, et al. Type 2 diabetes mellitus impaired nasal immunity and increased the risk of hyposmia in COVID-19 mild pneumonia patients. Int Immunopharmacol. 2021;93:1–7.
29. Castillo-López IY, Govea-Camacho LH, Rodríguez-Torres IA, Recio-Macías DA, Alobid I, Mullo J. Olfactory dysfunction in a mexican population outside of covid-19 pandemic: prevalence and associated factors (the OLFAMEX study). Curr Allergy Asthma Rep. 2020;20(12):1–10.
30. Najafloo R, Majidi J, Asghari A, Aleemardani M, Kamrava SK, Simorgh S, et al. Mechanism of anosmia caused by symptoms of COVID-19 and emerging treatments. ACS Chem Neurosci. 2021;12:3795–802.
31. Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. Nat Rev Cardiol. 2020;17(5):259–60.
32. Hussen H, Alemu ZA. Risk of covid-19 infection and associated factors among healthcare workers: A cross-sectional study at eka kotebe treatment center in Ethiopia. Int J Gen Med. 2021;14:1763–71.
33. Sulistyani, Winarsih DS. Characteristics of anosmia cases in Covid-19 patients : A systematic review. Int Summit Sci Technol Humanit. 2020;19–24.
34. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. J Adv Res. 2020;24:91–6.
35. Trougakos IP, Stamatopoulos K, Terpos E, Tsitsilis OE, Aivalioti E, Paraskevis D, et al. Insights to SARS-CoV-2 life cycle, pathophysiology, and rationalized treatments that target COVID-19 clinical complications. J Biomed Sci. 2021;28(9):1–10.
36. He F, Deng Y, Li W. Coronavirus disease 2019: What we know? J Med Virol. 2020;92(7):719–21.
37. Tsai PH, Lai WY, Lin YY, Luo YH, Lin YT, Chen HK, et al. Clinical manifestation and disease progression in COVID-19 infection. J Chinese Med Assoc. 2021;84(1):3–6.
38. PDPI, PERKI, PAPDI, PERDATIN, IDAI. Pedoman tatalaksana COVID-19. 4th ed. PPDI; 2022.

39. Morfi CW, Junaidi A, Elsesmita, Asrini DN, Pangestu F, Lestari DM, et al. Kajian terkini coronavirus disease 2019 (COVID-19). *J Ilmu Kesehat Indones.* 2020;1(1):2–4.
40. Pusparini. Tes serologi dan polimerase chain reaction (PCR) untuk deteksi SARS-CoV-2/COVID-19. *J Biomedika dan Kesehat.* 2020;3(2):46–7.
41. Syam AF, Zulfa FR, Karuniawati A. Manifestasi klinis dan diagnosis Covid-19. *eJournal Kedokt Indones.* 2021;8(3):224–6.
42. Desai M, Oppenheimer J. The importance of considering olfactory dysfunction during the covid-19 pandemic and in clinical practice. *J Allergy Clin Immunol Pract.* 2021;9(1):7–12.
43. Kumar V, Singla S, Gupta N, Bharati SJ, Garg R, Pandit A, et al. The incidence of anosmia in patients with laboratory-confirmed COVID 19 infection in India: An observational study. *J Anaesthesiol Clin Pharmacol.* 2021;37(1):52–5.
44. Cardoso CC, Rossi ÁD, Galliez RM, Faffe DS, Tanuri A, Castiñeiras TMPP. Olfactory dysfunction in patients with mild COVID-19 during gamma, delta, and omicron waves in Rio de Janeiro, Brazil. *JAMA.* 2022;328(6):582–3.
45. Mercier J, Osman M, Bouiller K, Tipirdamaz C, Gendrin V, Chirouze C, et al. Olfactory dysfunction in COVID-19, new insights from a cohort of 353 patients: The ANOSVID study. *J Med Virol.* 2022;94(10):4762–75.
46. Mubaraki AA, Alrbaiai GT, Sibyani AK, Alhulayfi RM, Alzaidi RS, Almalki HS. Prevalence of anosmia among COVID-19 patients in Taif City, Kingdom of Saudi Arabia. *Saudi Med J.* 2021;42(1):38–43.
47. Mangunkusumo E. Buku ajar ilmu kesehatan telinga hidung tenggorokan kepala & leher. In: Soepardi, E.A Iskandar, N Bashiruddin, J Rstuti RD, editor. 7th ed. Jakarta; 2020.
48. Meng X, Deng Y, Dai Z, Meng Z. COVID-19 and anosmia: A review based on up-to-date knowledge. *Am J Otolaryngol - Head Neck Med Surg.* 2020;41(5):102581.
49. Hwang C. Olfactory neuropathy in severe acute respiratory syndrome: report of a case. *Semant Sch.* 2006;15(1):26–7.
50. Hummel T, Whitcroft KL, Andrews P, Altundag A, Cinghi C, Costanzo RM, et al. Position paper on olfactory dysfunction. *Rhinology.* 2017;54(26):3–16.
51. Rodriguez-Sevilla JJ, Güerri-Fernández R, Bertran Recasens B. Is there less alteration of smell sensation in patients with omicron SARS-CoV-2 variant infection? *Front Med.* 2022;9:1–6.

52. Martin-Sanz E, Riestra J, Yebra L, Larran A, Mancino F, Yanes-Diaz J, et al. Prospective study in 355 patients with suspected COVID-19 infection: Value of cough, subjective hyposmia, and hypogeusia. *Laryngoscope*. 2020;130(11):2674–8.
53. Nouchi A, Chastang J, Miyara M, Lejeune J, Soares A, Ibanez G, et al. Prevalence of hyposmia and hypogeusia in 390 COVID-19 hospitalized patients and outpatients: a cross-sectional study. *Eur J Clin Microbiol Infect Dis*. 2021;40(4):691–7.
54. Bocksberger S, Wagner W, Hummel T, Guggemos W, Seilmaier M, Hoelscher M, et al. Temporary hyposmia in COVID-19 patients. *HNO*. 2020;68(6):440–3.
55. Thomas DC, Baddireddy SM, Kohli D. Anosmia: A review in the context of coronavirus disease 2019 and orofacial pain. *J Am Dent Assoc*. 2020;151(9):696–700.
56. Huriyati E, Nelvia T. Gangguan fungsi penghidu dan pemeriksaannya. *J Kesehat Andalas*. 2014;3(1):1–7.
57. Lao WP, Imam SA, Nguyen SA. Anosmia, hyposmia, and dysgeusia as indicators for positive SARS-CoV-2 infection. *World J Otorhinolaryngol - Head Neck Surg*. 2020;6:22–5.
58. Tian J, Pinto JM, Li L, Zhang S, Sun Z, Wei Y. Identification of viruses in patients with postviral olfactory dysfunction by multiplex reverse-transcription polymerase chain reaction. *Laryngoscope*. 2021;131(1):158–64.
59. Karamali K, Elliott M, Hopkins C. COVID-19 related olfactory dysfunction. *Curr Opin Otolaryngol Head Neck Surg*. 2022;30(1):19–23.
60. Porta-Etessam J, Núñez-Gil IJ, González García N, Fernandez-Perez C, Viana-Llamas MC, Eid CM, et al. COVID-19 anosmia and gustatory symptoms as a prognosis factor: a subanalysis of the HOPE COVID-19 (Health Outcome Predictive Evaluation for COVID-19) registry. *Infection*. 2021;49(4):677–84.
61. Roved J, Westerdahl H, Hasselquist D. Sex differences in immune responses: Hormonal effects, antagonistic selection, and evolutionary consequences. *Horm Behav*. 2017;88:95–105.
62. Diamond MS, Kanneganti TD. Innate immunity: the first line of defense against SARS-CoV-2. *Nat Immunol*. 2022;23(2):165–72.
63. Gori A, Leone F, Loffredo L, Cinicola BL, Brindisi G, De Castro G, et al. Covid-19-related anosmia: The olfactory pathway hypothesis and early intervention. *Front Neurol*. 2020;11:956.
64. Hopkins C, Surda P, Whitehead E, Nirmal Kumar B. Early recovery following new onset anosmia during the COVID-19 pandemic—an observational cohort study. 2020;49(1):1–5.

65. Kaliavaradan S, Bhat PS, Rajagopal M, Vinayagamoorthy V, Vimal M, Janakiraman S, et al. Course of hyposmia and hypogeusia and their relationship with severity of covid-19 disease among indian population. Indian J Otolaryngol Head Neck Surg. 2021;74(2):3078–84.
66. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet (London, England). 2020;395(10223):497–506.
67. Avci H, Karabulut B, Farasoglu A, Boldaz E, Evman M. Relationship between anosmia and hospitalisation in patients with coronavirus disease 2019: an otolaryngological perspective. J Laryngol Otol. 2020;134(8):1.
68. Kementerian Kesehatan RI (2018). Klasifikasi hipertensi. <https://p2ptm.kemkes.go.id/infographic/klasifikasi-hipertensi> - diakses 16 November 2022.
69. Ariani R, Muhammad A, Syahputra E, Sinaga N, Pratiwi D, Nuralita N, et al. Ragam penanganan dan pencegahan COVID-19 di rumah sakit dan klinik primer. 1st ed. Airlangga E, Ariani R, editors. medan: UMSUPRESS; 2020.
70. Rahayu LA, Admiyanti JC, Khalda YI, Adha FR, Agistany NFF. Hipertensi, diabetes melitus dan obesitas sebagai faktor komorbiditas utama terhadap mortalitas pasien Covid-19 : Sebuah studi literatur tinjauan pustaka hypertension, diabetes mellitus, and obesity as the main comorbidity factors of mortality in Covid-19. J Ilm Mhs Kedokt Indones. 2021;9:90–5.
71. Lv H, Zhang W, Zhu Z, Xiong Q, Xiang R, Wang Y, et al. Prevalence and recovery time of olfactory and gustatory dysfunction in hospitalized patients with covid-19 in wuhan, china. Int J Infect Dis. 2020;100:507–12.
72. Awwad AA, Abd Elhay OMM, Rabie MM, Awad EA, Kotb FM, Maghraby HM, et al. Impact of systemic diseases on olfactory function in covid-19 infected patients. Int J Gen Med. 2022;15:5681–91.
73. Ferraro S, Tuccori M, Convertino I, Valdiserra G, Cappello E, Maggi F, et al. Olfactory and gustatory impairments in COVID-19 patients : Role in early diagnosis and interferences by concomitant drugs. 2021;2:2186–8.
74. Guillén Martínez A, Andreu Gálvez M, Rodríguez Sanz S, Hernández Ruiz P, García Morillas A, Esteban Sánchez T. Incidence of smell and taste disorders and associated factors in patients with mild to moderate COVID-19. Otolaryngol Pol = Polish Otolaryngol. 2020;75(2):31–8.
75. Amer MA, Elsherif HS, Abdel-Hamid AS, Elzayat S. Early recovery patterns of olfactory disorders in COVID-19 patients; a clinical cohort study. Am J Otolaryngol. 2020;41(6):1–4.

76. Soelistijo SA, Suastika D, Decroli E, Permana H, Sucipto KW, Kusnadi Y, et al. Pedoman pengelolaan dan pencegahan diabetes melitus tipe 2 dewasa di Indonesia. In: Perkumpulan Endokrinologi Indonesia. PB. PERKENI; 2021. p. 6–46.
77. Fauzia NS. Faktor risiko terjadinya coronavirus disease 2019 pada pasien dengan komorbid diabetes melitus. Arter J Ilmu Kesehat. 2021;2(4):107–10.
78. Kim SJ, Windon MJ, Lin SY. The association between diabetes and olfactory impairment in adults: A systematic review and meta-analysis. Laryngoscope Investig Otolaryngol. 2019;4(5):465–75.
79. WHO (2021). Cardiovascular disease (CVD). [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)) - diakses 15 November 2022.
80. Crea F. European Society of Cardiology guidance for the management of cardiovascular disease during the pandemic and a focus on long COVID. Eur Heart J. 2022;43(11):1017–21.
81. Shafter AM, Shaikh K, Johanis A, Budoff MJ. De-risking primary prevention: role of imaging. Ther Adv Cardiovasc Dis. 2021;15:1–6.
82. Roh D, Lee DH, Kim SW, Kim SW, Kim BG, Kim DH, et al. The association between olfactory dysfunction and cardiovascular disease and its risk factors in middle-aged and older adults. Sci Rep. 2021;11(1):1–10.
83. Lim JU, Lee JH, Kim JS, Hwang Y Il, Kim TH, Lim SY, et al. Comparison of world health organization and asia-pacific body mass index classifications in COPD patients. Int J Chron Obstruct Pulmon Dis. 2017;12:2465–72.
84. Moraes AHDA, Passos TS, Vale SHDL, Maia JKDS, Maciel BLL. Obesity and the increased risk for covid-19: Mechanisms and nutritional management. Nutr Res Rev. 2021;34(2):1–9.
85. Hemalatha Rajkumar PB. The impact of obesity on immune response to infection and vaccine: An Insight into Plausible Mechanisms. Endocrinol Metab Syndr. 2013;2(2):2–8.
86. Peng M, Coutts D, Wang T, Cakmak YO. Systematic review of olfactory shifts related to obesity. Obes Rev. 2019;20(2):325–34.
87. WHO (2020). Tobacco use and COVID-19. <https://www.who.int/news/item/11-05-2020-who-statement-tobacco-use-and-covid-19> - diakses 19 Juni 2022.
88. Ismail N, Hassan N, Abd Hamid MHN, Yusoff UN, Khamal NR, Omar MA, et al. Association of smoking and severity of Covid-19 infection among 5,889 patients in Malaysia : A multi-center observational study. Int J Infect Dis. 2022 Mar 1;116:189–95.

89. Jiang C, Chen Q, Xie M. Smoking increases the risk of infectious diseases: A narrative review. *Tob Induc Dis.* 2020 Jul 1;18:1–9.
90. Las Casas Lima MH de, Cavalcante ALB, Leão SC. Pathophysiological relationship between COVID-19 and olfactory dysfunction: A systematic review. *Braz J Otorhinolaryngol.* 2022;88(5):794–801.
91. Kabbani N, Olds JL. Does covid19 infect the brain? if so, smokers might be at a higher risk. *Mol Pharmacol.* 2020;97(5):351–3.
92. Shastri MD, Shukla SD, Chong WC, KC R, Dua K, Patel RP, et al. Smoking and COVID-19: What we know so far. *Respir Med.* 2021 Jan 1;176:1–5.
93. Calina D, Hartung T, Mardare I, Mitroi M, Poulas K, Tsatsakis A, et al. COVID-19 pandemic and alcohol consumption: Impacts and interconnections. *Toxicol Reports.* 2021;8:529–33.
94. Wong DKC, Gendeh HS, Thong HK, Lum SG, Gendeh BS, Saim A, et al. A review of smell and taste dysfunction in COVID-19 patients. *Med J Malaysia.* 2020;75(5):575–80.
95. Cooper KW, Brann DH, Farruggia MC, Bhutani S, Pellegrino R, Tsukahara T, et al. COVID-19 and the chemical senses: Supporting players take center stage. *Neuron.* 2020;107(2):219–27.
96. Butowt R, von Bartheld CS. Anosmia in COVID-19: Underlying mechanisms and assessment of an olfactory route to brain infection. *Sage Journals.* 2020;27(6):582–97.
97. Rumeau C, Nguyen DT, Jankowski R. How to assess olfactory performance with the Sniffin' Sticks test. *Eur Ann Otorhinolaryngol Head Neck Dis.* 2016;133(3):203–6.
98. Niklassen AS, Ovesen T, Fernandes H, Fjaeldstad AW. Danish validation of sniffin' sticks olfactory test for threshold, discrimination, and identification. *Laryngoscope.* 2018;128(8):1759–66.
99. Fenólio GHM, Anselmo-Lima WT, Tomazini GC, Compagnoni IM, Amaral MSA d., Fantucci MZ, et al. Validation of the connecticut olfactory test (CCRC) adapted to Brazil. *Braz J Otorhinolaryngol.* 2022 Sep 1;88(5):725–31.
100. Neta FI, Fernandes ACL, Vale AJM, Pinheiro FI, Cobucci RN, Azevedo EP de, et al. Pathophysiology and possible treatments for olfactory-gustatory disorders in patients affected by Covid-19. *Curr Res Pharmacol Drug Discov.* 2021;2:2–8.
101. Whitcroft KL, Hummel T. Olfactory dysfunction in COVID-19: Diagnosis and management. *JAMA - J Am Med Assoc.* 2020;323(24):2512–4.
102. Dahlan MS. Besar sampel dan cara pengambilan sampel dalam penelitian kedokteran dan kesehatan. In: Edisi 3. Jakarta: Salemba Medika; 2010. p. 36–41.

103. Roberto P, Júnior S, Oliveira AL, Gomes R, Eugênio L, Coelho A, et al. Anosmia and COVID-19: perspectives on its association and the pathophysiological mechanisms involved. *Egypt J Neurol Psychiatry Neurosurg.* 2021;57(1):2–7.
104. Mercante G, Ferrel F, de Virgilio A, Gaino F, Di Bari M, Colombo G, et al. Prevalence of taste and smell dysfunction in coronavirus disease 2019. *JAMA Otolaryngol - Head Neck Surg.* 2020;146(8):723–8.
105. Biadsee A, Biadsee A, Kassem F, Dagan O, Masarwa S, Ormianer Z. Olfactory and oral manifestations of covid-19: sex-related symptoms-a potential pathway to early diagnosis. *Otolaryngol Neck Surg.* 2020;163(4):722–8.
106. Hüttenbrink KB, Hummel T, Berg D, Gasser T, Hähner A. Olfactory dysfunction: Common in later life and early warning of neurodegenerative disease. *Dtsch Arztbl Int.* 2013;110(1–2):1–5.
107. Nugraha MFB, Irfandy D, Yenny SW. Karakteristik pasien rinosinusitis kronik di poliklinik THT-KL RSUP DR. M. Djamil Padang 2017-2019. *J Ilmu Kesehat Indones.* 2021;2(4):279–85.
108. Sánchez-Vallecillo M V., Fraire ME, Baena-Cagnani C, Zernotti ME. Olfactory dysfunction in patients with chronic rhinosinusitis. *Int J Otolaryngol.* 2012;2012:1–4.
109. Dalton P. Olfaction and anosmia in rhinosinusitis. *Curr Allergy Asthma Rep.* 2004;4(3):230–5.
110. Irfandy D, Budiman BJ, Huryati E. Relationship between deviations of nasal septum and mucociliary transport time using saccharin test. *Otorinolaringologia.* 2019;69(1):30–5.
111. Alekseenko SI, Skalny A V., Ajsuvakova OP, Skalnaya MG, Notova S V., Tinkov AA. Mucociliary transport as a link between chronic rhinosinusitis and trace element dysbalance. *Med Hypotheses.* 2019;127:5–8.
112. Mullo J, Mariño-Sánchez F, Valls M, Alobid I, Marin C. The sense of smell in chronic rhinosinusitis. *J Allergy Clin Immunol.* 2020;145(3):773–6.
113. Tabah A, Ramanan M, Laupland KB, Buetti N, Cortegiani A, Mellinghoff J, et al. Personal protective equipment and intensive care unit healthcare worker safety in the COVID-19 era (PPE-SAFE): An international survey. *J Crit Care.* 2020;59:70–5.
114. Villarreal IM, Morato M, Martínez-RuizCoello M, Navarro A, Garcia-Chillerón R, Ruiz Á, et al. Olfactory and taste disorders in healthcare workers with COVID-19 infection. *Eur Arch Oto-Rhino-Laryngology.* 2021;278(6):2123–6.
115. Adelina M, Dwijayanti F. The infection of COVID-19 among health care workers in dharmais cancer hospital. *Indones J Cancer.* 2021;15(1):1.

116. Jha S, Singh RK, Kumar S, Singh T. Institutional evaluation of anosmia and ageusia in COVID-19 positive health care workers of clinical and non-clinical departments. *Int J Res Med Sci.* 2022;10(3):683–6.
117. Yoshimura T, Kitabatake M, Lefèvre N, Corazza F, Valsamis J, Delbaere A, et al. The number of X chromosomes influences inflammatory cytokine production following toll-like receptor stimulation. *Front Immunol.* 2019;10:1–10.
118. Ko CJ, Huang CC, Lin HY, Juan CP, Lan SW, Shyu HY, et al. Androgen-induced TMPRSS2 activates matriptase and promotes extracellular matrix degradation, prostate cancer cell invasion, tumor growth, and metastasis. *Cancer Res.* 2015;75(14):2949–60.
119. Mendonça CV, Mendes Neto JA, Suzuki FA, Orth MS, Machado Neto H, Nacif SR. Olfactory dysfunction in COVID-19: a marker of good prognosis? *Braz J Otorhinolaryngol.* 2022;88(3):439–43.
120. Lechien JR, Chiesa-Estomba CM, Place S, Van Laethem Y, Cabaraux P, Mat Q, et al. Clinical and epidemiological characteristics of 1420 European patients with mild-to-moderate coronavirus disease 2019. *Assoc Publ J Intern Med.* 2020;288:335–43.
121. Surendra H, Elyazar IR, Djaafara BA, Ekawati LL, Saraswati K, Adrian V, et al. Clinical characteristics and mortality associated with COVID-19 in Jakarta, Indonesia: A hospital-based retrospective cohort study. *Lancet Reg Heal - West Pacific.* 2021;9:100108.
122. Setiadi W, Rozi IE, Safari D, Daningrat WOD, Johar E, Yohan B, et al. Prevalence and epidemiological characteristics of COVID-19 after one year of pandemic in Jakarta and neighbouring areas, Indonesia: A single center study. *PLoS One.* 2022;17(5).
123. Htun YM, Win TT, Aung A, Latt TZ, Phyo YN, Tun TM, et al. Initial presenting symptoms, comorbidities and severity of COVID-19 patients during the second wave of epidemic in Myanmar. *Trop Med Health.* 2021;49(1):1–10.
124. Lechien JR, Chiesa-Estomba CM, Vaira LA, De Riu G, Cammaroto G, Chekkouri-Idrissi Y, et al. Epidemiological, otolaryngological, olfactory and gustatory outcomes according to the severity of COVID-19: a study of 2579 patients. *Eur Arch Oto-Rhino-Laryngology.* 2021;278(8):2851–8.
125. Lechien JR, Barillari MR, Jouffe L, Saussez S. Anosmia is a key symptom of COVID-19 infection and should be used as a diagnostic tool. *Ear, Nose Throat J.* 2020;99(9):577–8.
126. Talavera B, García-Azorín D, Martínez-Pías E, Trigo J, Hernández-Pérez I, Valle-Peña G, et al. Anosmia is associated with lower in-hospital mortality in COVID-19. *J Neurol Sci.* 2020;419:117163.

127. Klopfenstein T, Kadiane-Oussou NJ, Toko L, Royer PY, Lepiller Q, Gendrin V, et al. Features of anosmia in COVID-19. *Med Mal Infect*. 2020;50(5):436–9.
128. Khan AS, Hichami A, Khan NA. Obesity and COVID-19: Oro-naso-sensory perception. *J Clin Med*. 2020;9(7):1–7.
129. Paderno A, Schreiber A, Grammatica A, Raffetti E, Tomasoni M, Gualtieri T, et al. Smell and taste alterations in COVID-19: a cross-sectional analysis of different cohorts. *Int Forum Allergy Rhinol*. 2020;10(8):955–61.
130. Speth MM, Singer-Cornelius T, Oberle M, Gengler I, Brockmeier SJ, Sedaghat AR. Olfactory dysfunction and sinonasal symptomatology in COVID-19: Prevalence, severity, timing, and associated characteristics. *Otolaryngol Neck Surg*. 2020;163(1):114–9.
131. Babaei A, Iravani K, Malekpour B, Golkhar B, Soltaniesmaeli A, Hosseinihashemi M. Factors associated with anosmia recovery rate in COVID-19 patients. *Laryngoscope Investig Otolaryngol*. 2021;6(6):1248.

