

## DAFTAR PUSTAKA

- Achmad, F., & Rusmai, T. (2021). The effect of intradialytic stretching training on restless legs syndrome and sleep quality in hemodialysis patients. *Korean Journal of Adult Nursing*, 33(1), 37–43. <https://doi.org/10.7475/kjan.2021.33.1.37>
- Afshar, R., Emany, A., Saremi, A., Shavandi, N., & Sanavi, S. (2011). Effects of intradialytic aerobic training on sleep quality in hemodialysis patients. *Iranian Journal of Kidney Diseases*, 5(2), 119–123. <https://doi.org/10.206/1166201>
- Al Naamani, Z., Gormley, K., Noble, H., Santin, O., & Al Maqbali, M. (2021). Fatigue, anxiety, depression and sleep quality in patients undergoing haemodialysis. *BMC Nephrology*, 22(1), 1–8. <https://doi.org/10.1186/s12882-021-02349-3>
- Aliasgharpour, M., Abbasi, Z., Razi, S. P., & Kazemnezhad, A. (2016). The effect of stretching exercises on severity of restless legs syndrome in patients on hemodialysis. *Asian Journal of Sport Medicine*, 7(2). <https://doi.org/10.5812/asjms.31001.Research>
- Allen, R. P., Burchell, B. J., MacDonald, B., Hening, W. A., & Earley, C. J. (2009). Validation of the self-completed Cambridge-Hopkins questionnaire (CH-RLSq) for ascertainment of restless legs syndrome (RLS) in a population survey. *Sleep Medicine*, 10(10), 1097–1100. <https://doi.org/10.1016/j.sleep.2008.10.007>
- Allen, R. P., Picchiatti, D. L., Garcia-Borreguero, D., Ondo, W. G., Walters, A. S., Winkelman, J. W., Zucconi, M., Ferri, R., Trenkwalder, C., & Lee, H. B. (2014). Restless legs syndrome/willis-ekbom disease diagnostic criteria:

Updated international restless legs syndrome study group (IRLSSG) consensus criteria - history, rationale, description, and significance. *Sleep Medicine*, 15(8), 860–873. <https://doi.org/10.1016/j.sleep.2014.03.025>

Alligood, M. R. (2014). *Nursing theorists and their work*. Elsevier Mosby.

Amini, E., Goudarzi, I., Masoudi, R., Ahmadi, A., & Momeni, A. (2016). Effect of progressive muscle relaxation and aerobic exercise on anxiety, sleep quality, and fatigue in patients with chronic renal failure undergoing hemodialysis. *International Journal of Pharmaceutical and Clinical Research*, 8(12), 1634–1639. <https://doi.org/10.10.316/pms.2016.12.025>

Arismunandar, P., Barus, S. U. B., & Pangesti, E. D. (2021). The effect of intradialytic exercise on fatigue in hemodialysis patients at Dustira Hospital Cimahi. *Egyptian Rheumatology and Rehabilitation*, 48(1), 40–44. <https://doi.org/10.1007/2Fs40620-018-00565-z>

Arnold, R., Pussell, B. A., Howells, J., Grinius, V., Kiernan, M. C., Lin, C. S. Y., & Krishnan, A. V. (2014). Evidence for a causal relationship between hyperkalaemia and axonal dysfunction in end-stage kidney disease. *Clinical Neurophysiology*, 125(1), 179–185. <https://doi.org/10.1016/j.clinph.2013.06.022>

Asmadi. (2008). *Teknik prosedural konsep & aplikasi kebutuhan dasar klien*. Salemba Medika.

Azimpour, S., Hosseini, H. S., Eftekhari, A., & Kazemi, M. (2019). The effects of vibration and massage on severity of symptoms of restless leg syndrome and sleep quality in hemodialysis patients; A randomized cross-over clinical trial. *Journal of Renal Injury Prevention*, 8(2), 106–111.

<https://doi.org/10.15171/jrip.2019.20>

Baumgaertel, M. W., Kraemer, M., & Berlitz, P. (2014). Neurologic complications of acute and chronic renal disease. In *Neurologic Aspects of Systemic Disease Part I* (1st ed., Vol. 119). Elsevier B.V. <https://doi.org/10.1016/B978-0-7020-4086-3.00024-2>

Brouns, R., & De Deyn, P. P. (2004). Neurological complications in renal failure: A review. *Clinical Neurology and Neurosurgery*, 107(1), 1–16. <https://doi.org/10.1016/j.clineuro.2004.07.012>

Buyse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The pittsburgh sleep quality index : A new instrument psychiatric practice and research. *Psychiatry Research*. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4)

Chairunnisa, P., Syukri, M., & Muhsin. (2017). Hubungan lama terapi hemodialisis terhadap pasien usia. *Jurnal Ilmiah Mahasiswa Medisia*, 2(4), 11–16. <https://doi.org/10.7567/15918-1>

Damanik, H. (2020). Tingkat kecemasan pasien gagal ginjal kronik dalam menjalani hemodialisa di Rumah Sakit Imelda Pekerja Indonesia. *Jurnal Ilmiah Keperawatan Imelda*, 6(1), 80–85. <https://doi.org/10.52943/jikeperawatan.v6i1.365>

Darmawan, I. Putu E., Nurhesti, P. O. Y., & Suardana, I. K. (2019). Hubungan lamanya menjalani hemodialisis dengan fatigue pada pasien chronic kidney disease 1. *Community of Publishing in Nursing*, 139–146. <https://doi.org/10.2715/coping.2019.12.030>

Dharma, K. K. (2011). *Metodologi penelitian (panduan melaksanakan dan*



menerapkan hasil penelitian). *Trans Info Media*.

Dzulfachri, M., & Kurniati, T. R. I. (2020). Faktor-faktor yang berhubungan dengan kejadian restless leg syndrom pada pasien gagal ginjal kronik di ruang hemodialisa Rumah Sakit Palang Merah Indonesia Kota Bogor tahun 2020.

*Universitas Muhammadiyah Jakarta*. <https://doi.org/10.18002/fstream.5435>

Fauzi, A. (2018). Efektifitas intradialytic stretching exercise terhadap penurunan gejala restless leg syndrome dan peningkatan sleep quality pada pasien hemodialisis tahun 2018. *Jurnal Kesehatan Bakti Husada*. <https://doi.org/10.1915/bkths.2018.01.501>

Fawzy, M., Elhamed, A., & Ahmed, M. (2021). Restless leg syndrome in patients with chronic kidney disease: A hospital-based study from upper Egypt. *International Journal of Neuroscience*, 0(0), 000. <https://doi.org/10.1080/00207454.2021.1910256>

Forwaty, E., Malini, H., & Oktarina, E. (2019). Pengaruh intradialytic range of motion (ROM) exercise terhadap depresi, insomnia dan asupan nutrisi pada pasien hemodialisis. *Jurnal Kesehatan Andalas*, 8(3), 529. <https://doi.org/10.25077/jka.v8i3.1038>

Gheshlagh, R. G., Farajzadeh, M., Zarei, M., Baghi, V., Dalvand, S., & Sayehmiri, K. (2017). The prevalence of restless legs syndrome in patients undergoing hemodialysis: A systematic review and meta-analysis study. *Basic and Clinical Neuroscience*, 8(2), 105–112.

<https://doi.org/10.18869/nirp.bcn.8.2.105>

Giannaki, C. D., Hadjigeorgiou, G. M., Karatzaferi, C., Maridaki, M. D.,

Koutedakis, Y., Founta, P., Tsianas, N., Stefanidis, I., & Sakkas, G. K. (2013).

A single-blind randomized controlled trial to evaluate the effect of 6 months of progressive aerobic exercise training in patients with uraemic restless legs syndrome. *Nephrology Dialysis Transplantation*, 28(11), 2834–2840. <https://doi.org/10.1093/ndt/gft288>

Giannaki, C. D., Sakkas, G. K., Hadjigeorgiou, G. M., Karatzaferi, C., Patramani, G., Lavdas, E., Liakopoulos, V., Koutedakis, Y., & Stefanidis, I. (2010). Non-pharmacological management of periodic limb movements during hemodialysis session in patients with uremic restless legs syndrome. *ASAIO Journal*, 56(6), 538–542. <https://doi.org/10.1097/MAT.0b013e3181f1cc04>

Guraire, M. (2023). *what is restless legs syndrome*. <https://www.verywellhealth.com/restless-legs-syndrome-symptoms-diagnosis-treatment-7107159?print>

Hmwe, N. T. T., Subramanian, P., Tan, L. P., & Chong, W. K. (2015). The effects of acupressure on depression, anxiety and stress in patients with hemodialysis: A randomized controlled trial. *International Journal of Nursing Studies*, 52(2), 509–518. <https://doi.org/10.1016/j.ijnurstu.2014.11.002>

Hutagaol, A. Y. (2021). *Gambaran tingkat kecemasan berdasarkan karakteristik pasien gagal ginjal kronik dalam menjalani hemodialisa rsup haji adam malik medan tahun 2021* [STIKES Santa Elisabet Medan]. <https://doi.org/10.119/wpc.2022.01.021>

Indrawati, N. (2012). Perbandingan kualitas tidur mahasiswa yang mengikuti ukm dan tidak mengikuti ukm pada mahasiswa reguler fik ui. *Fakultas Ilmu Keperawatan*, 1–84. <https://doi.org/10.203/S42629.08.112>

Innes, K. E., Selfe, T. K., Montgomery, C., Hollingshead, N., Huysmans, Z.,

- Srinivasan, R., Wen, S., Hausmann, M. J., Sherman, K., & Klatt, M. (2020). Effects of a 12-week yoga versus a 12-week educational film intervention on symptoms of restless legs syndrome and related outcomes: An exploratory randomized controlled trial. *Journal of Clinical Sleep Medicine*, 16(1), 107–119. <https://doi.org/10.5664/JCSM.8134>
- Islamiyah, W. R. (2018). *Panduan tatalaksana gangguan tidur* (Cetakan I.). CV. Sagung Seto.
- Jieun, C., & Myungsun, Y. (2014). Symptom clusters and quality of life in patients on hemodialysis. *Journal of Korean Clinical Nursing Research Vol.20 No.1*, 123-133, 20(1), 5–7. <https://doi.org/10.1097/gme.0b013e31827c4627>
- Joni, Y. N., Nur, B. M., & Rayasari, F. (2019). Efektivitas exercise intradialisis menggunakan barbell dan range of motion (ROM) terhadap adekuasi hemodialisa pada pasien penyakit ginjal kronik. *Jurnal Keperawatan Silampari*, 2(2), 189–200. <https://doi.org/10.31539/jks.v2i2.533>
- Kaur, J., Venkatesan, M., Kaur, H., & Rawat, P. S. (2016). Effectiveness of muscle stretching exercise on quality of life of hemodialysis patients. *International Journal of Medical Research & Health Sciences*, 203–207. <https://doi.org/https://10.117/ijmrhs.2016.04.029>
- KDOQI. (2002). Clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. In *National Kidney Foundation* (Vol. 90, Issue 1).
- Keliat, B. A. (2021). *Asuhan Keperawatan Jiwa*. EGC.
- Kemenkes. (2018). Kementerian Kesehatan Republik Indonesia. *Kementerian Kesehatan RI*, 1(1), 1.



<https://www.kemkes.go.id/article/view/19093000001/penyakit-jantung-penyebab-kematian-terbanyak-ke-2-di-indonesia.html>

Kementrian Kesehatan Republik Indonesia. (2019). *Laporan Riskesdas 2018 Nasional.pdf* (p. 674). Bidang Penelitian dan Pengembangan Kesehatan.

Klingelhofer, L., Bhattacharya, K., & Reichmann, H. (2016). Restless legs syndrome. *Clinical Medicine, Journal of the Royal College of Physicians of London*, 16(4), 379–382. <https://doi.org/10.7861/clinmedicine.16-4-379>

Klingelhofer, P. L., Kalyan, S., Bhattacharya, B., & Reichmann, H. (2016). *Sindrom kaki gelisah*. 16(4), 379–382. <https://doi.org/10.7861/clinmedicine.16-4-379>

Kovesdy, C. P. (2022). Epidemiology of chronic kidney disease: An update 2022. *Kidney International Supplements*, 12(1), 7–11. <https://doi.org/10.1016/j.kisu.2021.11.003>

Kusuma, H. (2019). *Buku panduan mengenal penyakit ginjal kronis dan perawatannya*.

[http://eprints.undip.ac.id/81430/1/Buku\\_Panduan\\_Mengenal\\_Penyakit\\_Ginjal\\_Kronis\\_dan\\_Perawatannya\\_Henni\\_Kusuma%2C\\_Suhartini%2C\\_Untung\\_Sujianto%2C\\_Chandra\\_Bagus\\_Ropiyanto%2C\\_Wahyu\\_Hidayati.pdf](http://eprints.undip.ac.id/81430/1/Buku_Panduan_Mengenal_Penyakit_Ginjal_Kronis_dan_Perawatannya_Henni_Kusuma%2C_Suhartini%2C_Untung_Sujianto%2C_Chandra_Bagus_Ropiyanto%2C_Wahyu_Hidayati.pdf)

Ladesvita, F. (2016). Perbandingan latihan range of motion ( ROM ) dengan latihan pedal intradialisis terhadap adekuasi hemodialisa pada pasien hemodialisa kronik. *Universitas Indonesia*. <https://doi.org/10.1590/2175-8239>

Laoli, D., Hulu, V. P. P., Buulolo, Y., & ... (2019). Pengaruh latihan fisik selama hemodialisis terhadap kekuatan otot pada pasien gagal ginjal kronik di Rumah Sakit Royal Prima Medan. *Jurnal Keperawatan Stikes Hang Tuah Medan*,

9(2). <https://doi.org/10.136/2019.07.027>

LeMone, P. (2017). *Medical Surgical Nursing - Critical Thinking for Person Centered Care* (3rd ed., Vols. 1–3). Pearson Australia.

Lestari, W. F., Herman, & Mita. (2020). The effect of intradialytic exercise on stress in hemodialysis patients in Islamic Hospital Foundation (YARSI) Pontianak. *Jurnal ProNers*, 5(1), 2020. <https://doi.org/10.26418/jpn.v5i1.42911>

Lukman, & Ningsih, N. (2012). *Asuhan Keperawatan pada Klien dengan gangguan sistem muskuloskeletal*. Salemba Medika.

Mahaldar, A. R. (2014). Restless leg syndrome in chronic kidney disease. *Clinical Queries: Nephrology*, 3(1), 5–8. <https://doi.org/10.1016/j.cqn.2014.03.002>

Mahrova, A., & Svagrova, K. (2013). *Exercise therapy additional tool for managing physical and psychological problems on hemodialysis*. Intech Open Science. <https://doi.org/10.5772/53058>

Muna, U. W. (2022). Gambaran kejadian fatigue pada pasien chronic kidney disease di Rumah Sakit Islam Sultan Agung Semarang. *Jurnal Universitas Islam Sutan Agung*. <https://doi.org/10.465/26928>

Nur, A., Erika, K. A., & Sinrang, A. W. (2018). The effect of intradialysis stretching exercise on the scale of restless leg syndrome. *Journal of Islamic Nursing*, 3(2), 16. <https://doi.org/10.24252/join.v3i2.3673>

Nurhalimah. (2016). *Keperawatan Jiwa* (Vol. 1). Kementrian Kesehatan Republik Indonesia.

Nurhusna, Mekeama, L., & Sulistiawan, A. (n.d.). Analisis determinan faktor yang berhubungan dengan restless legs sindrom pada penderita chronic kidney disease end stage di RSUD Raden Mattaher Jambi. *Jambi Medical Jurnal*.



<https://doi.org/10.22437/jmj.v9i0001.12896>

Pardede, S. O., Yulman, A. R., & Andriana, J. (2016). Komplikasi neurologi penyakit ginjal konik pada anak. *Majalah Kedokteran*, 32(4), 189–195.

<https://doi.org/10.33541/mkvol34iss2pp60>

Perhimpunan Nefrologi Indonesia. (2013). *Konsensus Transplantasi Ginjal 2013 Vol I*. In Perhimpunan Nefrologi Indonesia

Pernefri. (2003). *Konsensus Dialisis*. Perhimpunan Nefrologi Indonesia.

Perry, A. G., & Potter, P. A. (2017). Fundamentals of nursing. In *Key to Success Staff Nurses Recruitment Exam*. [https://doi.org/10.5005/jp/books/12954\\_32](https://doi.org/10.5005/jp/books/12954_32)

Powe, N. R. (2019). CKD progression: Teasing out contributions of elements in the human exposome. *American Journal of Kidney Diseases*, 73(5), 581–582. <https://doi.org/10.1053/j.ajkd.2019.01.018>

Price, S. A., & Wilson, L. M. (2012). *Patofisiologi, konsep klinis proses-proses penyakit* (6th ed., Vol. 2). EGC.

Rady, S. E., Yousef, S. G., Elrefaey, S. R. I., & Abdelhady, M. S. (2020). Effect of muscle stretching and range of motion exercises on sleep quality and anxiety among hemodialysis patients. *Egyptian Journal of Health Care*, 11(4), 582–601. <https://doi.org/10.21608/ejhc.2020.156947>

Rahayu, G., Malini, H., & Oktarina, E. (2019). Hubungan kadar ureum terhadap restless legs syndrome pada pasien chronic kidney disease. *NERS Jurnal Keperawatan*, 15(2), 140. <https://doi.org/10.25077/njk.15.2.140-146.2019>

Rahimipour, M., Shahgholian, N., & Yazdani, M. (2015). Effect of hope therapy on depression, anxiety, and stress among the patients undergoing hemodialysis. *Iranian Journal of Nursing and Midwifery Research*, 20(6),

694–699. <https://doi.org/10.4103/1735-9066.170007>

Rahmawati, Hasanuddin, F., & Mokodompit, N. A. (2019). Latihan rom pada pasien gagal ginjal kronik dalam pemenuhan aktivitas. *Jurnal Media Keperawatan: Politeknik Kesehatan Makassar*, 10(2), 85–91.

<https://doi.org/10.32382/jmk.v10i2.1299>

Rochmawati, E., Utomo, E. K., & Makiyah, S. N. N. (2022). Improving dialysis adequacy and quality of life in patients undergoing hemodialysis with twice a week range of motion exercise. *Therapeutic Apheresis and Dialysis*, 26(1), 140–146. <https://doi.org/10.1111/1744-9987.13701>

Rompas, A. B., Tangka, J., & Rotti, J. (2013). Hubungan kadar hemoglobin dengan kualitas tidur pasien penyakit ginjal kronik dipoli ginjal dan hypertension BLU RSUP Prof. dr.R.D Kandou Manado. *Jurnal Keperawatan Universitas Sam Ratulangi*, 1. <https://doi.org/10.35790/jkp.v1i1.2160>

Rosdiana, I., Yetty, K., & Sabri, L. (2014). Kecemasan dan lamanya waktu menjalani hemodialisis berhubungan dengan kejadian insomnia pada pasien gagal ginjal kronik. *Jurnal Keperawatan Indonesia*, 17(2), 2354–9203. <https://doi.org/10.7454/jki.v17i2.440>

Sakitri, G., Makiyah, N., & Khoiriyati, A. (2017). Pengaruh intradialytic exercise terhadap fatigue pasien hemodialisis RSUP dr Soeradji Tirtonegoro Klaten. *Media Publikasi Penelitian*, 15(1), 58–64. <https://doi.org/123456789/35034>

Salama, A. H., Zaki, M. mahmoud, Radwan, H. A., Elfiky, E. R., & Zayed, D. A. (2022). Effect of intradialytic exercise on fatigue, psychological distress, and biochemical findings among hemodialysis children. *Egyptian Journal of Health Care*, 13(4), 1443–1457. <https://doi.org/10.21608/ejhc.2022.272201>

- Sananta, T., Gunadharma, S., Sadeli, H. A., Bandiara, R., Gamayani, U., Wibisono, Y., & Kurniani, N. (2022). Gambaran konduksi saraf tepi neuropati asimptomatik pada penyakit ginjal kronis stadium 3 dan 4. *Neurona Majalah Medis Dan Neuroscience*, 39, 4–8. <https://doi.org/10.52386/neurona.v39i2.205>
- Setiadi, & Irawandi, D. (2020). *Keperawatan Dasar*. Indomedia Pustaka.
- Sevim, S., Dogu, O., Kalegasi, H., Aral, M., Metin, O., & Çamdeviren, H. (2004). Correlation of anxiety and depression symptoms in patients with restless legs syndrome: A population based survey. *Journal of Neurology, Neurosurgery and Psychiatry*, 75(2), 226–230. <https://doi.org/10.1101/1102.02.04>
- Shahid, A., Wilkinson, K., Marcu, S., & Shapiro, C. M. (2012). Fatigue severity scale. *Stop, That and One Hundred Other Sleep Scales*, 1–406. <https://doi.org/10.1007/978-1-4419-9893-4>
- Silva, S. F. da, Pereira, A. A., Silva, W. A. H. da, Simoes, R., Neto, B., & Resende, J. de. (2013). Physical therapy during hemodialyse in patients with chronic kidney disease. *Jornal Brasileiro de Nefrologia : 'orgão Oficial de Sociedades Brasileira e Latino-Americana de Nefrologia*, 35(3), 170–176. <https://doi.org/10.5935/0101-2800.20130028>
- Siregar, S. (2015). *Statistik Parametrik Untuk Penelitian Kuantitatif* (1st ed.). Bumi Aksara.
- Soliman, M., & Mohamed, H. (2015a). Effect of intradialytic exercise on fatigue, electrolytes level and blood pressure in hemodialysis patients: A randomized controlled trial. *Journal of Nursing Education and Practice*, 5(11). <https://doi.org/10.5430/jnep.v5n11p16>



Soliman, M., & Mohamed, H. (2015b). Effect of intradialytic exercise on fatigue, electrolytes level and blood pressure in hemodialysis patients: A randomized controlled trial. *Journal of Nursing Education and Practice*, 5(11), 16–28. <https://doi.org/10.5430/jnep.v5n11p16>

Song, Y. Y, Hu, R. J, Diao, Y. S, Chen, L., & Jiang, X. L. (2018). Effects of exercise training on restless legs syndrome, depression, sleep quality, and fatigue among hemodialysis patients: A systematic review and meta-analysis. *Journal of Pain and Symptom Management*, 55(4), 1184–1195. <https://doi.org/10.1016/j.jpainsymman.2017.12.472>

Stuart, G. . (2016). *Keperawatan Kesehatan Jiwa*. Elsever.

Stuart, G. ., & Laraia, M. . (2005). *Principles and Practice of Psychiatric Nursing* (8th ed.). Mosby.Inc.

Sudrajat, A. (2019). Pengaruh range of motion dan taichi intradialysis exercise terhadap adekuasi pasien hemodialisis di rumah sakit pemerintah kabupaten lebak. *Jurnal Universitas Indonesia*. <https://doi.org/10.1016/j.enfcli.2019.07.021>

Sudrajat, A., Yetti, K., & Waluyo, A. (2021). The effect of the range of motion exercises combined with tai chi intradialysis on the adequacy of hemodialysis in patients at lebak district hospital. *Enfermeria Clinica*, 31, S113–S116. <https://doi.org/10.1016/j.enfcli.2020.12.003>

Sulistini, R., Yetti, K., & Hariyati, R. T. S. (2012). Faktor Faktor yang Mempengaruhi Fatigue Pada Pasien yang Menjalani Hemodialisis. *Jurnal Keperawatan Indonesia*, 15(2), 75–82. <https://doi.org/10.7454/jki.v15i2.30>

Sulistyaningrum, D. P. (2017). Pengaruh range of motion (ROM) intradialisis

terhadap aktivitas fisik pasien hemodialisa yang mengalami fatigue. *Jurnal Universitas Indonesia*. <https://doi.org/10.35728/jmkik.v5i1.117>

Sumah, D. F. (2020). Dukungan keluarga berhubungan dengan kepatuhan pasien gagal ginjal kronik yang menjalani terapi hemodialisa di RSUD dr. M. Haulussy Ambon. *Jurnal Biosainstek*, 2(01), 81–86. <https://doi.org/10.52046/biosainstek.v2i01.351>

Sutriyawan, A. (2021). *Metodologi Penelitian Kedokteran dan kesehatan*. PT Refika Aditama.

Tanvir, S., Butt, G.-U.-D., Taj, R., & Pak, A. (2013). Prevalence of depression and anxiety in chronic kidney disease patients on hemodialysis sohil tanvir et al prevalence of depression and anxiety in chronic kidney disease patients on haemodialysis. *Institute of Medical Sciences*, 9(2), 64–67. <https://doi.org/67.2089/2100>

Tsunoda, K., Kitano, N., Kai, Y., Uchida, K., Kuchiki, T., Okura, T., & Nagamatsu, T. (2015). Prospective study of physical activity and sleep in middle-aged and older adults. *American Journal of Preventive Medicine*, 48(6), 662–673. <https://doi.org/10.1016/j.amepre.2014.12.006>

Widianti, A. T., Hermayanti, Y., & Kurniawan, T. (2017). Pengaruh latihan kekuatan terhadap restless legs syndrome pasien hemodialisis. *Jurnal Keperawatan Padjadjaran*, 5(1), 47–56. <https://doi.org/10.24198/jkp.v5n1.6>

Yaseen, M., Jarullah, F. A., Yaqoob, S., Shakeel, H. A., Maqsood, H., & Naveed, S. (2021). Association of quality of life, anxiety, and depression with restless leg syndrome in the hemodialysis patients. *BMC Research Notes*, 14(1), 1–12. <https://doi.org/10.1186/s13104-021-05701-w>

Zuber, J., Le Quintrec, M., Soussan, R. S., Loirat, C., Bacchi, F., Ve'Ronique, V., Legendre, C., Krid, S., Bertoye, C., Gueutin, V., Lahoche, A., Heyne, N., Ardissino, G., Chatelet, V., Noël, L. H., Hourmant, M., Niaudet, P., Frémeaux-Bacchi, V. V., Rondeau, E., ... Gallon, L. (2017). Long term migalastat treatment stabilizes renal function in patients with fabry disease: results from a phase 3 clinical study. *Nephrology Dialysis Transplantation*, 3252. <https://doi.org/10.3252/pso.eu.54ERA.2017>

