

DAFTAR PUSTAKA

1. Bagchi A, Zapol WM. The pathophysiology of critical illness. In: Longnecker DE, Mackey SC, Newman MF, Sandberg WS, Zapol WM, editors. *Anesthesiology*, 3e. New York, NY: McGraw-Hill Education; 2017:1225-9.
2. Cerro G, Checkley W. Global analysis of critical care burden. *The Lancet Respiratory Medicine*. 2014;2(5):343–4.
3. Garland A, Olafson K, Ramsey CD, Yogendran M, Fransoo R. Distinct determinants of long-term and short-term survival in critical illness. *Intensive Care Med*. 2014;40(8):1097–105.
4. Vincent JL, Marshall JC, Namendys-Silva SA, François B, Martin-Loeches I, Lipman J, et al. Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *The Lancet Respiratory Medicine*. 2014;2(5):380–6.
5. Annane D, Pastores SM, Arlt W, Balk RA, Beishuizen A, Briegel J, et al. Critical illness-related corticosteroid insufficiency (CIRCI): a narrative review from a multispecialty task force of the society of critical care medicine (SCCM) and the european society of intensive care medicine (ESICM). *critical care medicine*. 2017;45(12):2089–98.
6. Evans L, Rhodes A, Alhazzani W, Antonelli M, Coopersmith CM, French C, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Intensive Care Med*. 2021; 47(11):1181–247.
7. Fleischmann C, Scherag A, Adhikari NKJ, Hartog CS, Tsaganos T, Schlattmann P, et al. Assessment of global incidence and mortality of hospital-treated sepsis. current estimates and limitations. *Am J Respir Crit Care Med*. 2016;193(3):259–72.
8. Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the global burden of disease study. *The Lancet*. 2020; 395(10219):200–11.
9. World Health Organization. Global report on the epidemiology and burden of sepsis: current evidence, identifying gaps and future direction. Geneva: World Health Organization; 2020:16-28.

10. Dugar S, Choudhary C, Duggal A. Sepsis and septic shock: guideline-based management. *CCJM*. 2020; 87(1):53–64.
11. Jawad I, Lukšić I, Rafnsson SB. Assessing available information on the burden of sepsis: global estimates of incidence, prevalence and mortality. *J Glob Health*. 2012; 2(1):010404.
12. KMK. Keputusan menteri kesehatan republik indonesia tentang pedoman nasional pelayanan kedokteran tata laksana sepsis. 2017.
13. Hidayati, Arifin H. kajian penggunaan antibiotik pada pasien sepsis dengan gangguan ginjal. *Jurnal Sains Farmasi & Klinis*. 2016;2(2):129–37.
14. Pardede DKB, Siahaan D. Eosinopenia sebagai penanda diagnosis sepsis. *cermin dunia kedokteran*. 2014;41(10).
15. Lambden S, Laterre PF, Levy MM, Francois B. The SOFA score—development, utility and challenges of accurate assessment in clinical trials. *Crit Care*. 2019; 23(1):374.
16. Marino LO, Souza HP. Dysfunction of the hypothalamic-pituitary-adrenal axis in critical illness: a narrative review for emergency physicians. *European Journal of Emergency Medicine*. 2020; 27(6):406–13.
17. Marik PE. *Handbook of evidence-based critical care*. New York, NY: Springer New York; 2010:427-33.
18. The Corticosteroid Guideline Task Force of SCCM and ESICM, Pastores SM, Annane D, Rochweg B. Guidelines for the diagnosis and management of critical illness-related corticosteroid insufficiency (CIRCI) in critically ill patients (Part II): society of critical care medicine (SCCM) and european society of intensive care medicine (ESICM) 2017. *Intensive Care Med*. 2018; 44(4):474–7.
19. University of the Philippines Manila-Philippine General, Arcellana AE, Lim KW, Arcegon M, Jimeno C. The Development of a protocol for critical illness-related corticosteroid insufficiency (CIRCI) at a tertiary hospital. *JAFES*. 2022; 37(1):14–23.
20. Kromah F, Tyroch A, McLean S, Hughes H, Flavin N, Lee S. Relative adrenal insufficiency in the critical care setting: debunking the classic myth. *World J Surg*. 2011; 35(8):1818–23.

21. Téblick A, Peeters B, Langouche L, Van den Berghe G. Adrenal function and dysfunction in critically ill patients. *Nat Rev Endocrinol*. 2019; 15(7):417–27.
22. Salgado D, Verdeal J, Rocco J. Adrenal function testing in patients with septic shock. *Crit Care*. 2006;10(5):R149.
23. Song JH, Kim JH, Lee SM, Lee J. Prognostic implication of adrenocortical response during the course of critical illness. *Acute Crit Care*. 2019; 34(1):38–45.
24. Annane D. Cortisol replacement for severe sepsis and septic shock: what should I do? *Crit Care*. 2002; 6(3):190.
25. Goodman S, Sprung CL, Ziegler D, Weiss YG. Cortisol changes among patients with septic shock and the relationship to ICU and hospital stay. *Intensive Care Med*. 2005; 31(10):1362–9.
26. Alzahrani AS, Mukhtar N, Aljomaiah A, Aljamei H, Bakhsh A, Alsudani N, et al. The impact of covid-19 viral infection on the hypothalamic-pituitary-adrenal axis. *Endocrine Practice*. 2021; 27(2):83–9.
27. Longnecker DE, Newman MF, Mackey S, Sandberg WS, Zapol WM. *Anesthesiology*. 2017.1131-36
28. Jackson M, Cairns T. *Care of the critically ill patient*. Surgery (Oxford). 2021; 39(1):29–36.
29. DeMerle KM, Angus DC, Baillie JK, Brant E, Calfee CS, Carcillo J, et al. Sepsis subclasses: a framework for development and interpretation. *Crit Care Med*. 2021; 49(5):748–59.
30. Kobbe P, Bläsius FM, Lichte P, Oberbeck R, Hildebrand F. Neuroendocrine modulation of the immune response after trauma and sepsis: does it influence outcome? *JCM*. 2020; 9(7):2287.
31. Yang Y, Liu L, Jiang D, Wang J, Ye Z, Ye J, et al. Critical illness–related corticosteroid insufficiency after multiple traumas: A multicenter, prospective cohort study. *Journal of Trauma and Acute Care Surgery*. 2014; 76(6):1390–6.
32. The WHO Rapid Evidence Appraisal for COVID-19 Therapies (REACT) Working Group, Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, et al. Association between administration of systemic corticosteroids and mortality

among critically ill patients with covid-19: a meta-analysis. *jama*. 2020; 324(13):1330.

33. Jameson JL. Mechanisms of hormone action. In: Jameson JL, Fauci AS, Kasper DL, Hauser SL, Longo DL, Loscalzo J, editors. *Harrison's Principles of Internal Medicine*, 20e. New York, NY: McGraw-Hill Education; 2018; 2653-9.
34. Arlt W. Disorders of the adrenal cortex. In: Jameson JL, Fauci AS, Kasper DL, Hauser SL, Longo DL, Loscalzo J, editors. *Harrison's Principles of Internal Medicine*, 20e. New York, NY: McGraw-Hill Education; 2018; 2719-38.
35. Melmed S, Auchus RJ, Goldfine AB, Koenig RJ, Rosen CJ. *Williams textbook of endocrinology*. 14th ed. Philadelphia: Elsevier, Inc; 2019:480-99.
36. Fang F, Zhang Y, Tang J, Lunsford LD, Li T, Tang R, et al. Association of corticosteroid treatment with outcomes in adult patients with sepsis: a systematic review and meta-analysis. *JAMA Intern Med*. 2019; 179(2):213–23.
37. Wu YP, Lin CK, Hamaya R, Huang FY, Chien YS, Hsu YT, et al. Intravenous corticosteroid treatment in adult patients with sepsis defined by the Sepsis-3 criteria: a systematic review and meta-analysis. *Intensive Care and Critical Care Medicine*; 2021.
38. Liang H, Song H, Zhai R, Song G, Li H, Ding X, et al. Corticosteroids for Treating Sepsis in Adult Patients: A Systematic Review and Meta-Analysis. *Front Immunol*. 2021; 12:709155.
39. Sprung CL, Annane D, Keh D, Moreno R, Singer M, Freivogel K, et al. Hydrocortisone therapy for patients with septic shock. *N Engl J Med*. 2008; 358(2):111–24.
40. Roquilly A. Hydrocortisone therapy for patients with multiple trauma: the randomized controlled hypolyte study. *JAMA*. 2011; 305(12):1201.
41. Annane D, Bellissant E, Bollaert PE, Briegel J, Keh D, Kupfer Y, et al. Corticosteroids for treating sepsis in children and adults. *Cochrane Emergency and Critical Care Group*, editor. *Cochrane Database of Systematic Reviews*. 2019

42. Park HY, Suh GY, Song JU, Yoo H, Jo IJ, Shin TG, et al. Early initiation of low-dose corticosteroid therapy in the management of septic shock: a retrospective observational study. *Crit Care*. 2012;16(1).
43. Katsenos CS, Antonopoulou AN, Apostolidou EN, Ioakeimidou A, Kalpakou GT, Papanikolaou MN, et al. Early administration of hydrocortisone replacement after the advent of septic shock: impact on survival and immune response. *Crit Care Med*. 2014; 42(7):1651–7.
44. Hall JE. *Guyton and hall textbook of medical physiology*. 14th ed. Philadelphia: Elsevier; 2020. 1570-9
45. Barrett KE, Barman SM, Brooks HL, Yuan JXJ. The adrenal medulla & adrenal cortex. In: *Ganong's Review of Medical Physiology*, 26e. New York, NY: McGraw-Hill Education; 2019. 1563-72.
46. Turpeinen U, Hämäläinen E. Determination of cortisol in serum, saliva and urine. *Best Practice & Research Clinical Endocrinology & Metabolism*. 2013; 27(6):795–801.
47. Rifai N. *Tietz Textbook of clinical chemistry and molecular diagnostics - E-Book*. Saintt Louis: Elsevier Health Sciences; 2017. 1442-47
48. Mignot-Evers L, Raaijmakers V, Buunk G, Brouns S, Romano L, van Herpt T, et al. Comparison of SIRS criteria and qSOFA score for identifying culture-positive sepsis in the emergency department: a prospective cross-sectional multicentre study. *BMJ Open*. 2021;11(6).
49. Kilinc Toker A, Kose S. Comparison of SOFA Score, SIRS, qSOFA, and qSOFA + L criteria in the diagnosis and prognosis of sepsis. *Eurasian J Med*. 2021; 53(1):40–7.
50. Lakbar I, Einav S, Lalevée N, Martin-Loeches I, Pastene B, Leone M. Interactions between gender and sepsis—implications for the future. *Microorganisms*. 2023; 11(3):746.
51. Nasir N, Jamil B, Siddiqui S, Talat N, Khan FA, Hussain R. Mortality in sepsis and its relationship with gender. *Pak J Med Sci*. 2015;31(5):1201–6.
52. Kweon OJ, Choi JH, Park SK, Park AJ. Usefulness of presepsin (sCD14 subtype) measurements as a new marker for the diagnosis and prediction of

- disease severity of sepsis in the Korean population. *Journal of Critical Care*. 2014; 29(6):965–70.
53. Klouche K, Cristol JP, Devin J, Gilles V, Kuster N, Larcher R, et al. Diagnostic and prognostic value of soluble CD14 subtype (Presepsin) for sepsis and community-acquired pneumonia in ICU patients. *Ann Intensive Care*. 2016; 6(1):59.
54. Rabee HA, Tanbour R, Nazzal Z, Hamshari Y, Habash Y, Anaya A, et al. Epidemiology of sepsis syndrome among intensive care unit patients at a tertiary university hospital in palestine in 2019. *Indian Journal of Critical Care Medicine*. 2020; 24(7):551–6.
55. Joy AP, Murali AB, Joshi MA, Parambil JC. Absolute eosinophil count as a diagnostic and prognostic marker compared to C- reactive protein and Procalcitonin in patients with sepsis. *Clinical Epidemiology and Global Health*. 2020; 8(2):632–6.
56. Jeon K, Na SJ, Oh DK, Park S, Choi EY, Kim SC, et al. Characteristics, management and clinical outcomes of patients with sepsis: a multicenter cohort study in Korea. *Acute Crit Care*. 2019; 34(3):179–91.
57. Nugent K, Edriss H, Selvan K. Hyperglycemia and outcomes in patients with sepsis. *J Thorac Dis*. 2016; 8(7): E575–7.
58. Ganesh K, Sharma R, Varghese J, Pillai MGK. A profile of metabolic acidosis in patients with sepsis in an intensive care unit setting. *Int J Crit Illn Inj Sci*. 2016; 6(4):178.
59. Dupuis C, Bouadma L, Ruckly S, Perozziello A, Van-Gysel D, Mageau A, et al. Sepsis and septic shock in France: incidences, outcomes and costs of care. *Ann Intensive Care*. 2020; 10(1):145.
60. Chauvet JL, El-Dash S, Delastre O, Bouffandeau B, Jusserand D, Michot JB, et al. Early dynamic left intraventricular obstruction is associated with hypovolemia and high mortality in septic shock patients. *Crit Care*. 2015; 19(1):262.
61. Vucelić V, Klobučar I, Đuras-Cuculić B, Gverić Grginić A, Prohaska-Potočnik C, Jajić I, et al. Sepsis and septic shock – an observational study of

the incidence, management, and mortality predictors in a medical intensive care unit. *Croat Med J.* 2020; 61(5):429–39.

62. Gonçalves-Pereira J, Pereira JM, Ribeiro O, Baptista JP, Froes F, Paiva JA. Impact of infection on admission and of the process of care on mortality of patients admitted to the intensive care unit: the INFAUCI study. *Clinical Microbiology and Infection.* 2014; 20(12):1308–15.
63. Shankar-Hari M, Phillips GS, Levy ML, Seymour CW, Liu VX, Deutschman CS, et al. Developing a new definition and assessing new clinical criteria for septic shock: for the third international consensus definitions for sepsis and septic shock (sepsis-3). *JAMA.* 2016; 315(8):775.
64. Chatterjee S, Bhattacharya M. Epidemiology of adult-population sepsis in India: A single center 5 year experience. *Indian Journal of Critical Care Medicine.* 2017; 21(9):573–7.
65. Paoli CJ, Reynolds MA, Sinha M, Gitlin M, Crouser E. Epidemiology and costs of sepsis in the united states—an analysis based on timing of diagnosis and severity level*: *critical care medicine.* 2018 ; 46(12):1889–97.
66. Dewi RS, Radji M, Andalusia R. Evaluation of Antibiotic Use Among Sepsis Patients in an Intensive Care Unit: A cross-sectional study at a referral hospital in Indonesia. *Sultan Qaboos Univ Med J.* 2018; 18(3):367.
67. Do SN, Dao CX, Nguyen TA, Nguyen MH, Pham DT, Nguyen NT, et al. Sequential Organ Failure Assessment (SOFA) Score for predicting mortality in patients with sepsis in Vietnamese intensive care units: a multicentre, cross-sectional study. *BMJ Open.* 2023; 13(3):e064870.
68. Liu C, Suo S, Luo L, Chen X, Ling C, Cao S. SOFA Score in relation to Sepsis: Clinical Implications in Diagnosis, Treatment, and Prognostic Assessment. Chen G, editor. *Computational and Mathematical Methods in Medicine.* 2022; 2022:1–8.
69. Yébenes JC, Ruiz-Rodríguez JC, Ferrer R, Clèries M, Bosch A, et al. Epidemiology of sepsis in Catalonia: analysis of incidence and outcomes in a European setting. *Ann Intensive Care.* 2017; 7(1):19.

70. Bouza C, López-Cuadrado T. Epidemiology and Trends of Sepsis in Young Adults Aged 20–44 Years: A Nationwide Population-Based Study. *JCM*. 2019; 9(1):77.
71. Sinapidis D, Kosmas V, Vittoros V, Koutelidakis IM, Pantazi A, Stefos A, et al. Progression into sepsis: an individualized process varying by the interaction of comorbidities with the underlying infection. *BMC Infect Dis*. 2018; 18(1):242.
72. Whiles B, Deis A, Miller P, Simpson S. Comorbid Conditions Predict Outcomes in Patients With Severe Sepsis. *Chest*. 2016; 149(4):A170.
73. Umemura Y, Ogura H, Takuma K, Fujishima S, Abe T, Kushimoto S, et al. Current spectrum of causative pathogens in sepsis: A prospective nationwide cohort study in Japan. *International Journal of Infectious Diseases*. 2021; 103:343–51.
74. Leligdowicz A, Dodek PM, Norena M, Wong H, Kumar A, Kumar A. Association between Source of Infection and Hospital Mortality in Patients Who Have Septic Shock. *Am J Respir Crit Care Med*. 2014; 189(10):1204–13.
75. Abe T, Ogura H, Kushimoto S, Shiraishi A, Sugiyama T, et al. Variations in infection sites and mortality rates among patients in intensive care units with severe sepsis and septic shock in Japan. *J Intensive Care*. 2019; 7(1):28.
76. Peeters B, Langouche L, Van Den Berghe G. Adrenocortical Stress Response during the Course of Critical Illness. In: Terjung R, editor. *Comprehensive Physiology*. 1st ed. Wiley; 2017. p. 283–98.
77. Peeters B, Meersseman P, Vander Perre S, Wouters PJ, Vanmarcke D, Debaveye Y, et al. Adrenocortical function during prolonged critical illness and beyond: a prospective observational study. *Intensive Care Med*. 2018; 44(10):1720–9.
78. Annane D, Pastores SM, Arlt W, Balk RA, Beishuizen A, Briegel J, et al. Critical Illness-Related Corticosteroid Insufficiency (CIRCI): A Narrative Review from a Multispecialty Task Force of the Society of Critical Care Medicine (SCCM) and the European Society of Intensive Care Medicine (ESICM). *Critical Care Medicine*. 2017; 45(12):2089–98.

79. Arcellana AE, Lim KW, Arcegon M, Jimeno C. Critical Illness-Related Corticosteroid Insufficiency (CIRCI) among Patients with COVID-19 at a Tertiary Hospital: Clinical Characteristics and Outcomes. *JAFES*. 2023; 38(1):90–9.
80. Levy-Shraga Y, Pinhas-Hamiel O. Critical Illness-Related Corticosteroid Insufficiency in Children. *Horm Res Paediatr*. 2013;80(5):309–17.
81. Ok YJ, Lim JY, Jung SH. Critical Illness-Related Corticosteroid Insufficiency in Patients with Low Cardiac Output Syndrome after Cardiac Surgery. *Korean J Thorac Cardiovasc Surg*. 2018;51(2):109–13.
82. Odom DC, Gronowski AM, Odom E, Clutter W, Thaelke M. A Single, Post-ACTH Cortisol Measurement to Screen for Adrenal Insufficiency in the Hospitalized Patient. *J Hosp Med*. 2018.
83. Kawahara T, Tsuji M, Tominaga N, Toyama N, Toda M. Frequency of Adrenal Insufficiency in Patients With Hypoglycemia in an Emergency Department: A Cross-sectional Study. *Journal of the Endocrine Society*. 2022;6:bvac119.
84. Alabdrabnabi FM, Saeed ZAAL, Elamin YA. Hypoglycemia As Initial Presentation In Patient With Isolated Acth Deficiency. *AACE Clinical Case Reports*. 2020;6:e338–41.
85. Ammar MA, Ammar AA, Wieruszewski PM, Bissell BD, T. Long M, Albert L, et al. Timing of vasoactive agents and corticosteroid initiation in septic shock. *Ann Intensive Care*. 2022;12:47.