

DAFTAR PUSTAKA

- Abbas, B., Tjoli, I., & Munarti. (2020). Genetic diversity of sago palm (*Metroxylon sagu*) accessions based on plastid cpDNA matK gene as DNA barcoding. *BIODIVERSITAS*, 21(1), 219-225.
- Abdullah, M. A., Rahmah, A. U., & Man, Z. (2010). Physicochemical and sorption characteristics of Malaysian *Ceiba pentandra* (L.) Gaertn. as a natural oil sorbent. *Journal of Hazardous Materials*, 177, 683-691.
- Abengmeneng, C. S., Ofori, D., Kumapley, P., Akromah, R., Jamnadass, R., & Quain, M. (2016). Genetic relationship among 36 genotypes of *Ceiba pentandra* (L.) as revealed by RAPD and ISSR markers. *American Journal of Agriculture and Forestry*, 4(4), 86-96.
- Agoes, A. (2010). *Tanaman Obat Indonesia*. Jakarta: Salemba Medika.
- Almerekova S., Mukhitdinov N., & Abugalieva, S. (2017). Phylogenetic study of the endemic species *Oxytropis almaatensis* (Fabaceae) based on nuclear ribosomal DNA ITS sequences. *BMC Plant Biology*, 17(1), 173.
- Alvarado, C. R., Alvarado, C. A., & Mendoza, O. O. (2002). *Ceiba pentandra* (L.) Gaertn. In: J. A. Vozzo (Ed.), *Tropical tree seed manual* (pp. 394-396). Washington DC, United States: USDA, Forest Service Publication.
- Alvarez, I., & Wendel, J. F. (2003). Ribosomal ITS sequences and plant phylogenetic inference. *Molecular Phylogenetics and Evolution*, 29, 417-434.
- Aneklaphakij, C., Bunsupa, S., Sirichamorn, Y., Bongcheewin, B., & Satitpatipan, V. (2020). Taxonomic notes on the 'Mahat' (*Artocarpus lacucha* and *A. thailandicus*, Moraceae) species complex in Thailand. *Plants*, 9(3), 391.
- Azani N., Bruneau A., Wojciechowski M. F., & Zarre S. (2017). Molecular phylogenetics of annual *Astragalus* (Fabaceae) and its systematic implications, *Botanical Journal of the Linnean Society*, 184(3), 347-365.
- Azuma, H., Thien, L. B., & Kawano, S. (1999). Molecular phylogeny of Magnolia (Magnoliaceae) inferred from cpDNA sequences and evolutionary divergence of the floral scents. *Journal of Plant Research*, 112, 291-306.
- Badan Pusat Statistik (BPS). (2020). *Produksi Tanaman Perkebunan Rakyat (Ton)*.
- Badotti, F., de Oliveira, F. S., Garcia, C. F., Vaz, A. B. M., Fonseca, P. L. C., Nahum, L. A., Oliveira, G., & Goes-Neto, A. (2017). Effectiveness of ITS and sub-regions as DNA barcode markers for identifying Basidiomycota (Fungi). *BMC Microbiology*, 17(42), 1-12.

- Baker, H. G. (1955). Self-compatibility and establishment after 'long-distance' dispersal. *Evolution*, 9(3), 347-349.
- Baker, H. G., & Harris, B. J. (1959). Bat-pollination of the silk-cotton tree, *Ceiba pentandra* (L.) Gaertn. (*sensu lato*), in Ghana. *Journal of the West African Science Association*, 4, 1-9.
- Baldwin, B. G. (1992). Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: an example from the compositae. *Molecular Phylogenetics Evolution*, 1(1), 3-16.
- Baldwin, B. G., Sanderson, M. J., Porter, J. M., Wojciechowski, M. F., Campbell, C. S., & Donoghue, M. J. (1995). The ITS region of nuclear ribosomal DNA: A valuable source of evidence on angiosperm phylogeny. *Annals of the Missouri Botanical Garden*, 82(2), 247-277.
- Bandelt, H. J., Forster, P., & Rohl, A. (1999). Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology Evolution*, 16(1), 37-48.
- Bates, D. M. (2004). *Malvales*. Encyclopedia Britannica.
- Bayer, C., Fay, M. F., De Bruijn, A. Y., Savolainen, V., Morton, C. M., Kubitzki, K., Alverson, W. S., & Chase, M. W. (1999). Support for an expanded family concept of Malvaceae within a circumscribed order Malvales: a combined analysis of plastid *atpB* and *rbcL* DNA sequences. *Botanical Journal of the Linnean Society*, 129(4), 267-303.
- Beeck, M., Lievens, B., Busschaert, P., Declerck, S., Vangronsveld, J., & Colpaert, J. V. (2014). Comparison and validation some ITS primer pairs useful for fungal metabarcoding studies. *PLOS ONE*, 9(6).
- Bellarosa, R., Simeone, M. C., Papini, A., & Schirone, B. (2005). Utility of ITS sequence data for phylogenetic reconstruction of Italian *Quercus* spp. *Molecular Phylogenetics and Evolution*, 34(2), 355-370.
- Bellemain, E., Carlsen, T., Brochmann, C., Coissac, E., Taberlet, P., & Kuserud, H. (2010). ITS as an environmental DNA barcode for fungi: an in-silico approach reveals potential PCR biases. *BMC Microbiology*, 10, 189.
- Brondani, R. P. V., Gaiotto, F. A., Missiaggia, A. A., Kirst, M., Gribel, R., & Grattapaglia, D. (2003). Microsatellite markers for *Ceiba pentandra* (Bombaceae), an endangered tree species of the Amazon Forest. *Molecular Ecology*, 3, 177-179.
- Bryois, J., Buil, A., Ferreira, P. G., Panousis, N. I., Brown, A. A., Vinuela, A., Planchon, A., Bielser, D., Small, K., Spector, T., & Dermitzakis, E. T. (2017). Time-dependent genetic effects on gene expression implicate aging processes. *Genome Research*, 27(4), 545-552.

- CBOL Plant Working Group. (2009). A DNA barcode for land plants. *Proceedings of the National Academy of Sciences*, 106, 12794-12797.
- Chan, E. W. C., Yeong, S. W., Wong, C. W., Soo, O. Y. M., Phua, A. C. Y., & Ng, Y. K. (2023). *Ceiba pentandra* (L.) Gaertn.: An overview of its botany, uses, reproductive biology, pharmacological properties, and industrial potentials. *Journal of Applied Biology & Biotechnology*, 11(1), 1-7.
- Chase, M. W., & Fay, M. F. (2009). Barcoding of plants and fungi. *Science*, 325, 682-683.
- Chen, S., Yao, H., Han, J. (2010). Validation of the ITS2 region as a novel DNA barcode for identifying medicinal plant species. *PLoS ONE*, 5, e8613.
- Cheng, Z., Lu, B., Baldwin, B. S., Sameshima, K. and Chen, J. 2001. Comparative studies of genetic diversity in kenaf (*Hibiscus cannabinus* L.) varieties based on analysis of agronomic and RAPD data. *Hereditas*, 136, 231-232.
- Choi, J. H., Jeong, D. G., Oh, J. N., Kim, S., Lee, Y. H., UngChoi, Y., Myoung, J. G., & Kim, C. G. (2020). DNA barcoding of coral reef fishes from Chuuk State, Micronesia. *Mitochondrial DNA B: Resour*, 5(3), 201-217.
- Church, G. M. (2006). Genomes for all. *Scientific American*, 294, 46-54.
- Cox, P. A. (1983). Observations on the natural history of Samoan bats. *Mammalia*, 47, 519-521.
- Deng, Y., Liu, T., Xie, Y., Wei, Y., Xie, Z., Shi, Y., & Deng, X. (2020). High genetic diversity and low differentiation in *Michelia shiluensis*, an endangered magnolia species in South China. *Forests*, 11, 1-15.
- Dick, C. W., Bermingham, E., Lemes, M. R., & Gribel, R. (2007). Extreme long-distance of the lowland tropical rainforest tree *Ceiba pentandra* L. (Malvaceae) in Africa and the Neotropics. *Molecular Ecology*, 16, 3039-3049.
- Dkhar, J., Kumaria, S., Rama, Rao, S., & Tandon, P. (2012). Sequence characteristics and phylogenetic implications of the nrDNA internal transcribed spacers (ITS) in the genus *Nymphaea* with focus on some Indian representatives. *Plant Systematics and Evolution*, 298, 93-108.
- Doh E. J., Paek S. H., Lee G., Lee M. Y., and Oh S. E. (2016). Application of partial internal transcribed spacer sequences for the discrimination of *Artemisia capillaris* from other *Artemisia* species. *Evidence based complementary and alternative medicine*, 2016, 7043436.
- Doyle, J. J., & Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin*, 19(1), 11-15.

- Duarte, M. C., Esteves, G. L., Salatino, M. L. F., Walsh, K. C., & Baum, D. A. (2011). Phylogenetic analyses of *Eriotheca* and related genera (Bombacoideae, Malvaceae). *Systematic Botany*, 36(3), 690-701.
- Duke. (2008). *Duke's handbook of medical plants of Latin America*. CRC Press.
- El-Metwally, S., Ouda, O. M., & Helmy, M. (2014). New horizons in next-generation sequencing. In: *Next-generation sequencing technologies sand challenges in sequence assembly* (pp. 51-59).
- Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetic*, 17(7), 422-433.
- Elmqvist, T., Cox, P. A., Rainey, W. E., & Pierson, E. D. (1992). Restricted pollination on oceanic islands: Pollination of *Ceiba pentandra* by flying foxes in Samoa. *Biotropica*, 24(1), 15-23.
- Elumalai, A., Mathani, N., Didala, A., Kasarla, R., & Venkatesh, Y. (2012). A review on *Ceiba pentandra* and its medical features. *Asian Journal Pharmaceutical Technology*, 2, 83-86.
- Enaberuel, L. O., Obisesan, I. O., Okolo, E. C., Akinwale, R. O., Aisueni, N. O., & Ataga, C. D. (2014). Genetic diversity of shea butter tree (*Vitellaria paradoxa* C. F. Gaertn.) in the Guinea Savanna of Nigeria based on morphological markers. *American-Eurasian Journal Agriculture & Environment Sciences*, 14(7), 615-623.
- Enguito, R. Z. C., & Novero, A. U. (2018). Genetic variation of sago palm (*Metroxylon sagu* Rottb.) samples from mindano, philippines using internal transcribed spacer analysis. *Malaysian Applied Biology*, 47(3), 39-48.
- Eren, K., Taktakoglu, N., & Pirim, I. (2022) DNA sequencing methods: from past to present. *The Eurasian Journal of Medicine*, 54, 47-56.
- Fajarningsih, N. D. (2016). Internal transcribed spacer (ITS) as DNA barcoding to identify fungal species: a review. *Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology*, 11(2), 37-44.
- Fama, P., Olsen, J.J., Stam, W.T., & Procaccini, G. (2000). High levels of intra- and inter individual polymorphism in the rDNA ITS1 of *Caulerpa racemosa* (Chlorophyta). *European Journal of Phycology*, 35, 349-356.
- Fan, X., Liu, J., Sha, L. N., Sun, G. L., Hu, Z. Q., Zeng, J., Kang, H. Y., Zhang, H. Q., Wang, Y., Wang, X., Zhang, L., Ding, C. B., Yang, R. W., Zheng, Y. L., & Zhou, Y. H. (2014). Evolutionary pattern of rDNA following polypoidy in *Leymus* (Triticeae: Poaceae). *Molecular Phylogenetics and Evolution*, 77, 296-306.

- Fang, G., Hammar, S., & Grumet, R. (1992). A quick and inexpensive method for removing polysaccharides from plant genomic DNA. *Biofeedback*, 13, 52-53.
- Frankham, R., Ballou, J. D., & Briscoe, D. A. (2002). *Introduction to conservation genetics*. Cambridge, UK: Cambridge University Press.
- Freitas, A. D. S., Silva, M. L. R. B., Mergulhao, A. C. E. S., & Lyra, M. C. C. P. (2013). The genetic variability using sequencing of the ribosomal internal transcribed spacer (ITS) region in cultivars of the cowpea [*Vigna unguiculata* L. (Walp)]. *African Journal of Agricultural Research*, 8(32), 4365-4373.
- Furlan, E. J., Stoklosa, J., Griffith, N., Gust, R., Ellis, R. M., Huggins, & A. R. Weeks. (2011). Small population size and extremely low levels of genetic diversity in island populations of the platypus, *Ornithorhynchus anaticus*. *Ecology and Evolution*, 844-857.
- Gao, T., Yao, H., Song, J., Liu, C., Zhu, Y., Ma, X., Pang, X., Xu, H., & Chen, S. (2010). Identification of medicinal plants in the family Fabaceae using a potential DNA barcode ITS2. *Journal of Ethnopharmacology*, 130, 116-121.
- Garcia-Robledo, C., Erickson, D. L., Staines, C. L., Erwin, T. L., & Kress, W. J. (2013). Tropical plant-herbivore networks: reconstructing species interactions using DNA barcodes. *PLOS ONE*, 8.
- Gawali, A. (2014). Litterfall and carbon, nutrient returns in stands of *Ceiba pentandra* (L.) Gaertn. sub-humids tropics of Eastern India. *International Journal for Crop Improvement*, 5, 1-8.
- Gawali, A., Putri, S., & Swamy, S. L. (2015). Evaluation growth and yield of wheat varieties under *Ceiba pentandra* (L.) based agrisilviculture system. *Universal Journal of Agricultural Research*, 3, 173-181.
- GBIF. 2023. GBIF Backbone Taxonomy *Ceiba pentandra* (L.) Gaertn. Diakses pada 20 April 2023, dari <https://www.gbif.org/species/5406697>.
- Gentry, A. H. (1993). Diversity and floristic composition of lowland tropical forest in Africa and South America. In: P. Goldblatt, *Biogeography of Africa and South America* (pp. 500-546). New Haven: Yale University Press.
- Ghada, B., Ahmed, B. A., Messaoud, M., & Amel, S. H. (2013). Genetic diversity and molecular evolution of the internal transcribed spacer (ITSs) of nuclear ribosomal DNA in the Tunisian fig cultivars (*Ficus carica* L.; Moraceae). *Biochemical Systematics and Ecology*, 48, 20-33.
- Grattapaglia, D. (2007). Marker-assisted selection in Eucalyptus. In: E. P. Guimaraes, J. Ruane, B. D. Scherf, A. Sonnino, J. D. Dargie (Ed.), *Current*

status and future perspective in crops, livestock, forestry, and fish (pp. 252-281). Rome: Food and Agriculture Organization of the United Nations.

Gribel, R., Gibbs, P. E., & Queiroz, A. L. (1999). Flowering phenology and pollination biology of *Ceiba pentandra* (Bombaceae) in Central Amazonia. *Journal of Tropical Ecology*, 15, 247-263.

Gudbjartsson, D. F., Helgason, H., Gudjonsson, S. A., Zink, F., Oddson, A., Gylfason, A., Besenbacher, S., Magnusson, G., Halldorsson, B. V., Hjartarson, E., Sigurdsson, G. T., Stacey, S. N., Frigge, M. L., Holm, H., Saemundsdottir, J., Helgadottir, H. T., Johannsdottir, H., Sigfusson, G., Thorgeirsson, G., Sverrisson, J. T., Gretarsdottir, S., Walters, G. B., Rafnar, T., Thjodleifsson, B., Bjornsson, E. S., Olafsson, S., Thorarinsdottir, H., Steingrimsdottir, T., Gudmundsdottir, T. S., Theodors, A., Jonasson, J. G., Sigurdsson, A., Bjornsdottir, G., Jonsson, J. J., Thorarensen, O., Ludvigsson, P., Gudbjartsson, H., Eyjolfsson, G. I., Sigurdardottir, O., Olafsson, I., Arnar, D. O., Magnusson, O. T., Kong, A., Masson, G., Thorsteinsdottir, U., Helgason, A., Sulem, P., & Stefansson, K. (2015). Large-scale whole-genome sequencing of the Icelandic population. *Nature Genetics*, 47(5), 435-444.

Hafizah, R. A. Adawiyah, R., Harahap, R. F., Hannum, S. Santoso, P. C. 2017. Aplikasi marka ISSR pada keanekaragaman genetik durian (*Durio zibethinus* Murr.) di Kabupaten Deli Serdang, Sumatera Utara. *Journal of Biology*, 11(1), 49-56.

Hall, B. G. (2001). *Phylogenetic trees made easy: a how - to manual for molecular biologist*. USA: Sinauer Associates, Inc.

Hall, N. (2007). Advanced sequencing technologies and their wider impact in microbiology. *Journal of Experimental Biology*, 209, 1518-1525.

Hall, T. A. (1999). Bioedit: A user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. *Nucleic Acids Symposium Series*, 41, 95-98.

Hamrick, J. L. & Godt, M. J. W. (1996). Effects of life history traits on genetic diversity in plant species. *Philosophical Transactions of the Royal Society London B*, 351, 1291-1298.

Hedrick, P. W., & Fredrickson. R. (2010). Genetic rescue guidelines with examples from mexican wolves and florida panthers. *Conservation Genetics*, 11, 615-626.

Hefzi, F. (2022). Variasi genetik tiga populasi kapuk randu (*Ceiba pentandra* (L.) Gaertn.) di Sumatera Barat dengan menggunakan penanda RAPD. *Skripsi*. Universitas Andalas.

- Hefzi, F., Mansyurdin, M., & Maideliza, T. (2023). Analysis of genetic variations of *Ceiba pentandra* (L.) Gaertn. on several critical lands in West Sumatra using RAPD molecular markers. *Biosaintifika: Journal of Biology & Biology Education*, 15(1), 26-35.
- Heithaus, E. R., Fleming, T. H., & Opler, P. A. (1975). Foraging patterns and resource utilization in seven species of bats in a seasonal tropical forest. *Ecology*, 56(4), 841-854.
- Hershkovitsz, M. A. & Zimmer, E. A. (1996). Conservation patterns in angiosperm rDNA ITS2 sequences. *Nucleic Acid Research*, 24(15), 2857-2867.
- Hershkovitz, M. A. & Lewis, L. A. (1996). Deep-level diagnostic value of the rDNA-ITS region. *Molecular Biology Evolution*, 13(9), 1276-1295.
- Hidalgo, O., Garnatje, T., Susanna, A., & Mathez, J. (2004). Phylogeny of Valerianaceae based on matK and ITS markers, with reference to matK individual polymorphism. *Annals of Botany*, 93, 283-293.
- Hidayat, T., Priyandoko, D., Wardiny, P. Y., & Islami, D. K. (2016). Molecular phylogenetic screening of *Withania somnifera* relative from Indonesia based on internal transcribed spacer region. *HAYATI Journal of Biosciences*, 1-4.
- Hidayati, N. I. (2020). Analisis swot agribisnis kapuk randu di Desa Soekarmojo Kecamatan Purwosari Kabupaten Pasuruan. *AGROMIX*, 11(1), 87-100.
- Hikam, A. G. (2021). Karakteristik genetik invertebrata laut filum arthropoda, mollusca, dan echinodermata di perairan papua. *Skripsi*. Universitas Islam Malang.
- Horisawa, S., Sakuma, Y., & Doi, S. (2009). Qualitative and quantitative PCR methods using species-specific primer for detection and identification of wood rot fungi. *Journal of Wood Science*, 55(2), 133-138.
- Houle, A. (1998). Floating islands: a mode of long-distance dispersal for small and medium-sized seed plants. *Diversity and Distributions*, 4, 201-216.
- Immanuel, R. R., & Ganapathy, M. (2007). Growth and physiological attributes of *Ceiba pentandra* (L.) Gaertn. seeds and seedlings under salt stress. *Journal of Agricultural and Biological Science*, 2(6), 12-16.
- Immanuel, R. R., & Ganapathy, M. (2020). Restoration of soil microbial biomass using multipurpose trees. *Plant Archives*, 20, 2641-2645.
- Irawanti, S., Race, D., Stewart, H., Parlinah, N., & Suka, A. P. (2017). Understanding the timber value chain in community-based forestry in

Indonesia: Analysis of Sengon in Central Java. *Journal of Sustainable Forestry*, 36(8), 847–862.

Janson, C. H., Terborgh, J., & Emmons, L. H. (1981). Non-flying mammals as pollinating agents in the Amazonian Forest. *Biotropica*, 13(2), 1-6.

Jobes, D. V. & Thien, L. B. (1997). A conserved motif in the 5.8S ribosomal RNA (rRNA) gene is a useful diagnostic marker for plant internal transcribed spacer (ITS) sequences. *Plant Molecular Biology Reporter*, 15, 326-334.

John, M. E. (1992). An efficient method for isolation of RNA and DNA from plants containing polyphenolics. *Nucleic Acids Research*, 20(9), 2381.

Kan, X. Z., Wang, S. S., Ding, X., & Wang, X. Q. (2007). Structural evolution of nrDNA ITS in Pinaceae and its phylogenetic implications. *Molecular Phylogenetics and Evolution*, 44, 765-777.

Kaneko, K., & Furusawa, C. (2005). An evolutionary relationship between genetic variation and phenotypic fluctuation. *Journal of Theoretical Biology*.

Kim S. J., Cho K.S., Yoo K. O., Lim K. B., Hwang Y. J., Chang D. C., & Kim K. S. (2015). Sequence analysis of the internal transcribed spacer (ITS) region of the nuclear ribosomal DNA (nrDNA) *Chrysanthemum* species in Korea. *Horticulture, Environment, and Biotechnology*. 56, 44-53.

Kuruvilla, J., & Anilkumar, M. (2018). Pharmacognostical studies in the leaves of *Ceiba pentandra* (L.) Gaertn. *Journal of Pharmacognosy and Phytochemistry*, 7(6), 46-54.

Kuzmina, M. L., Johnson, K. L., Barron, H. R., & Hebert, P. D. (2012). Identification of the vascular plants of Churchill, Manitoba, using a DNA barcode library. *BMC Ecology*, 12, 25.

Lackie, J. M. (2013). *The dictionary of cell & molecular biology* (5th ed.). Academic Press.

Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A., McWilliam, H., Valentin, F., Wallace, I. M., Wilm, A., Lopez, R., Thompson, J. D., Gibson, T. J., & Higgins, D. G. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics*, 23, 2947-2948.

Li, D.-Z., Gao, L.-M., Li, H.-T., Wang, H., Ge, X.-J., Liu, J.-Q., Chen, Z.-D., Zhou, S.-L., Chen, S.-L., Yang, J.-B., Fu, C.-X., Zeng, C.-X., Yan, H.-F., Zhu, Y.-J., Sun, Y.-S., Chen, S.-Y., Zhao, L., Wang, K., Yang, T., & Duan, G.-W. (2011). Comparative analysis of a large dataset indicates that internal transcribed spacer (ITS) should be incorporated into the core barcode for seed plants. *Proceedings of the National Academy of Sciences*, 108(49), 19641-19646.

- Li, X., Xie, R., Lu, Z., & Zhou, Z. (2010). The origin of cultivated citrus as inferred from internal transcribed spacer and chloroplast DNA sequence and amplified fragment length polymorphism fingerprints. *Journal of the American Society for Horticultural Science*, 135(4), 341-350.
- Lim, T. K. (2012). *Ceiba pentandra*. In: *Edible medicinal and non-medicinal plants Vol. 3* (pp. 540-549). Germany: Fruits, Springer Science and Business Media.
- Liu, J. S. & Schardl, C. L. (1994). A conserved sequence in internal transcribed spacer 1 of plant nuclear rRNA genes. *Plant Molecular Biology*, 26(2), 775-778.
- Liu, J., Moller, M., Gao, L. M., Zhang, D. Q., & Li, D. Z. (2011). DNA barcoding for the discrimination of Eurasian yews (*Taxus* L., Taxaceae) and the discovery of cryptic species. *Molecular Biology Resources*, 11, 89-100.
- Liu, Z., Liu, Y., Liu, S., Ding, X., Yang, Y., & Huang, H. (2009). Analysis of the Sequence ITS1-5.8S- ITS2 Regions of the three species of *Fructus evodiae* in Guizhou province of China and identification of main ingredients of their medicinal chemistry. *Journal of Computer Science and Systematic Biology*, 2, 200-207.
- Maggini, F., Frediani, M., & Gelati, M. T. (2000). Nucleotide sequence of the internal transcribed spacers of ribosomal DNA in *Picea abies* Karst. *DNA Sequence*, 11(1-2), 87-89.
- Maley, J. & Livingstone, D. A. (1983). Late Pleistocene and early Holocene extension of a mountain element in Southern Ghana (West Africa): Preliminary pollen data. *Comptes Rendus de L'Academie des Sciences Serie II*, 296, 1287-1292.
- Maras, M., Aksoz, E., & Menemen, Y. (2006). The structural features and phylogenetic utility of the ITS in *Ferulago* W. Koch (Umbelliferae) genus. *International Journal of Botany*, 2(1), 17-22.
- Mawarni, S. N., Khairunnisa, D., Larasati, I., Rizqo, N., Erfianti, T., Kusumaningrum, H. P., Ferniah, R. S., & Pujiyanto, S. (2021). Application of Doyle and Doyle method for DNA isolation from pamelu yellow orange (*Citrus maxima* Merr), lime orange (*C. limon*) and sunkist orange (*C. sinensis*). *Journal of Physics: Conference Series*, 1943, 1-5.
- Milot, E., Weimerskirch, H., Duchesne, P., & Bernatchez, L. (2007). Surviving with low genetic diversity: the case of albatrosses. *Proceedings of the Royal Society B*, 274, 779-787.

- Mir, B. A., Koul, S., Kumar, A., Kaul, M. K., Soodan, A. S., & Raina, S. N. (2010). Intraspecific variation in the internal transcribed spacer (ITS) regions of rDNA in *Withania somnifera* (Linn.) Dunal. *Indian Journal Biotechnology*, 9, 325-328.
- Mukhamedov, R. S., Martsinkovskaia, A. I., Shu, K., & Abdugarimov, A. A. (1994). Nucleotide sequence of internal transcribed spacers and 5.8S rDNA for the ribosomal operon from alfalfa *Medicago sativa* and cotton *Gossypium hirsutum* L. *Molecular Biology*, 28(1), 204-209.
- Munirah, S., Mat, N., Hasan Nudin, N.F., Mazlan, N., Arumugam, N., Bakar, M.F.A., Wahid, M.A.A., & Ali, A.M. (2014). The Distribution of *Dioscorea hispida* Dennst. germplasm in Terengganu and phylogenetic relationships of *Dioscorea* spp. using internal Transcribed (ITS). *Journal of Agrobiotechnology*, 5, 31-47.
- Murawski, D. A., & Hamrick, J. L. (1992). Mating system and phenology *Ceiba pentandra* (Bombacaceae) in central Panama. *Journal of Heredity*, 83(6), 401-404.
- Murphy B. P., & Tranel P. J. (2018). Identification and validation of *Amaranthus* species-specific SNPs within the ITS region: applications in quantitative species identification. *Crop Science*, 58, 304-311.
- Mursyidin, D. H., & Setiawan, A. (2023). Assessing diversity and phylogeny of Indonesian breadfruit (*Artocarpus* spp.) using internal transcribed spacer (ITS) region and leaf morphology. *Journal of Genetic Engineering and Biotechnology*, 21(15), 1-12.
- Nei, M., & Kumar, S. (2000). *Molecular Evolution and Phylogenetics*. New York: Oxford University Press, p. 333.
- Ningrum, W. D. A., Atmaja, M. B., Daryono, B. S., & Purnomo. (2020). Genetic variability of *Begonia longifolia* Blume from Indonesia based on Nuclear DNA internal transcribed spacer (ITS) sequence data. *BIODIVERSITAS*, 21(12), 5778-5785.
- Nkoum, G. B., Adjoh, G., Leudeu, C. B. T., Kouebou, C., Tchiegang, C., & Kapseu, C. (2017). Local uses of kapok (*Ceiba pentandra* Gaertn.) tree from the northern part of Cameroon. *International Journal of Environment, Agriculture, and Biotechnology*, 2(4), 2214-2219.
- Nyffeler, R., & Baum, D. A. (2000). Phylogenetic relationships of the durians (Bombacaceae-Durioneae or /Malvaceae/Helicteroideae/Durioneae) based on chloroplast and nuclear ribosomal DNA sequences. *Plant Systematic Evolution*, 224, 55-82.

- Orwa, C. M. (2009). *Agroforestry database: a tree reference and selection guide version 4.0*. Kenya: World Agroforestry Centre.
- Pang, X., Luo, H., & Sun, C. (2012). Assessing the potential of candidate DNA barcodes for identifying non-flowering seed plants. *Plant Biology*, *14*, 839-844.
- PROSEA (Plant resources of South-East Asia 17). (2003). In: M. Brink & R. P. Escobin (Ed.), *Fibre plants* (pp. 99-101). Leiden: Backhuys Publishers.
- Putri, E. M. (2012). Biodiesel production from kapok seed oil (*Ceiba pentandra*) through the transesterification process by using CaO as catalyst. *Global Journal of Researches in Engineering Chemical Engineering*, *12*(2), 6-11.
- Qin, Y., Li, M., Cao, Y., Gao, Y., & Zhang, W. (2017). Molecular thresholds of ITS2 and their implications for molecular evolution and species identification in seed plants. *Scientific Reports*, *7*, 17316.
- Queiroz, C. S., Batista, F. R. C., & Oliveira, L. O. (2011). Evolution of the 5.8S nrDNA gene and internal transcribed spacers in *Carapichea ipecacuanha* (Rubiaceae) within a phylogeographic context. *Molecular Phylogenetics and Evolution*, *59*, 293-302.
- Rahmadara, G., Hanifah, N. F., Rismayanti, Purwoko, D., Rochandi, A., & Tajuddin, T. (2022). Comparison of DNA isolation methods that derived from leaves of a potential anti-cancer rodent tuber (*Typhonium flagelliforme*) plant. *International Journal of Agriculture System*, *10*(2), 93-103.
- Rojas-Sandoval, J., Lobo, J. A., & Quesada, M. (2008). Phenological patterns and reproductive success of *Ceiba pentandra* (Bombacaceae) in tropical dry and wet forests of Costa Rica. *Revista Chilena de Historia Natural*, *81*, 443-454.
- Rozas, J., Ferrer-Mata, A., Sanchez-DelBarrio, J. C., Guirao-Rico, S., Librado, P., Ramos-Onsins, S. E., & Sanchez-Gracia, A. (2017). DnaSP 6: DNA sequence polymorphism analysis of large data sets. *Molecular Biology and Evolution*, *34*(12), 3299-3302.
- Sahid, M., & Zeven, A. C. (2003). *Ceiba pentandra* (L.) Gaertn. In: Brink, M., Escobin, R. P. (Ed.), *Plant resources of South-East Asia No. 17* (pp. 399-400). Netherlands: Backhuys Publishers.
- Saini, A., Reddy, S. K., & Jawali, N. (2018). Intra-individual and intra-species heterogeneity in nuclear rDNA ITS region of *Vigna* species from subgenus *Ceratopics*. *Genetic Research Cambridge*, *90*, 299-316.
- Salimi, H., Bahar, M., Mirlohi, A., & Talebi, M. (2016). Assessment of the genetic diversity among potato cultivars from different geographical areas using the

- genomic and EST microsatellites. *Iran Journal Biotechnology*, 14(4), 270-277.
- Samson, R. A., Houbraken, J., Thrane, U., Frisvad, J. C., & Andersen, B. (2010). *Food and indoor fungi*. Centraalbureau voor Schimmelcultures.
- Sass, C., Little, D. P., Stevenson, D. W., & Specht, C. D. (2007). DNA barcoding in the cycadales: testing the potential of proposed barcoding markers for species identification of cycads. *PLOS ONE*, 2.
- Sayed, L. M. (2018). Utilization of internal transcribed spacer (ITS) as a molecular marker for phylogenetic relationship of Solanaceae family. *Egyptian Journal of Genetics and Cytology*, 46, 167-178.
- Senavirathna, H. M. T. N., Ranaweera, L. T., Mudannayake, M. M. A. W. P., Nawanjana, P. W. I., Wijesundara, W. M. D. A., Jayarathne, H. S. M., Ratnasuriya, M. A. P., Weebadde, C. K., & Sooriyapathirana, S. D. S. S. (2020). Assessment of the taxonomic status of the members of genus *Artocarpus* (Moraceae) in Sri Lanka. *Genetic Resources and Crop Evolution*, 67, 1163-1179.
- Shaw, A. J., McDaniel, S. F., Werner, O., & Ros, R. M. (2002). Phylogeography and phylodemography. *The Bryologist*, 105(3), 373-383.
- Sherwin, W. B., & Moritz, C. (2000). Managing and monitoring genetic erosion. In: A. G. Young & G. M. Clarke (Ed.), *Genetics, demography and viability of fragmented populations* (pp. 9-34). Cambridge, UK: Cambridge University Press.
- Shoaie, S. (2022). A short notes on point mutations and its types and function. *European Journal of Experimental Biology*, 12(5), 137.
- Singaraleven, N., & Marimuthu, G. (2004). Nectar feeding and pollen carrying from *Ceiba pentandra* by pteropodid bats. *Journal of Mammalogy*, 85(1), 1-7.
- Slotta, T. A. B. (2000). *Phylogenetic analysis of Iliamna (Malvaceae) using the internal transcribed spacer region*. Thesis. Virginia Polytechnic Institute and State University.
- Standley, P. C. (1931). *The forests and flora of British Honduras*. Field Museum of Natural History.
- Suh, Y., Thien, L. B., & Zimmer, E. A. (1992). Nucleotide sequences of the internal transcribed spacers and 5.8S rRNA gene in *Canella winterana* (Magnoliales; Cannellaceae). *Nucleic Acids Research*, 20, 6101-6102.

- Sultana, S., Lim, Y. P., Bang, J. W., & Choi, H. W. (2011). Internal transcribed spacer (ITS) and genetic variations in *Lilium* native to Korea. *Horticulture Environment and Biotechnology*, 52, 502-510.
- Takhtajan, A. (1986). *Floristic regions of the world*. Berkeley: University of California Press.
- Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38, 3022-3027.
- Tate, J. A., Aguilar, J. F., Wagstaff, S. J., Duke, J. C. L., Slotta, T. A. B., & Simpson, B. B. (2005). Phylogenetic relationships within the tribe Malveae (Malvaceae, subfamily Malvoideae) as inferred from ITS sequence data. *American Journal of Botany*, 92(4), 584-602.
- Terra V., Garcia F. C. P., Queiroz L. P. de, Bank M., Miller J. T. (2017). Phylogenetic Relationships in *Senegalia* (Leguminosae-Mimosoideae) Emphasizing the South American Lineages. *Systematic Botany*, 42(3), 458-464.
- Terryana, R. T., Nugroho, K., Rijzaani, H., & Lestari, P. (2018). Karakterisasi keragaman genetik 27 genotipe cabai berdasarkan marka SSR (Simple Sequence Repeat). *Berita Biologi*, 17(2), 183-194.
- Thorne, R. F. (1973). Floristic relations between tropical Africa and tropical America. In: Meggers, B., Ayesu, E., Duckworth, W (Ed.), *Tropical Forest Ecosystems in Africa and South America: A Comparative Review*, pp. 27-47. Washington DC: Smithsonian Institution Press.
- Toha, A. H. A. (2001). Keragaman genetik bulu babi (Echinoidea). *Biota*, 12(2), 131-135.
- Tokouka, T. (2008). Molecular phylogenetic analysis of Violaceae (Malpighiales) based on plastid and nuclear DNA sequences. *Journal of Plant Research*, 121, 253-260.
- Toledo, V. M. (1977). Pollination of some rain forest plants by non-hovering birds in Veracruz, Mexico. *Biotropica*, 9(4), 262-267.
- Topik, H., Yukawa, T., & Ito, M. (2005). Molecular phylogenetics of subtribe Aeridinae (Orchidaceae): Insights from plastid matK and nuclear ribosomal ITS sequences. *Journal of Plant Research*, 18, 271-284.
- Tripathi, A. M., Tyagi, A., Kumar, A., Singh, A., Singh, S., Chaudhary, L. B., & Roy, S. (2013). The internal transcribed spacer (ITS) region and trnH-psbA are suitable candidate loci for DNA barcoding of tropical tree species of India. *PLOS ONE*, 8(2), e57934.

- Tsai, C. C., Chou, C. H., Wang, H. V., Ko, Y. Z., Chiang, T. Y., & Chiang, Y. C. (2015). Biogeography of the *Phalaneopsis amabilis* species complex inferred from nuclear and plastid DNAs. *BMC Plant Biology*, *15*(202), 1-16.
- Turaki, A., A., Ahmad, B., Magaji, U. F., Abdulrazak, U. K., Yusuf, B. A., & Hamza, A. B. (2017). Optimised cetyltrimethylammonium bromide (CTAB) DNA extraction method of plant leaf with high polysaccharide and polyphenolic compounds for downstream reliable molecular analyses. *African Journal of Biotechnology*, *16*(24), 1354-1365.
- Ubaidillah, R. & Sutrisno, H. (2009). Pengantar biosistemik: teori dan praktikum. Jakarta: LIPI Press.
- Vieira, F.C., de Laia, M.L., Marques, A., Goncalves, J.F., & Veloso, R.V.S. (2017). Use of the internal transcribed spacer (ITS) regions to characterize Macaw progenies. *Biotechnology Journal International*, *18*(2), 1-8.
- Wang, D. Y., Wang, Q., Wang, Y. L., Xiang, X. G., Huang, L. Q., & Jin, X. H. (2017). Evaluation of DNA barcodes in *Codonopsis* (Campanulaceae) and in some large angiosperm plant genera. *PLOS ONE*, *12*(2), e0170286.
- Warseno, T., Efendi, M., Chasani, A. R., Daryono, B. S. (2022). Genetic variability and phylogenetic relationships of *Begonia multangula* based on atpB-rbcL non-coding spacer of cpDNA sequences. *BIODIVERSITAS*, *23*(10), 5491-5501.
- White, T., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: M.A. Innes, D. H. Gelfand, J. S. Sninsky, & T. J. White (Ed.), *PCR protocols: a guide to methods and applications* (pp. 315-322). London: Academic Press.
- Wibawa, H. (2022). *Introduction to DNA sequencing technology*. Yogyakarta: Universitas Gadjah Mada.
- Wilson, J., Lowe, A. J., & Cavers, S. (1997). Assesment of levels and dynamic of intra specific genetic diversity of tropical trees: final scientific report. *European Commission*, 1-76.
- Wulandari, E. F., Mawikere, N. L., & Abbas, B. (2021). Keragaman morfologi dan genetik beberapa aksesi tanaman sagu (*Metroxylon sagu* Rottb.) berdasarkan penanda molekuler gen mat-K. *CASSOWARY*, *4*(1), 68-86.
- Xu, P., Jiang, Y., Xu, J., Li, J., & Sun, X. (2016). Genomics in the common carp. In: *Genomics in aquaculture* (pp. 247-274). Academic Press.
- Zhang, J. Q., Meng, S. Y., Wen, J., & Rao, G. Y. (2015). DNA barcoding of *Rhodiola* (Crassulaceae): a case study on a group of recently diversified

medical plants from the Qinghai-Tibetan Plateau. *PLOS ONE*, 10(3), e0119921.

Zhang, J., Chiodini, R., Badr, A., & Zhang, G. (2001). The impact of next-generation sequencing on genomics. *Journal of Genetics and Genomics*, 38(3), 95-109.

Zhou B., Tu T., Kong F., Wen J., and Xu X. (2018). Revised phylogeny and historical biogeography of the cosmopolitan aquatic plant genus *Typha* (Typhaceae). *Scientific reports*, 8, 8813.

Zhou, Z., Jiang, Y., Wang, Z., Gou, Z., Lyu, J., Li, W., Yu, Y., Shu, L., Zhao, Y., Ma, Y., Fang, C., Shen, Y., Liu, T., Li, C., Li, Q., Wu, M., Wang, M., Wu, Y., Dong, Y., Wan, W., Wang, X., Ding, Z., Gao, Y., Xiang, H., Zhu, B., Lee, S., Wang, W., & Tian, Z. (2015). Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. *Nature Biotechnology*, 33(4), 408-441.

