

## DAFTAR PUSTAKA

- Agbor, G.A., Vinson, J.A., & Donnelly, P.E. (2014). Folin-Ciocalteu reagent for polyphenolic assay. *Int. J. Food Sci. Nutr. Diet*, 3, 8, 147-156.
- Agoes, G. (2007). *Teknologi bahan alam (sf-2)* (ed. revisi). Bandung: Penerbit ITB.
- Ampofo, S.A., & Waterman, P.G. (1986). Xanthenes from three garcinia species. *Phytochemistry*, 25, 10, 2351-2355.
- Badan Pengawasan Obat dan Makanan RI. (2008). *Acuan sediaan herbal* (vol. 4, ed. 1). Jakarta: Badan Pengawasan Obat dan Makanan.
- Badan Pengawasan Obat dan Makanan RI. (2013). *Pedoman teknologi formulasi sediaan berbasis ekstrak*. Jakarta: Badan Pengawasan Obat dan Makanan.
- Benzie, I.F.F., & Strain, J.J. (1999). Ferric reducing antioxidant power assay. *Method in Enzymology*, 299, 15-27.
- Cadenas, E., & Packer, L. (2002). *Handbook of antioxidants*. Switzerland: Marcel Dekker Inc.
- Dachriyanus. (2004). *Analisis struktur senyawa organik secara spektrofotometri* (ed. 1). Padang: Andalas University Press.
- Departemen Kesehatan Republik Indonesia. (1995). *Farmakope Indonesia* (ed. 4). Jakarta: Departemen Kesehatan.
- Departemen Kesehatan Republik Indonesia. (2000). *Parameter standar umum ekstrak tumbuhan obat*. Jakarta: Departemen Kesehatan.
- Departemen Kesehatan Republik Indonesia. (2008). *Farmakope herbal Indonesia* (ed. 1). Jakarta: Departemen Kesehatan.
- Dharmaratne, H.R.W., & Wanigasekera, W.M.A.P. (1996). Xanthenes from root bark of *Calophyllum thwaitesii*, *Phytochemistry*, 42, 1, 249-250.
- Fessenden, R.J., & Fessenden, J.S. (1986). *Kimia organik* (ed. 3, jilid 1). Penerjemah: A.H. Pudjaatmaka. Jakarta: Erlangga.
- Gandjar, I.G., & Rohman, I. (2007). *Kimia farmasi analisis*. Yogyakarta: Pustaka Pelajar.

- Harborne, J.B. (1987). *Metode fitokimia, penuntun cara modern menganalisis tumbuhan*. Penerjemah: K. Patmawinata, I. Soediro. Bandung: Penerbit ITB.
- Harmita. (2004). Petunjuk pelaksanaan validasi metode dan cara perhitungannya. *Ilmu Kefarmasian*, 1, 117-135.
- Harris, D.C. (2007). *Quantitative chemical analysis* (7<sup>th</sup> ed.). New York: W. H. Freeman and Company.
- Huang, D., Ou, B., & Prior, R.L. (2005). The chemistry behind antioxidant capacity assays. *Agricultural and Food Chemistry*, 53, 1841-1856.
- Jantan, I., Pizar, M., Idris, Taher, M., & Ali, R.M., (2002). In vitro inhibitory effect of rubraxanthone isolated from *Garcinia parvifolia* on platelet-activating factor receptor binding. *Planta Med.*, 68, 1133-1134.
- Jeffery, G. H., Basset, J., Mendham, J., & Denney, R. C. (1989). *Vogel's textbook of quantitative chemical analysis* (5<sup>th</sup> ed.) .New York: John Wiley Sons Inc.
- Jena, B.S., Jayaprakasha, G.K., & Sakariah, K.K. (2002). Organic acids from leaves, fruits, and rinds of *Garcinia cowa*. *Agricultural and Food Chemistry*, 50, 12, 3431-3434.
- Lee, K.W., Kim, Y.J., Lee, H.J., & Lee, C.Y. (2003). Cocoa has more phenolic phytochemicals and a higher antioxidant capacity than teas and red wine. *Agricultural and Food Chemistry*, 51, 7292-7295.
- Lim, K.L. (2012). *Edible medicinal and non-medicinal plants* (vol. 2). Berlin: Springer.
- Lisdawati, V., Wiryowidagdo, S., & Kardono, B.S. (2007). Isolasi dan elusidasi struktur senyawa lignan dan asam lemak dari ekstrak daging buah *Phaleria macrocarp*. *Penel Kesehatan*, 35, 3, 115-124.
- Mahabusarakam, W., Chairerk, P., & Taylor, W.C. (2004). Xanthones from *Garcinia cowa* Roxb. Latex. *Phytochemistry*, 66, 1148-1153.
- Marinova, G., & Batchvarov, V. (2011). Evaluation of the methods for determination of the free radical scavenging activity by DPPH. *Bulgarian Agricultural Science*, 17, 1, 11-24.
- Murakami, A., Jiwajiinda, S., Koshimizu, K., & Ohigashi, H. (1995). Screening for in vitro anti-tumor promoting activities of edible plants from Thailand. *Cancer Lett*, 95, 137-146.

- Mustafa, R., Hamid, A., Mohamed, S., & Bakar, F. (2010). Total phenolic compounds, flavonoids, and radical scavenging activity of 21 selected tropical plants. *J. of Food Science*, 75, C28-C35.
- Na Pattalung, P., Thongtheeraparp, W., Wiriyaichitra, P., & Taylor, W.C. (1994). Xanthenes of *Garcinia cowa*. *Planta Med.*, 60, 365-368.
- Panthong, K., Pongcharoen, W., Phongpaichit, S., & Taylor, W.C. (2006). Tetraoxygenated xanthenes from the fruits of *Garcinia cowa*. *Phytochemistry*, 67, 999-1004.
- Pietta, P.G. (2000). Reviews: Flavonoids as Antioxidants. *Journal of Natural Products*, 63, 7, 1035-1042.
- Pratt, D.E. (1992). Natural antioxidants from plant material. *Food and Agriculture Organisation*, 507, 54-71.
- Prior, R.L., Wu, X., & Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.*, 53, 4290-4302
- Rebeta, M.S. & Nur, F.R. 2013. Total phenolic content and ferric reducing antioxidant of the leaves and fruits of *Garcinia atroviridis* and *Cynometra cauliflor*. *International Food Research*, 20, 1691-1696.
- Ritthiwigrom, T., Laphookieo, S., & Pyne, G. (2013). Chemical constituents and biological activities of *Garcinia cowa* Roxb. *Maejo Int. J. of Sci. and Tech.*, 7, 2, 212-231.
- Rohman, A. (2014). *Validasi dan penjaminan mutu; metode analisa kimia*. Yogyakarta: Gadjah Mada University Press.
- Silverstein, R.M., Bassler, G.C., & Morrill, T.C. (1981). *Spectrometric Identification of Organic Compound* (4th ed.). Singapore: John Wiley and Sons.
- Singleton, V.L., & Rossi, J.A. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Enology and Viticulture*, 16,3, 144-158.
- Song, Q.S., Na, Z., Hu, H.B. (2013). A new prenylated xanthone from latex of *Garcinia cowa* Roxb. *Academy of Chemistry of Globe Publications*, 7, 3, 220-224.
- Suryohudoyo, P. (2000). *Oksidan, antioksidan, dan radikal Bebas*. Jakarta: Infomedika.

- Terry, L.A., Vicente, A., & Cools, K. (2011). Methodologies for extraction, isolation, characterization and quantification of bioactive compounds. *CAB Int. J.*, 375-376.
- Tjitrosoemo, G. (1993) Taksonomi tumbuhan (spermatophyta), Yogyakarta: Gadjah Mada University Press.
- Trisuwan, K., & Ritthiwigrom T. (2012). Benzophenone and xanthone derivatives from the inflorescent of *Garcinia cowa*. *Arch Pharm Res*, 35, 1733-1738.
- Tursiman, Puji A., & Risa N. (2012). Total fenol fraksi etil asetat dari buah asam kandis (*Garcinia dioica* Blume). *JKK*, 1, 1, 45-48.
- Umayah, E.U. & Moch A.H., (2007), Uji Aktivitas antioksidan ekstrak buah naga (*Hylocereus undatus* Haw. *Ilmu Dasar*, 8, 1, 83-90.
- Vichitphan, S., Vichitphan, K., & Sirikhansaeng P. (2007). Flavonoid content and antioxidant activity of krachai-dum (*Kaempferia parviflora*) wine. *Sci. Tech. J.*, 7, 2, 97-105.
- Wahyuni, F.S., Byrne, L.T., Dachriyanus, Dianita, R., Jubahar, J., & Lajis, N.H. (2004). A new ring-reduced tetraprenyltoluquinone and a prenylated xanthone from *Garcinia cowa*. *Aust. Chemistry*, 57, 3, 223-226.
- Wahyuni, F.S., Shaari, K., Stanslas, J., Lajis, N.H., & Dachriyanus. (2015). Cytotoxic xanthenes from the stem bark of *Garcinia cowa* Roxb. *J. of Chemical and Pharmaceutical Research*, 7, 1, 227-230.
- Waterhouse, A. L. (2002). *Determination of total phenolic*. California: University of California Press.
- Werdhasari, A. (2014). Peran antioksidan bagi kesehatan. *Biotek Medisiana Indonesia*, 3, 2, 59-68.
- Winarsi, H. (2007). *Antioksidan alami dan radikal bebas*. Yogyakarta: Kanisius.