

DAFTAR PUSTAKA

- Abhinandan KS, Vedavathi R, 2013, "Usefulness of Sequential Organ Failure Assessment (Sofa) and Acute Physiology And Chronic Health Evaluation II) Score in Analysing Patients With Multiple Organ Dysfunction Syndrome in Sepsis". *Journal of Evolution of Medical and Dental Sciences*; Vol2, Issue 49: 9591-9605.
- Angele MK, Pratscke S, Hubbard WJ, Chaudry IH, 2014, Gender Differences in Sepsis, Cardiovascular and Immunological Aspects, *Virulence* 5:1, 12-19.
- Angus DC, Poll T, 2013, Severe Sepsis and Septic Shock, *N Engl J Med* :369;9.3
- Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR, 2001. Epidemiology of Severe Sepsis in the United States: Analysis of Incidence, Outcome, and Associated Costs of Care. *Crit Care Med*, 29: 1303-1310
- Bale C, Kakrani AL, Dabadghao VS, Sharma ZD, 2013, Sequential Organ Failure Assessment Score As Prognostic Marker in Critically Ill Patients in A Tertiary Care Intensive Care Unit. *Int J Med Public Health* ;3:155–8.
- Bayram H, Tunger O, Civi M, Yuceyar MH, Ulman C, Horason GD, et al, 2015, Diagnostic and prognostic value of Procalcitonin and sTREM-1 level in Sepsis, *Turk J Med Sci*, 45: 578-86.
- Bochud P, Calandra T, 2003, Pathogenesis of Sepsis: New Concept and Implication for Future Treatment. *BMJ*; 326: 262-6.
- Buchori, Prihatini, 2006, Diagnosis Sepsis Menggunakan Procalcitonin (*Sepsis Diagnosis by Procalcitonin*), *Indonesian Journal of Clinical Pathology and Medical Laboratory*, Vol. 12, No. 3,: 131-137.
- Castelli G, Pognani C, Meisner M, et al, 2004, Procalcitonin and C-Reactive Protein During Systemic Inflammatory Response Syndrome, Sepsis and Organ Dysfunction. *Critical Care*, 8: 234-242.
- Centers for Disease Control and Prevention(CDC). (2010). QuickStats: Hospitalization Rates for Patients Aged ≥ 65 Years with Septicemia or Sepsis, by Age Group. National Hospital Discharge Survey, United States, 2000–2007. *MMWR* 59(34):110.
- Cohen J, 2002, The immunopathogenesis of Sepsis, *Nature*, 420, 885-891.
- Crain MC, Muller B, 2005, Procalcitonin in Bacterial Infection- Hype, Hope, More or Les?, *Swiss Med Wkly*, 135: 451-460. Daley BJ, 2017, Peritonitis and Abdominal Sepsis, diunduh dari <http://www.emedicine.medscape.com>.
- Dombrovskiy VY, Martin AA, Sunderram J, Paz HL (2007). Rapid Increase in Hospitalization and Mortality Rates for Severe Sepsis in the United States: a Trend Analysis from 1993 to 2003. *Crit Care Med*, 35: 1244-1250.

- El Maraghi S, Yehia A, Mahmoud O, El Hamid A, 2014, Procalcitonin Versus C-Reactive Protein at Different SOFA Scores in I.C.U. Sepsis: Diagnostic Value and Therapeutic Implications, *Med. J. Cairo Univ*, Vol. 82, No. 1, 29-36
- Endo S, Fujishima S, Aikawa N, Sekine I, Kogawa K, Yamamoto Y, *et al*, 2008, Usefullness of procalcitonin Serum Level for the Discrimination of Severe Sepsis from Sepsis: a Multicentre Prospective Study, *Journal of Infection and Chemotherapy*, vol 14(3), 224-49.
- Jain S, Sinha S, Sharma S, Samantaray JC, Aggrawal P, Vikram NK, *et al*, 2014, Procalcitonin as a prognostic Marker for Sepsis: a Prospective Observational Study, *BMC Research Notes*, 7:458.
- National Clinical Effectiveness Committee (NCEC), 2014, Sepsis Management National Clinical Guideline No. 6, An Roine Slainte Departement of Health, Ireland.
- Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, *et al.*, 2013, Surviving Sepsis Campaign: International Guidelines for Management of Severe Sepsis and Septic Shock: 2012, *Crit Care Med*; 41(2) :580–637.
- Ferreira FL, Bota DP, Bross A, Mélot C, Vincent JL, 2001, Serial Evaluation of The SOFA Score to Predict Outcome in Critically Ill Patients, *JAMA*;286:1754–8.
- Gomez CR, Nomellini V, Faunce DE, Kovacs EJ, 2008,. Innate Immunity and Aging. *Exp Gerontol*, 43: 718-728.
- Guntur A. 2009. Sepsis. Dalam: Sudoyo A, Setiyohadi B, Alwi I, Simadibrata M, S s, (eds), Buku Ajar Ilmu Penyakit Dalam. Edisi V. Interna Publishing: Jakarta.
- Harbarth S, Holeckova K, Froidevaux C, Pittet D, Ricou B, Grau GE, *et al*, 2001, Diagnostic Value of Procalcitonin, Interleukin-6, and Interleukin-8 in Critically Ill Patients Admitted with Suspected Sepsis. *Am J Respir Crit Care Med*;164:396-402.
- Hidayati, Arifin H, Raveinal, 2016, Kajian Penggunaan Antibiotik pada Pasien Sepsis dengan Gangguan Ginjal, *Jurnal Sains Farmasi & Klinis*, 2(2), 129-137.
- Kuncoro H, Suta IB, 2015, Peranan Prokalsitonin Dalam Bidang Pulmonologi, *J Respir Indo* Vol. 35 No. 3.
- Kurosawa DJS, Osuchowski MF, Valentine C, Kurosawa S, Remick DG, 2011, The Pathogenesis of Sepsis, *Annu Rev Pathol* . 2011 ; 6: 19–48.
- Leclerc F, Leteurtre S, Noizet O, Dorkenoo A, Sadik A,Cremer R, Fourier C, 2002, Procalcitonin as a Prognostic Marker in Children with Meningococcal Septic Shock. *Arch Dis Child* , 87:450
- Lee H, 2013, Procalcitonin as a Biomarker of Infectious Disease, *Korean J Intern Med* ;28:285-91.

- Levy M, Fink FM, Marshall JC, Abraham E, Angus D, Cook D et al, 2003, 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference, *Intensive Care Med* ;29:530–538.
- Martin GS, 2012, Sepsis, Severe Sepsis and Septic Shock: Changes in Incidence, Pathogens and Outcomes, *Expert Rev Anti Infect Ther*; 10(6): 701–06.
- Martin GS, Mannino DM, Eaton S, Moss M, 2003, The epidemiology of sepsis in the United States from 1979 through 2000, *N Engl J Med* ; 348:1546–54
- Maruna P, Nedelkinova K, Gurlich R, 2000, Physiology and Genetics of Procalcitonin, *Physiol.Res.*49 (suppl.1):S57-S61.
- Meissner M, Tschaikowsky K, Palmaers T, Schmidt J, 1999, Comparison of Procalcitonin (PCT) and C-Reactive Protein (CRP) Plasma Concentrations at Different SOFA Scores During the Course of Sepsis and MODS, *Crit Care* , 3:45–50
- Meissner M, 2010, Procalcitonin-Biochemistry and Clinical Diagnosis, 1st ed. UNIMED, International Medical Publisher, London.
- Meissner M, 2014, Update on Procalcitonin Measurement, *Ann Lab Med*; 34: 263-273.
- Muller B, Becker KL, Schachinger H, Rickenbacher PR, Zimmerli W, Ritz R, 2000, Calcitonin Precursors Are Reliable Markers of Sepsis in A Medical Intensive Care Unit. *Crit Care Med*; 977-983
- Munford RS, 2008, Severe sepsis and septic shock. In: A. Fauci, *et al.*, (eds), *Harrison's Principles of Internal Medicine*, 17th ed. New York: McGraw-Hill Medical, Ch. 265.
- Nasa P, Juneja D, Singh O, 2012, Severe Sepsis and Septic Shock in the Elderly: An Overview, *World J Crit Care Med*, 1(1): 23-30.
- Nair R, Bhandary NM, dan D'Souza AD, 2016. Initial Sequential Organ Failure Assessment score versus Simplified Acute Physiology score to analyze Multiple Organ Dysfunction in Infectious Diseases in Intensive Care Unit, *Indian J Crit Care Med.*, 20, p: 210-5.
- Nargis W, Ahamed BU, 2014, Procalcitonin Versus C-Reactive Protein: Usefulness As Biomarker of Sepsis in ICU Patient, *International Journal of Critical Illness and Injury Science*: 4(3); 195-9.
- Picard KM, O'Donoghoe SC, Young-Kershaw DA, Russell KJ, 2006, Developmental and Implementation of A Multidisciplinary Sepsis Protokol, *Criticalcarenurse* Vol 26, No. 3.
- Perrakis A, Müller V, Hohenberger W, Croner RS, 2012, The Role of Procalcitonin in the Early Diagnosis of Septic and Non-Septic Complications in the Immediate Postoperative Setting After Orthotopic Liver Transplantation, Intech open Science, chapter 9, pp: 187-204.

- Jin M, Khan AI, 2010, Procalcitonin: Uses in the Clinical Laboratory for the Diagnosis of Sepsis, *LabMedicine*: 41(3), pp: 173-7.
- Jones AE, Trzeciak S, Kline JA, 2009, The Sequential Organ Failure Assessment Score for Predicting Outcome in Patients With Severe Sepsis And Evidence Of Hypoperfusion at The Time of Emergency Department Presentation, *Crit Care Med* ; 37(5): 1649–1654
- Saha R, Das S, Chatterjee R, Kaur I, 2010, The Pathophysiology of Septic Shock, International Journal of Pharma and Bio Science, vol 1(2).
- Silva E, Passos R, Ferri MB, Francisco P, Fiquiredo P, 2008, Sepsis: From Bench to Bedside, *Clinics*;63(1): 109-20.
- Singer M, Deutschman C, Seymour CW, Hari MS, Annane D, Bauer M, 2016, The Third International Consensus for Defenition of Sepsis And Septic Shock, *JAMA*;315(8):801-810.
- Seeley EJ, Matthay MA, Wolters PJ, 2012, Inflection Points in Sepsis Biology: From Local Defense to Systemic Organ Injury, *Am J Physiol Lung Cell Mol Physiol* 303: L355–L363.
- Shiferaw B, Bekele E, Kumar K, Boutin A, Frieri M, 2016, The Role of Procalcitonin as A Biomarker in Sepsis, *J.Infect Dis Epidemiol*, vol 2(1);2:006.
- Snowden C, Kirkmen E, 2002, The Pathophysiology of Sepsis, *British Journal of Anasthesia*, CEPD Review: vol 2, No 1.
- Szederjesi J, Almasy E, Lazar A, Huțanu A, Badea I, Georgescu A, 2015, An Evaluation of Serum Procalcitonin and C-Reactive Protein Levels as Diagnostic and Prognostic Biomarkers of Severe Sepsis, *The Journal of Critical Care Medicine*;1(4):147-153
- Vincent JI, Moreno R, Takala J, Willats S, Mendonca AD, Bruining H, 1996, The SOFA (Sepsis Related Organ Failure Assesment) Score to Describe Organ dysfunction/Failure, *Intensive Care Med*;22: 707-710.
- Vincent JL, Rello J, Marshall J, et al, 2009, International Study of the Prevalence and Outcomes of Infection in Intensive Care Units. *JAMA* ; 302 : 2323-9.
- Vosylius S, Sipylaitė J, Ivaskevicius J, 2004, Sequential Organ Failure Assessment Score as The Determinant of Outcome for Patients with Severe Sepsis. *Croat Med J* ;45:715–20.
- Weksler ME, Goodhardt M, Szabo P, 2002, The Effect of Age on B cell Development and Humoral Immunity, *Springer Semin Immunopathol*, 24: 35-52.
- Yang Y, Xie J, Guo F, Longhini F, Gao Z, Huang Y, et al, 2016, Combination of C-Reactive Protein, Procalcitonin and Sepsis Related Organ Failure Score for The Diagnosis of Sepsis in Criticall Patient, *Annals of Intensive Care*, 6:51.