

## REFERENCES

- Adam, F.P., Moenandir, J. and Santoso, M., 2008. Pengaruh pencampuran herbisida dan persiapan lahan terhadap pertumbuhan dan hasil panen padi (the influence of herbicide mix and land preparation on the growth and rice crops yield). *J. Agritek*, 16, pp.1601-1615.
- Akbar, N., Jabran, K. and Ali, M.A., 2011. Weed management improves yield and quality of direct seeded rice. *Australian Journal of Crop Science*, 5(6), p.688.
- Antralina, M., Istina, I.N. and Simarmata, T., 2015. Effect of difference weed control methods to yield of lowland rice in the SOBARI. *Procedia Food Science*, 3, pp.323-329.
- Anwar, M.P., Juraimi, A.S., Man, A., Puteh, A., Selamat, A. and Begum, M., 2010. Weed Suppressive Ability of Rice ('Oryza sativa'L.) Germplasm under Aerobic Soil Conditions. *Australian Journal of Crop Science*, 4(9), p.706.
- Anwar, M.P., Juraimi, A.S., Mohamed, M.T.M., Uddin, M.K., Samedani, B., Puteh, A. and Man, A., 2013. Integration of agronomic practices with herbicides for sustainable weed management in aerobic rice. *The Scientific World Journal*, 2013.
- Awan, T.H., Cruz, P.C.S. and Chauhan, B.S., 2015. Efficacy and economics of different herbicides, their weed species selectivity, and the productivity of mechanized dry-seeded rice. *Crop Protection*, 78, pp.239-246.
- Beckie, H.J., 2006. Herbicide-resistant weeds: Management tactics and practices. *Weed Technology*, 20(3), pp.793-814.
- Begum, M., Juraimi, A.S., Amartalingam, R., Bin Man, A. and Bin Syed Rastans, S.O., 2006. The effects of sowing depth and flooding on the emergence, survival, and growth of *Fimbristylismiliacea* (L.) Vahl. *Weed Biology and Management*, 6(3), pp.157-164.
- Beltran, J.C., Pannell, D.J., Doole, G.J. and White, B., 2012. A bioeconomic model for analysis of integrated weed management strategies for annual barnyardgrass (*Echinochloa crus-galli* complex) in Philippine rice farming systems. *Agricultural Systems*, 112, pp.1-10.

Buhler, D.D., 2002. 50th Anniversary—Invited Article: Challenges and opportunities for integrated weed management. *Weed Science*, 50(3), pp.273-280.

Caton, BP. M Mortimer.J.EHill.E Johnson. 2010. A Practical Field Guide to Weeds of Rice in Asia. Second edition. International Rice Research Institute. Los Banos.

Chauhan, B.S. and Johnson, D.E., 2011. Row spacing and weed control timing affect yield of aerobic rice. *Field Crops Research*, 121(2), pp.226-231.

Chauhan, B.S., 2012. Weed ecology and weed management strategies for dry-seeded rice in Asia. *Weed Technology*, 26(1), pp.1-13.

Chauhan, B.S., 2013. Strategies to manage weedy rice in Asia. *Crop Protection*, 48, pp.51-56.

Chin, D.V., 2001. Biology and management of barnyardgrass, red sprangletop and weedy rice. *Weed Biology and Management*, 1(1), pp.37-41.

Dass, A., Shekhawat, K., Choudhary, A.K., Sepat, S., Rathore, S.S., Mahajan, G. and Chauhan, B.S., 2017. Weed management in rice using crop competition-a review. *Crop Protection*, 95, pp.45-52.

Diggle, A.J., Neve, P., Smith, F.P., 2003. Herbicides used in combination can reduce the probability of herbicide resistance. *Weed Res.* 43. pp 371-382.

DuPont Agricultural Product Indonesia.2015.Herbisida Ally Plus 77 WP.Beltway Office Park. Jakarta. <http://faedahjaya.com/herbisida/ally-plus77WP>. 2016. Ally plus 77 WP. Dupont. Jakarta

Ellis, J.M., Griffin, J.L., Linscombe, S.D. and Webster, E.P., 2003.Rice (*Oryza sativa*) and Corn (*Zea mays*) Response to Simulated Drift of Glyphosate and Glufosinate1. *Weed Technology*, 17(3), pp.452-460.

European Food Safety Authority. 2011. Review of the existing maximum residue levels (MRLs) for 2,4-D. Article 12 of Regulation (EC) No 396/2005 . EFSA Journal. 9 (11): 2431.

Extension Toxicology Network. 1993. A Pesticide Information Project of Cooperative Extension Offices of Cornell University, Michigan State University, Oregon State University, and University of California at Davis.  
<http://pmep.cce.cornell.edu/profiles/extoxnet/metiram-propoxur/metsulfuron-methyl-ext.html>

FAO (Food and Agriculture Organization of the United Nations).1998.Report of the Fifth External Program and Management Review of International Rice Research Institute (IRRI).Natural Resources Management and Environment Department.<http://www.fao.org/wairdocs/tac/x5801e/x5801e00.htm#Contents>

Galinato, M.I., Moody, K. and Piggin, C.M., 1999. *Upland rice weeds of South and Southeast Asia.* Int. Rice Res. Inst..

Harker, K.N. and O'donovan, J.T., 2013. Recent weed control, weed management, and integrated weed management. *Weed Technology*, 27(1), pp.1-11.

*Harvey, W.A.; Robblns, W.W. 1947. 2,4-D as a Weed Killer. University of California at Berkeley.*

Holm L. G., Plucknett D. L., Pancho J. V., Herberger J. P. 1991. The world's worst weeds.Distribution and Biology.East-West Center by the University Press. Hawaii.

Indonesia Investment. 2016. Rice. <http://www.indonesia-investments.com/business/commodities/rice/item183?>

International Agency for Research on Cancer. 2015. IARC Monographs evaluate DDT, lindane, and 2,4-D. World Health Organization. Press Release N° 236.

IRIR's Rice Almanac, 2013. <http://ricepedia.org/Indonesia>, read 21/06/2016

Jabran, K. and Chauhan, B.S., 2015. Weed management in aerobic rice systems. *Crop Protection*, 78, pp.151-163.

Jamilah. 2013. IlmuGulmah. Agriculture Faculty, Universitas Tamansiswa. Padang. Sumatra Barat. Indonesia

Janiya, J.D. and Moody, K., 1989. Weed populations in transplanted and wet-seeded rice as affected by weed control method. *International Journal of Pest Management*, 35(1), pp.8-11.

Karim, R.S., Man, A.B. and Sahid, I.B., 2004. Weed problems and their management in rice fields of Malaysia: An overview. *Weed Biology and Management*, 4(4), pp.177-186.

Khanh, T.D., Xuan, T.D., Chin, D.V., Chung, I.M., Abdelghany, E.A. and Tawata, S., 2006. Current status of biological control of paddy weeds in Vietnam. *Weed Biology and Management*, 6(1), pp.1-9.

Khush, G.S., 1997. Origin, dispersal, cultivation and variation of rice. In *Oryza: From molecule to plant* (pp. 25-34). Springer Netherlands.

Kim PV, Hoang VT, Thuy TTT, Hoa LH, Hoai NB. 1981. Host range of *Pyricularia oryzae* and *Thanatephorus cucumeris* in the Vietnamese Mekong Delta. Int. Rice Res. News. p 6-10.

Linares, O.F., 2002. African rice (*Oryzopsis glaberrima*): history and future potential. *Proceedings of the National Academy of Sciences*, 99(25), pp.16360-16365.

Maps of World. World's top 10 Rice Producing Countries. <http://www.mapsofworld.com/world-top-ten/rice-producing-countries>.

Mishra, J.S. and Singh, V.P., 2012. Tillage and weed control effects on productivity of a dry seeded rice-wheat system on a Vertisol in Central India. *Soil and Tillage Research*, 123, pp.11-20.

Moody K. 1989. Weeds reported in rice in South and Southeast Asia. Manila (Philippines): International Rice Research Institute.

Mubeen, K., Nadeem, M.A., Tanveer, A. and Jhala, A.J., 2014. Effects of seeding time and weed control methods in direct seeded rice (*Oryza sativa* L.). *The Journal of Animal & Plant Sciences*, 24(2), pp.534-542.

Nguyen HuuTruc. 2012. *Giaotinh co dai*. Ho Chi Minh City University of Agriculture and Forestry, Viet Nam.

Pesticide Properties DataBase. 2016. University of Hertfordshire.  
<http://sitem.herts.ac.uk/aeru/ppdb/en/Reports/1145.htm>

Rao, A.N., Johnson, D.E., Sivaprasad, B., Ladha, J.K. and Mortimer, A.M., 2007. Weed management in direct-seeded rice. *Advances in Agronomy*, 93, pp.153-255.

Renu, S., Thomas, C.G. and Abraham, C.T., 2000. Stale seedbed technique for the management of *Sacciolepis interrupta* in semi-dry rice. *Indian Journal of Weed Science*, 32(3/4), pp.140-145.

Rice Knowledge Bank. Chemical weed control.  
[http://www.knowledgebank.irri.org/index.php?option=com\\_zoo&view=item&layout=item&Itemid=710](http://www.knowledgebank.irri.org/index.php?option=com_zoo&view=item&layout=item&Itemid=710)

Sindhu, P.V., Thomas, C.G. and Abraham, C.T., 2010. Seedbed manipulations for weed management in wet seeded rice. *Indian Journal of Weed Science*, 42(3/4), pp.173-179.

Singh, Y., Singh, V.P., Singh, G., Yadav, D.S., Sinha, R.K.P., Johnson, D.E. and Mortimer, A.M., 2011. The implications of land preparation, crop establishment method and weed management on rice yield variation in the rice-wheat system in the Indo-Gangetic plains. *Field Crops Research*, 121(1), pp.64-74.

Solopos.com. 2015. *Beras Jenis Baru Petani Magelang Produksi Beras Rendah Gula*. Creative Information Network. <http://www.solopos.com/2015/03/11/beras-jenis-baru-petani-magelang-produksi-beras-rendah-gula-583846>

Thakur, R., Majhi, S., Upasani, R.R., Pal, S.K. and Singh, M.K., 2011. Impact of integrated weed management on dry matter partitioning and productivity of direct seeded rice. Pp 45-54.

TnauAgritech Portal, Agriculture (2016).  
[http://agritech.tnau.ac.in/agriculture/agri\\_weedmgt\\_rice.html](http://agritech.tnau.ac.in/agriculture/agri_weedmgt_rice.html)

Tshewang, S., Sindel, B.M., Ghimiray, M. and Chauhan, B.S., 2016. Weed management challenges in rice (*Oryza sativa L.*) for food security in Bhutan: A review. *Crop Protection*, 90, pp.117-124.

Tu et al. 2001. Weed Control Methods Handbook. The Nature Conservancy. University of Georgia Center for Invasive Species and Ecosystem Health.

Tuat, N. V., Shane, H. and Bruce, A., 1997. *Prospect for biological weed control in Viet Nam*, Report on Regional Workshop on Improved Weed Management in Rice, FAO Publication, 1997.

United States Environmental Protection Agency. Ingredients Used in Pesticide Products 2,4-D. <http://www2.epa.gov/EPA>

Van Nguyen, N. and Ferrero, A., 2006. Meeting the challenges of global rice production.4. Pp 1-9.

Vaughan, D.A., 1994. *The wild relatives of rice: a genetic resources handbook*. Int. Rice Res. Inst.