

## DAFTAR PUSTAKA

- Alexander JM, Leveno KJ, Hauth J, Landon MB, Thom E, Spong CY, *et al.* (2006). Fetal injury associated with cesarean delivery. *ACOG*, 108 (4): 886-887.
- Annisa SA (2011). Faktor-faktor risiko persalinan seksio sesarea di RSUD Dr. Adjidarmo Lebak pada bulan oktober-desember 2010. Universitas Islam Negeri Syarif Hidayatullah Jakarta. Skripsi.
- Armstrong C (2011). ACOG updates recommendations on vaginal birth after previous cesarean delivery. *AFP*, 83 (2): 201.
- Balitbang Kemenkes RI (2013). Riset kesehatan dasar 2013. Jakarta: Balitbang Kemenkes RI.
- Bogaert LJV (2004). Mode of delivery after one cesarean section. *IJGO*, 87: 9-13.
- Caughey AB, Robinson JN, Norwitz ER (2008). Contemporary diagnosis and management of preterm premature rupture of membranes. *Rev Obstet Gynecol*, 1(1): 11-22.
- Cecatti JG, Pires HMB, Faundes A, Duarte Osis MJ (2005). Factors associated with vaginal birth after previous cesarean section in Brazilian women. *Rev Panam Salud Publica*, 18 (2): 107-113.
- Chhabra S, Arora G (2006). Delivery in women with previous cesarean section. *J Obstet Gynecol India*, 56 (4): 304-307.
- Cunningham FG, Bangdiwala S, Brown SS, Dean TM, Frederiksen M, Rowland, *et al.* (2010). NIH consensus development conference statement: Vaginal birth after cesarean: New insights. *Obstetric & Gynecology*, 115 (6): 1279-1295.
- Cunningham FG, Leveno KJ, Bloom SL, Spong CY, Dashe JS, Hoffman BL, *et al.* (2014). *Williams obstetrics*, 24th ed. New York: McGraw-Hill Education, pp: 587-608.
- Dahlan MS (2009). Besar sampel dan cara pengambilan sampel. Jakarta: Salemba Medika, pp: 46-60.
- Dahlan MS (2011). *Statistik untuk kedokteran dan kesehatan*. Jakarta: Salemba Medika, pp: 16-23.
- Dinas K, Mavromatidis G, Dovas D, Giannoulis C, Tantanasis T, Loufopoulos A, *et al.* (2008). Current caesarean delivery rates and indications in a major public hospital in northern Greece. *ANZJOG*, 48: 142-146.

- Dodd JM, Crowther CA, Hiller JE, Haslam RR, Robinson JS (2007). Birth after caesarean study – planned vaginal birth or planned elective repeat caesarean for women at term with a single previous caesarean birth: Protocol for a patient preference study and randomized trial. *BMC Pregnancy and Childbirth*, 7 (17): 3.
- Fagerberg MC, Marsal K, Ekstrom P, Kallen K (2012). Indications for first caesarean and delivery mode in subsequent trial of labour. *PPE*, 27: 72–80.
- Gabbe SG, Niebyl JR, Simpson JL, Landon MB, Galan HL, Jauniaux ERM, *et al.* (2012). *Obstetrics: Normal and problem pregnancies*. Edisi ke 6. China: Elsevier Saunders, p: 445.
- Gates S, McHarg KM, Hurley P (2001). Effects of surgical techniques of caesarean section on maternal health. Cambridge University Press, 12 (2): 105-137.
- Gilliam M (2006). Cesarean delivery on request: Reproductive consequences. *Semin Perinatol*, 30: 257-260.
- Goel SS, Tiwari M, Hariharan C, Shrivastava DS (2013). Outcome of post caesarean pregnancy and comparison of maternal and foetal outcome following vaginal birth versus repeat caesarean section in a rural hospital. *IJRCOG*, 2(1): 16-22.
- Halimatusyaadiah S, Trisnawati NN (2014). Hubungan persalinan bekas seksio sesarea dengan cara persalinan, luaran janin. *MBI*, 8 (6): 54-62.
- Institute of Obstetricians and Gynaecologists, Royal College of Physicians of Ireland (2011a). Delivery after previous caesarean section. Clinical Strategy and Programmes Directorate, Health Service Executive, 5 (1): 10.
- Institute of Obstetricians and Gynaecologists, Royal College of Physicians of Ireland (2011b). The diagnosis and management of pre-eclampsia and eclampsia. Clinical Strategy and Programmes Directorate, Health Service Executive; 3(1): 14.
- Islam A, Ehsan A, Arif S, Murtaza J, Hanif A (2011). Evaluating trial of scar in patients with a history of caesarean section. *NAJMS*, 3 (4): 201-205.
- Karalti MO, Tinar S, Ozturk NT, Ozturk DC (2012). A case with vesicouterine fistula: Mini review. *Arch Gynecol Obstet*, 285: 667–670.
- Kennare R, Dip G, Tucker G, Heard A, Chan A (2007). Risks of adverse outcomes in the next birth after a first caesarean delivery. *ACOG*, 109 (2): 275.

- Khalil A, Syngelaki A, Maiz N, Zinevich Y, Nicolaides KH (2013). Maternal age and adverse pregnancy outcome: A cohort study. *Ultrasound Obstet Gynecol*, 42: 634–643.
- Kurniawan (2005). Gambaran kasus persalinan seksio sesarea di bagian Obstetrik dan Ginekologi Perjan RS. dr. M. Djamil Padang tahun 2003. Universitas Andalas Padang. Skripsi.
- Landon MB, Hauth JC, Leveno KJ, Spong CY, Leindecker S, Varner MW, *et al.* (2004). Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *NEJM*, 351 (25): 2583-2588.
- Landon MB, Leindecker S, Spong CY, Hauth JC, Bloom S, Varner MW, *et al.* (2005). The MFMU cesarean registry: Factors affecting the success of trial of labor after previous cesarean delivery. *AJOG*, 193: 1016-23.
- Li Z, McNally L, Hilder L, Sullivan EA (2011). Australia's mothers and babies 2009. Perinatal statistics series no. 25. Cat. no. PER 52. Sydney. AIHW National Perinatal Epidemiology and Statistics Unit, 40-43.
- Liu S, Liston RM, Joseph KS, Heaman M, Sauve R, Kramer MS (2007). Maternal mortality and severe morbidity associated with low-risk planned cesarean delivery versus planned vaginal delivery at term. *CMAJ*, 176 (4): 455-460.
- Manuaba IBG (2001). Kapita selekta penatalaksanaan rutin obstetri ginekologi dan KB. Jakarta: EGC, p: 206.
- Manuaba IBG, Manuaba IAC, Manuaba IBGF (2003). Pengantar kuliah obstetri. Jakarta: EGC, p: 876.
- Marshall NE, Fu R, Guise JM (2011). Impact of multiple cesarean deliveries on maternal morbidity: A systematic review. *AJOG*, 205:262e1-8.
- Martel MJ, Mackinnon CJ (2005). Guidelines for vaginal birth after previous caesarean birth. *JOGC*, 27 (2): 164–174.
- Minkoff H, Chervenak FA (2003). Elective primary cesarean delivery. *NEJM*, 348 (10): 946-950.
- Mohamed HA (2015). Influence of the indications and outcomes of previous cesarean section on planned mode of delivery in women with one previous cesarean section. University of Nairobi. Thesis.
- Mulyawati I, Azam M, Ningrum DNA (2011). Faktor tindakan persalinan operasi sectio caesarea. *KEMAS*, 7 (1): 14-21.
- Mylonas I, Friese K (2015). Indications for and risks of elective cesarean section. *Dtsch Arztebl Int*, 112: 489–495.

- Olagbuji B, Ezeanochie M, Okonofua F (2010). Predictors of successful vaginal delivery after previous caesarean section in a Nigerian tertiary hospital. *Journal of Obstetrics and Gynaecology*, 30 (6): 582–585.
- Pereira APE, Tharaux CD, Pereira MN, Saucedo M, Colle MHB, Leal MDC (2016). Caesarean delivery and postpartum maternal mortality: A population-based case control study in Brazil. *PLoS ONE*, 11 (4): 1-13.
- Prawirohardjo S (2008). Ilmu kebidanan sarwono prawirohardjo. Edisi ke IV. Jakarta: Bina Pustaka Sarwono Prawirohardjo, p: 616.
- Queensland Clinical Guidelines (2015). Vaginal birth after caesarean section (VBAC). Queensland Clinical Guideline. MN15.12-V4-R20.
- Rajamaheswari N, Chhikara AB (2012). Vesicouterine fistulae: Our experience of 17 cases and literature review. *Int Urogynecol J*.
- Rini WP (2014). Gambaran indikasi dan karakteristik ibu bersalin secara seksio sesarea di RSI Jemursari Surabaya. Universitas Nahdlatul Ulama Surabaya. Skripsi.
- RCOG (2005). WHO systematic review of maternal mortality and morbidity: The prevalence of uterine rupture. *BJOG*, 112: 1221–1228.
- Royal College of Obstetricians and Gynaecologists (2015). Birth after previous caesarean birth. RCOG Green-top Guideline No. 45.
- Rowe T (2014). Placenta previa. *JOGC*, 36 (8): 667–668.
- Saxena R (2010). *Bedside obstetrics and gynecology*. New Delhi: Jaypee Brothers Medical Publishers, p: 121.
- Souza JP (2010). Caesarean section without medical indication increases risk of short-term adverse outcome for mothers: The 2004-2008 WHO global survey on maternal and perinatal health. *BMC Medicine*, 8 (1): 71.
- Stavrou EP, Ford JB, Shand AW, Morris JM, Roberts CL (2011). Epidemiology and trends for caesarean section births in New South Wales, Australia: A population-based study. *BMC Pregnancy and Childbirth*, 11 (8): 3.
- Statistics Indonesia (Badan Pusat Statistik—BPS) and Macro International (2008). *Indonesia demographic and health survey 2007*. Calverton, Maryland, USA: BPS and Macro International.
- Sumelung V, Kundre R, Karundeng M (2014). Faktor-faktor yang berperan meningkatnya angka kejadian *sectio caesarea* di rumah sakit umum daerah liun kendage tahuna. *Ejournal keperawatan (e-Kp)*, 2 (1): 1-7.

Thilagavathi P, Geetha V (2017). Study of vaginal delivery following cesarean section. IOSR-JDMS, 16 (1): 7-11.

Villar J, Valladares E, Wojdyla D, Zavaleta N, Carroli G, Velazco A, *et al.* (2006). Caesarean delivery rates and pregnancy outcomes: The 2005 WHO global survey on maternal and perinatal health in Latin America. ProQuest, 367 (9525): 1825-1826.

Women's Health Committee (2015). Birth after previous cesarean section. RANZCOG; C Obs 38.

Zaitoun MM, Nour Eldin SA, Mohammad EY (2013). A Prediction score for safe and successful vaginal birth after cesarean delivery: A prospective controlled study. J Women's Health Care, 2 (3): 1-7.

