

DAFTAR PUSTAKA

- Abboud O. Hypertension and pregnancy. Heart views.vol.1 no.8 june-august 2000:309-313.
- Achan.V.et al. Asymmetric dimethylarginine causes hypertension and cardiac dysfunction in humans and is actively metabolized by dimethylargininedimethyl aminohydrolase, Arteriosclerosis, Thombosis, and Vascular Biology, vol. 23, no. 8, pp. 14551459, 2003.
- Adiga U, D'souza V, Kamath A, Mangalore N. 2007. Antioxidant activity and lipid peroxidation in preeclampsia. J Chin Med Assoc; 70 (10): 435-438.
- Agrawal S, Walia GK. Prevalence and risk factors for Pre-eclampsia in Indian women: a national cross sectional study. Population Association of America 2014 Annual Meeting Program2014. p. 1-23.
- Ahmed.A, Dunk.C, Ahmed.S, Khaliq.A.Regulation of placental vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF) and soluble Flt-1 by oxygen--a review. *Placenta*. 2000 Mar-Apr;21 Suppl A:S16-24
- Akolekar R, Zaragosa E, Poon LCY, Pepes S, Nicolaides KH (2008): Maternal serum placental growth factor at 11+0 to 13+6 weeks of gestation in the prediction of pre-eclampsia. *Ultrasound Obstet Gynecol*. 32: 732-739.
- Akolekar R, Cruz JD, Foidart JM, Munaut C, Nicolaides KH : Maternal serum placental soluble fms-like tyrosine kinase-1 and free vascular endothelial growth factor at 11 to 13 weeks of gestation in pre-eclampsia. *Prenat Diagn* 2010;30:191-7.
- Amelda; Gambaran Karekteristik Ibu Hamil dengan Preeklampsia, 2009.fileswordpress.com. Diakses 15 Maret 2013.
- Anas MN. Hubungan Lingkar Lengan Atas (Lila) pada Ibu Hamil dengan Angka Kejadian Preeklampsia di RS. PKU Muhammadiyah Surakarta. Surakarta: Universitas Muhammadiyah Surakarta; 2013
- Anonim. Isoprostane. Wikipedia the free encyclopedia, available at <http://en.wikipedia.org/wiki/Isoprostane>. Last update 28 May 2010.
- Anonim. Oxysterol. Wikipedia the free encyclopedia, available at <http://en.wikipedia.org/wiki/Oxysterol>. Last update 15 May 2010.
- Anonim. Malondialdehyde. Wikipedia the free encyclopedia, available at <http://en.wikipedia.org/wiki/Malondialdehyde>. Last update 18 August 2009.
- Anonim. 4-hydroxynonenal. Wikipedia the free encyclopedia, available at <http://en.wikipedia.org/wiki/4-Hydroxynonenal>. Last update 2 April 2010.

Anonim. *Lipid peroxidation*. Wikipedia the free encyclopedia, available at http://en.wikipedia.org/wiki/Lipid_peroxidation. Last update 20 April 2010.

Anonim. *Lipid*. Wikipedia the free encyclopedia, available at <http://en.wikipedia.org/wiki/Lipid>. Last update 30 may 2010.

Arias.F et al. Preterm Labor in Practical Guide to high risk pregnancy and delivery. Mosby Year book.P.71-99.

Arias F, Daftary S, Bhide A. Hypertensive Disorders in Pregnancy. In: Arias F, Daftary S, Bhide A, editors. Practical Guide to High-Risk Pregnancy and Delivery: A South Asian Perspective. New Delhi: Elsevier; 2008. p. 397-45.

Artikasari K,2009. Hubungan antara Primigravida dengan Angka Kejadian Preeklamsia/Eklamsia di RSUD Dr. Moewardi Surakarta Periode 1 Januari – 31 Desember 2008. Universitas Muhammadiyah Surakarta

Atis A et al. Troponin I and Homocysteine levels in mild and sever preeclampsia. *Clin Exp Obstet Gynecol*2010; 37(1) :21-3.

Badria LF, Amarin ZO. Pre-eclampsia is it a different disease in primiparous and multiparous women. *Arch Gynecol Obstet*.2005;273:26-31.

Barden, A., Ritchie, J., Walters, B., Michael, C., Rivera, J., Mori, T., et al. 2001. Study of Plasma Factors Associated With Neutrophil Activation and Lipid Peroxidation in Preeclampsia. *Journal of Hypertension*, 38:803-808.

Bargale.A, Jayashree.VU, Dhiraj.JT, Nitin.N, Rakesh.M, Aparna.S: Serum hsCRP and Uric acid as indicator of severity in Preeclampsia.*Int.J of Pharma and Bio science*.Vol.2.issue 3. July-Sept 2011.

Becker E. Novel therapies for the preventive and treatment of Preeclampsia and eclampsia. *Path.Org*,p:10.May 2014.

BilanoVL, Ota E, Ganchimeg T, Mori R, Souza JoP. Risk Factors of Preeclampsia /Eclampsia and Its Adverse Outcomes in Low- and Middle-Income Countries: A WHO Secondary Analysis. *PLOS ONE*. 2014;9(3):1-9.

Benaitreau D, Marie-Noëlle Dieudonné, Esther Dos Santos, Marie-Christine Leneveu, Philippe de Mazancourt and René Pecquery : Antiproliferative Effects of Adiponectin on Human Trophoblastic Cell Lines JEG-3 and BeWo. *Biology of Reproduction* June 1, 2009 vol. 80 no. 6 1107-1114

Berg C.J, et al. Preventability of Pregnancy-Related Deaths Results of a State-Wide Review. *Obstetric & Gynaecology*, Vol. 106, No. 6, December 2005. pp : 1228- 34.

Block, G., Dietrich, M., Norkus, E.P., Morrow, J.D., Hudes, M., Bette, C., et al.

2002. Factors Associated with Oxidative Stress in Human Population. *American Journal of Epidemiology*, 156: 274-85.

Bobak, Lowdermilk, Jensen. Buku ajar keperawatan maternis. ed.4. EGC, Jakarta. 2004.

Boger R.H, S. M. Bode-Boger, A. Szuba et al., "Asymmetric dimethylarginine (ADMA): a novel risk factor for endothelial dysfunction: its role in hypercholesterolemia," *Circulation*, vol.98, no. 18, pp. 1842–1847, 1998.

Boyle P.J. What are the effects of peroxisome proliferator-activated receptor agonists on adiponectin, tumor necrosis factor-alpha, and other cytokines in insulin resistance? *Clinical cardiology*. Volume 27, Issue Supplement S4, pages 11–16, July 2004. Article first published online: 5 JUL 2007.

Brown G C, Borutaite V, Nitric Oxide, Mitochondria, and Cell Death. *IUBMB Life*, 52:189–195, 2001

Burton GJ, Yung HW, Davies TC, Charnock-Jones DS. 2009. Placental Endoplasmic Reticulum Stress and Oxidative Stress in the Pathophysiology of Unexplained

Intrauterine Growth Restriction and Early Onset Preeclampsia. *Placenta* 30, Supplement A, Trophoblast Research; 23: S43–S48.

Buke Bans, Topcu HO, Ustun YE, Darusman N: Comparasion of serum maternal adiponectin concentration in women with isolated intraurine growth retardation and intra uterine growth retardation concomitant with preeclampsia. *J Turk Ger Gynecol Assoc*. 2014;15: 173-6.

Bujold E, S. Robege, Y. Lacasse et al., "Prevention of preeclampsia and intrauterine growth restriction with aspirin started in early pregnancy: a meta-analysis," *Obstetrics and Gynecology*, vol. 116, no. 2, pp. 402–414, 2010.

Carolina Valdiviezo, Vesna D, Pamela Ouyang. Preeclampsia and hypertensive disease in pregnancy; Their contributions to cardiovascular risk. *Clin Cardiol* 35 (3) 2002. p:160-162

Cedergren, M., 2007. *Total Weight Gain and Pattern of Weight Gain in Pregnancy*. Division of Obstetric Gynecology, Department of Molecular and Clinical Medicine, Faculty Science, Sweden University.

Chappel, S. Morgan, L. 2006. Searching for genetic clues to the causes of preeclampsia. *Clinical science* 110: 443-458

Chang Jj, Muglia LJ, Macones GA. Association of early onset preeclampsia in first pregnancy with normotensive second pregnancy outcomes. A population based study. *BJOG*. 2010;117(8):946-953

Chen, B.; Longtine, M.S.; Nelson, D.M. Hypoxia induces autophagy in primary human trophoblasts. *Endocrinology* 2012, 153, 4946–4954

Cindrova-Davies, T. 2009. Gabor Than Award Lecture 2008: Pre-eclampsia from Placental Oxidative Stress to Maternal Endothelial Dysfunction. *Placenta*, 23: S55-65.

Corwin, Elizabeth, J., 2000. Sistem Kardiovaskular. Dalam: Buku Saku Patofisiologi. Ed 3. Jakarta: Penerbit Buku Kedokteran EGC, 358-359.

Cracowski, J.L., Baguet, J.P., Ormezzano, O., Bessard, J., Stanke-Labesque, F., Bessard, G., dkk. 2003. Lipid Peroxidation is Not Increased in Patients with Untreated Mild-to-moderate Hypertension. *Journal of Hypertension*, 41:286 –288.

Cunningham FG et al. Hypertensive disorder in pregnancy. William Obstetrics 22th. United State of America . Mc Graw-Hill, 2009: 761-808

Cunningham FG, Leveno KJ, Bloom SL, Spong CY, Dashe JS, Hoffman BL, et al. Hypertensive Disorders. In: Cunningham FG, Leveno KJ, Bloom SL, Spong CY, Dashe JS, Hoffman BL, et al., editors. Williams Obstetrics 24th. United States: McGraw-Hill Education; 2014. p. 728-79.

Czajkowska E.P, B. Marciak, Z. Kimber-Trojnar, B. Leszczynska-Gorzelak, and J. Oleszczuk, “Nitric oxide in normal and preeclamptic pregnancy,” Current Pharmaceutical Biotechnology, vol. 12, no. 5, pp. 743–749, 2011.

Dalle Donne, I., Rossi, R., Colombo, R., Giustarini, D., Milzani, A. 2006. Biomarkers of Oxidative Damage in Human Disease. *Clinical Chemistry*, 52(4):601-623.

D'Anna R, Baviera G, Corrado F, Giordano D, Di Benedetto A, Jasonni VM. Plasma Adiponectin Concentration in Early Pregnancy and Subsequent Risk of Hypertensive Disorders. *Obstet Gynecol* 2005; 106:340–344.

Danforth.Obstetric and Gynecology.10 th Ed.2011

Departemen Kesehatan RI. Statistik Rumah Sakit di Indonesia. [cited 2006]. Available from URL: <http://www.yanmedikdepkes.net/statistik.2006>.

Dekker GA, Sibai BM: Etiology and pathogenesis of preeclampsia: current concepts. Am J Obstet Gynecol. 1998 Nov;179(5):1359-75.

Dikenoy E, O. Balat, S. Pence, A. Balat, M. Cekmen, and M. Yurekli, “Th changes of plasma malondialdehyde, nitric oxide and adrenomedullin levels in patients with preeclampsia,” *Hypertension in Pregnancy*, vol. 28, no. 4, pp. 383–389, 2009.

Dini.DJ,Bambang Kuntarto.Hubungan umur dan paritas ibu dengan kejadian preeklamsi.E-Jurnal Obstretika Vol 1 no 1 (Januari-Juni, 2013)

Dominique .R,Ferdinand L,Megan K,Richard B,Ravi R,Steven K,Walter J,Bruce L,Loe ell S,Ruben M,Dinesh A: Troponin I level in Preeclampsia patient.The American Journal Medicine.VOLUME 120,issue 9, p 819, september 2007.

Dotan Y, Lichtenberg D, Pinchuk I. 2004. Lipid peroxidation cannot be used as a universal criterion of oxidative stress. Progress in Lipid Research; 43: 200–227.

English FA, Kenny LC, McCarthy FP. Risk factors and effective management of preeclampsia. Dove Press Journal: Integrated Blood Pressure Control 2015;8:7-12.

Fam, S.S., Morrow, J.D. 2003. The Isoprostanes: Unique Products of Arachidonic Acid Oxidation : A Review. *Current Medicinal Chemistry*, 10:1723-1740.

Farer. Preeclampsia: an excessive maternal inflammatory response to pregnancy.Am J Obstet Gynecol.1999 Feb;180(2Pt1):499-506.

Farooqui, A.A., Horrocks, L.A. 2007. Nonenzymic Metabolites of Arachidonate and Docosahexanoate in Brain. In :*Glycerophospholipids in Brain : Phospholipases A2 in Neurological Disorders*. Springer science + Bussiness.

Fasshauer M, and Paschke R : Regulation of adipocytokines and insulin resistance. Diabetologia 2003; 46: 1594–1603.

Gaskins AJ, et al. Biocyte Study Group.: Adherence to a Mediterranean diet and plasma concentrations of lipid peroxidation in premenopausal women associated with plasma adiponectin concentrations in diabetic women Am J Clin Nutr. 2010 Dec; 92(6):1461-7.

Gatot S, Lawrence, Irawan Yusuf, Ani Miaya, Syarifuddin Wahid. Kadar adiponektin rendah pada toleransi glukosa terganggu: implikasi vaskuler awal. Medical Faculty of Hasanuddin University' June 11' 2006.

Gilstrap LC, Ramin MS, Diagnosis and Management of preeklampsia and eclampsia. American College of Obstetricians and Gynaecologist 2002; 67-159.

Girouard J et al. Previous hypertensive disease of pregnancy is associated with alterations of markers of insulin resistance. *Hypertension*, 2007, 49:1056–1062.

Goldstein et al. A novel adipokin Linking Adipocyte and Vaskular Function. J Clin Endocrinol Metab 2004.

Gratacos E, Casals E, Deulofeu R, Cararach V,Pedro A, Fortuny A. Lipid peroxide and vitamin E patterns in pregnant women with different types of hypertension in pregnancy. American Journal Obstetrics Gynecology, 1998;178; 1072-1076.

Grill S, Rusterholz C, Dalenbach RZ, Tercanli S, Holzgreve W, Hahn S. 2009. Potential markers of preeclampsia-a review. Reprod biol endocrinol; 7: 70.

Guerrier G, Oluyide B, Keramarou M, Grais RF. Factors associated with severe preeclampsia and eclampsia in Jahun, Nigeria. International Journal of Women's Health. 2013 (5):509-13.

Gupta, S., Agarwal, A., Sharma, R.K. 2005. The Role of Placenta Oxidative Stress and Lipid Peroxidation in Preeclampsia. *Obstetrical and Gynecological Survey*, 60(12):807-816.

Gupta, S., Aziz, N., Sekhon, L., Agarwal, R., Mansour, G., Li, J., Agarwal, A. 2009. Lipid Peroxidation and Antioxidant Status in Preeclampsia, A Systematic Review. *Obstetrical and Gynecological Survey*, 64(11):750-759.

Hanifa W: Ilmu Kebidanan. Ed. ketiga. Cetakan keempat. Jakarta. Yayasan Bina Pustaka Sarwono Prawiroharjo. 1997; 24: 281-301.

Harsem, N.K., Roald, B., Braekke, K., Staff, A.C. 2007. Acute Atherosclerosis in Decidual Tissue: Not Associated with Systemic Oxidative Stress in Preeclampsia. *Placenta*, 28 : 958-964.

Hawfield.A and Freedman B.I ; Pre-eclampsia: the pivotal role of the placenta in its pathophysiology and markers for early detection. *Ther Adv Cardiovasc Dis*. 2009 Feb; 3(1): 65-73.

Henriksen, T. 2000. The Role of Lipid Oxidation and Oxidative Lipid Derivatives in the Development of Preeclampsia. *Seminars in Perinatology*, 24(1):29-32.

Hermanto Quedarusman 2 John Wantania 2 Juneke J. Kaeng. Hubungan indeks massa tubuh ibu dan peningkatan berat badan saat kehamilan dengan preeklampsia. *Jurnal e-Biomedik (eBM)*, Volume 1, Nomor 1, Maret 2013, hlm. 305-311

Hermenegildo, C., Garcia-Martinez, M.C., Tarin, J.J., Cano, A. 2002. Estradiol Reduces F2 α -Isoprostane Production in Cultured Human Endothelial Cells. *American Journal Physiological - Heart and Circulation Physiology*, 283:H2644-H2649.

Hybel CA, Mc Laughlin MKA, Evans RWA, Hauth BAA, Sims CJA, Roberts JMA. Fasting serum triglycerides, free fatty acids, and malondialdehyde are increased in preeclampsia, are positively correlated, and decreased within 48 hours post partum. *American Journal of Obstetrics and Gynecology*, 1996; 174:975-982.

Hubel CA. Oxidative stress in the pathogenesis of preeclampsia. *Journal of Experimental Biology and Medicine*; 1999;222-235.

Hubel CA. Dyslipidemia and Pre-eclampsia. In *Preeclampsia Etiology and Clinical Practice*, New York: Cambridge University Press 2007 : 164-82.

Hung JH. Oxidative Stress and Antioxidant in Preeclampsia. *J Chin Med Assoc* 2007; 70(10).

Hung, T.H., Bruton, G.J. 2006. Hypoxia and Reoxygenation : A Possible Mechanism for Placental Oxidative Stress in Preeclampsia. *Taiwanese Journal of Obstetrics and Gynecology*, 43(3):189-200.

Huppertz.B. Placental origin of preeclampsia challenging the current hypothesis. *Hypertension* 2008; 51:970-5.

Ichida K et al. Plasma adiponectin concentrations and placental adiponectin expression in pre-eclamptic women. *Gynecological Endocrinology*, 2007, 23:238–243.

Ishihara, O., Hayashi, M., Osawa, H., Kobayashi, K., Takeda, S., Vessby, B., et al. 2004. Isoprostanes, Prostaglandins and Tocopherols in Pre-eclampsia, Normal Pregnancy and Non-pregnancy. *Free Radical Research*, 38(9):913 918.

Ika Fidianingsih. Sel lemak dan peranannya dalam penyakit. Fakultas Kedokteran Universitas Islam Indonesia'2000.

Israel Hendlar, The levels of leptin, adiponectin, and resistin in normal weight, overweight, and obese pregnant women with and without preeclampsia, *American Journal of Obstetrics and Gynecology*, 2005; 193 -979.

Janicka, M., Kot-Wasik, A., Kot, J., Namieśnik, J. 2010. Isoprostanes-Biomarkers of Lipid Peroxidation: Their Utility in Evaluating Oxidative Stress and Analysis. *International Journal of Molecular Sciences*, 11: 4631-4659.

Jaffe et all. A Biomarker Odyssey. *Clin Chim Acta* 1999;284:197-211.

Janero DR.1990. *Malondialdehyde and thiobarbituric acid reactivity as diagnosis indices of lipid peroxidation and peroxidative tissue injury*. FreeRadical Biology & Medicine; 9: 515-540.

Jones DL. Dasar-dasar Obstetric dan Ginekologi. 6 ed. Jakarta: Hipokrates; 2001.

Kabukçu.A.S ; Önderoglu.L ; Laleli.Y ; Durukan.T. Women with Elevated Second Trimester HumanChorionic Gonadotropin Level Are at IncreasedRisk for Preeclampsia. *Turk J Med Sci*.28 (1998) 273-276,

Kadowaki T and Toshimasa Yamauchi. Adiponectin and Adiponectin receptor. *Endocrine Reviews*, May 2005, 26(3):439–451

Kato K, Osawa H, Ochi M, et al : Serum total and high molecular weight adiponectin levels are correlated with the severity of diabetic retinopathy and nephropathy. *Clin Endocrinol* 2008; 68: 442–449.

Kajantie E, Kaaja R, Ylikorkala O, Andersson S, Laivuori H. Adiponectin concentrations in maternal serum: elevated in preeclampsia but unrelated to insulin sensitivity. *Journal of the Society for Gynecologic Investigation*. 2005; 12:433–9.

Karyati S. Faktor-Faktor yang Berkontribusi Pada Kejadian Pre Eklampsia. JIKK. 2014;5

Kelley D & Mandarino L: Fuel selection in human skeletal muscle in insulin resistace: A Rexamination Diabetes.2004,49.677-83.

Kementerian Kesehatan RI. Survey kesehatan rumah tangga;angka kematian ibu dan anak. 2011.

Khalil A, N. J.Cowans, K. Spencer, S. Goichman, H. Meiri.and K.Harrington : First-trimester markers for the prediction of pre-eclampsia in women with a-priori high risk. *Ultrasound Obstet Gynecol* 2010; 35: 671–679 . Published online in Wiley InterScience.

Khaliq F, Singhal U, Arshad Z, Hossain MM. 1999. Thyroid functions in pre-eclampsia and its correlation with maternal age, parity, severity of blood pressure and serum albumin. *Indian J Physiol Pharmacol* 43: 193–198.

Khan K.S, Daniel Wojdyla, Lale Say, A Metin Gümezoglu, Paul F A Van Look: WHO analysis of causes of maternal death: a systematic review. *Lancet* 2006; 367: 1066–74.

Kim SY, Ryu HM,Yang JH,Kim MY,Han JY,Kim JO.Increase sFlt-1 to PLGF ratio in women who subsequently develop preeclampsia.*J Korean Med Sci*.2007;22:873-7.

Kirwan, J.P., S.H.D. Mouzon, J. Lepercq, J.C. Challier and L. Huston-Presley et al: TNF- α is a predictor of insulin resistance in human pregnancy. *Diabetes*. 2002;51: 2207-2213.

Koga M, Kasayama S (2010). "Clinical impact of glycated albumin as another glycemic control marker". *Endocrine Journal* 57 (9): 751–762.

Koopmans CM,D.Bijlenga,H.Groen et al., "Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre, open-label randomised controlled trial," *The Lancet*, vol. 374, no. 9694, pp. 979–988, 2009.

Krauss.T,Pauer.H U, Augustin H G. Prospective Analysis of Placenta Growth Factor (PIGF) Concentrations in the Plasma of Women with Normal Pregnancy and Pregnancies Complicated by Preeclampsia . *Hypertension in Pregnancy*.Volume 23, Issue 1, 2004,pages 101-111.

Levine RJ et al.Circulating angiogenic factors and the risk of preeclampsia.*N Engl J Med* 2004;350: 672-83.

Levine, R.J. et al. (2005) Urinary placental growth factor and risk for preeclampsia. *JAMA* 293:77.

Linda J. Heffner, Danny J. Schust: psychosocial stress and chronic hypertension with preeclampsia. *Am J*. 2005 Sep; 106(3):444-5.

Lindheimer MD,Taler SJ,Cunningham FG.Hypertension in pregnancy.*J of Americ Soc of Hypertension*.2008;2:484-94.

Lu D, Yang X, Wu Y, Wang H, Huang H, Dong M.: Serum adiponectin, leptin and soluble leptin receptor in preeclampsia. *Int J Gynaecol Obstet* 2006;95:121-6.

Luealon, P., Phupong, V., 2010. *Risk Factors of Preeclampsia in Thai Women*. J Med Assoc Thai. Department of Obstetrics and Gynecology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Llurba E, Grataco E, Galla MP, Cabero L, Dominguez C. 2004. *A comprehensive study of oxidative stress and antioxidant status in preeclampsia and normal pregnancy*. Free Radical Biology & Medicine; 37(4): 557 – 570.

Marc.Montminy: "How Obesity Increases The Risk For Diabetes." Salk Institute.ScienceDaily. ScienceDaily, 22 June 2009.

Marshall DL, MD, and Jason GU, MD, PhD.: Explaining and predicting preeclampsia. N Engl J Med. 2006; 355:105610.

Matsuda M, Shimomura I, Sata M, Arita Y, Nishida M, Maeda N, Kumada M, Okamoto Y, Nagaretani H, Nishizawa H, Kishida K, Komuro R, Ouchi N, Kihara S, Nagai R, Funahashi T, Matsuzawa Y 2002 : Role of adiponectin in preventing vascular stenosis. The missing link of adipo-vascular axis. J Biol Chem 277:37487–37491

Matsuzawa Y.et al Adiponectin and metabolic syndrome. Arteriosclerosis, thrombosis, and vascular biology. 2004, 24; 29-33

Meis P, Goldenberg R and Mercer B (1998): The preterm prediction study: Risk factors for indicated preterm birth. Am J Obst Gynecol, 179: 562-7.

Michael M, et al. Physiological, Pharmacological and Nutritional regulation of circulating adiponectin concentration in humans. Mary ann Liebert, Vol.6, number 2, 2008.

Michael M. Swarbrick, Ph.D. and Peter J. Havel, D.V.M., Ph.D.corresponding author: Physiological, Pharmacological, and Nutritional Regulation of Circulating Adiponectin Concentrations in Humans. Metab Syndr Relat Disord. 2008 Jun; 6(2): 87–102.

Mijal RS,Holzman CB,Rana S,Karumanchi A,Wang J,Sikorskii A.Mild pregnancy levels of angiogenic markers in relation with maternalcharacteristic.Am J Obstet Gynecol.2012;204(3):244,el-244.12.

Mikat B,A. Gellhaus, N.Wagner, C.Birdir, R.Kimmig, and A.Koninger": Early Detection of Maternal Risk for Preeclampsia. International Scholarly Research Network ISRN obstetrics and Gynecology Volume 2012, Article ID 172808,pages.1-7.

Milne, G.L., Musiek, E.S., Morrow, J.D. 2005. F2-Isoprostanes as markers of oxidative stress in vivo: An overview. *Biomarkers*, 10 (Suppl. 1):S10-S23. media LCC : New York. Hal 178-182.

Mitayani. Asuhan keperawatan maternitas. Salemba medika.Jakarta.2009.

Montuschi, P., Barnes, P.J., Roberts, L.J. 2004. Isoprostanes: markers and mediators of oxidative stress. *The FASEB Journal*, 18:1792-1800.

Muniyappa R,Sower JR : Role of insulin resistance in endothelial dysfunction.Rev.Endocr metab disorder.2013;14: 5-12.

Myatt L,Clifton RG,Roberts JM,et al.The utility uterine artery Doppler velocimetry in prediction of preeclampsia in low risk population.*Obstet-gynecol* 120 (4):815,2012(b).

Myles W, Kettyle E, Sandler L. et al. Obesity and preeclampsia: the potential role of inflammation. *Obstet Gynecol* 2001; 98:757-62.

Nagamatsu, T. *et al.* (2004) Cytotrophoblasts up-regulate soluble fms-like tyrosine kinase-1 expression under reduced oxygen: an implication for the placental vascular development and the pathophysiology of preeclampsia. *Endocrinology*.145: 4835.

Nakatsukasa H, Masuyama H, Takamoto N, Hiramatsu Y. : Circulating leptin and angiogenic factors in preeclampsia patients. *Endocr J* 2008;55:565–73.

Naruse K, Yamasaki M, Umekage H, Sado T, Sakamoto Y, Morikawa H. Peripheral blood concentrations of adiponectin, an adipocyte-specific plasma protein, in normal pregnancy and preeclampsia. *J Reprod Immunol* 2005; 65:65–75.

Neslihan Bukan,Omer Kandemir,Tuncay Nas,Ozlem Gulbahar,Adem Unal& Banu Cayci : Maternal cardiac risks in pre-eclamptic patients. *The Journal of Maternal- Fetal and Neonatal Medicine*, 2012; 25(7): 912–914.

Nicolaides K.H : “Turning the pyramid of prenatal care,” *Fetal Diagnosis and Therapy*, vol. 29, no. 3, pp. 183–196, 2011.

Nien JK, Mazaki-Tovi S, Romero R, Erez O, Kusanovic JP,Gotsch F, *et al.* Adiponectin in severe preeclampsia. *J Perinat Med* 2007;35:503–12.

Niki E, Yoshida Y, Saito Y, Noguchi N. 2005. Lipid peroxidation: Mechanisms, inhibition, and biological effects. *Biochemical and Biophysical Research Communication*; 338: 668–676.

Niki E. 2009. *Lipid peroxidation: Physiological levels and dual biological effects.* *Free Radical Biology & Medicine*; 47: 469–484.

Norwitz ER, Robinson J, Repke J : Prevention of preeclampsia: is it possible.Clin obstet & Gynecol.1999.P.436-54.

O'Brien TE, Ray JG, Chan WS.maternal body mass index and the risk of preeclampsia: a systematic overview.Epidemiology. 2003 May;14(3):368-74.

Ogge G,Chaiworapongsa T,Romero R,Youssef H,Kusanovic JP,Yeo L.Placental lesion associated with maternal underperfusion are more frequent in early onset than in late onset preeclampsia.*J Perinat Med*.2011;39(6):641-52.

Opitasari C, Andayasari L. Parity, education level and risk for (pre-) eclampsia in selected hospitals in Jakarta. *Health Science Indones.* 2014;5:35-9.

Ouyang, Yanqiong Reduced plasma adiponectin and elevated leptin in pre-eclampsia, *International Journal of Gynecology and Obstetrics* 2007; 98–114.

Ouyang YQ et al. Plasma sFLt-1 to PIGF ratio is correlated with . inflammatory but not with oxidative stress in Chinese preeclamptic women. *Archives of Gynecology and Obstetrics*, 2008, 280:91–97.

Pasaoglu H, Bulduk G, Agus E, Pasaoglu A, Onalan G. Nitric oxide, lipid peroxides, and uric acid levels in pre-eclampsia and eclampsia. *Tohoku Journal of Experimental Medicine*, 2004; 202; 87- 92.

Patil SB, Kodliwadmath MV, Kodliwadmath SM. Correlation between lipid peroxidation and Nonenzymatic antioxidants in pregnancy induced hypertension. *Indian Journal of Clinical Biochemistry*, 2008; 23(1);45-48.

Patrignani, P., Tacconelli, S. 2005. Isoprostanes and Other Markers of Peroxidation in Atherosclerosis. *Biomarkers*, 10 (Suppl 1):S24-S29.

Pajvani, UB Structure-function studies of the adipocyte-secreted hormone acrp30 I adiponectin. *The American Society for Biochemistry and Molecular Biology*. December 20, 2002 ; Vol.278 No'1 1

Pijnenborg R, Vercruyse L, Hanssens M, Van Assche FA. Trophoblast invasion in preeclampsia and other pregnancy disorders. In *Pre-eclampsia Etiology and clinical Practice*, New York : Cambridge University Press 2007 : 1-20.

Poon.LC; K.H Nicolaides : Early Prediction of Preeclampsia. *Obstetric and Gynecology International*.Vol 2014, Article ID 297397,p.1-7.

Prawirohardjo.S : Hipertensi dalam kehamilan Ilmu Kebidanan.Edisi IV.Yayasan Bina Pustaka Sarwono Prawirohardjo.2002

Quedarusman H, Wantania J, et al. Hubungan Indeks Massa Tubuh Ibu dan Peningkatan Berat Badan Saat Kehamilan dengan Preeklampsia. *Jurnal e-Biomedik (eBM)*. 2013;1.

Ramsay JE,et al. Paradoxical elevation in adiponectin concentrations in women with preeclampsia. *American Heart Association*. September 2003 ; 42:891-894.

Roberts CL, Ford JB, Algert CS, et al. *BMJ Open* 2011. Published online 2011 May 24.

Robert.JM : Endothelial dysfunction in preeclampsia. *Reprod Endocrinol*.1998;16: 5-15.

Robert.JM; Hilary Gammil : Insulin Resistance in Preeclampsia. *Hypertension*.March.2006; 47; 341-342.

Roberts JM, Hubel CA. The two stages model of preeclampsia: Variations on theme. *placenta trophoblast research*. 2009;23:32-7.

Roberts J, Lain K. Recent insights into the pathogenesis of pre-eclampsia. *Placenta* 2002;23:359-72.

Roberge S, Y. Giguere, P. Villa et al., "Early administration of low-dose aspirin for the prevention of severe and mild preeclampsia: a systematic review and meta-analysis," *American Journal of Perinatology*, vol. 29, no. 7, pp. 551-556, 2012

Robinson S & Phil Pemberton & Ian Laing & Luciano G. Nardo : Low grade inflammation, as evidenced by basal high sensitivity CRP, is not correlated to outcome measures in IVF, *Assist Reprod Genet*, 2008 ; 25:383-388.

Robson J, Dostal I, Mathur R, Hippisley-Cox J, Coupland. Predicting cardiovascular risk in England and Wales. 2008.

Robins, Cotran, et al. (2006), *Blood Vessel. Pathologic basis of disease* 7th edition Elsevier: 511- 554.

Romero R, Nien JK, Espinoza J, Todem D, Fu W, Chung H, Kusanovic JP, Gotsch F, Erez O, Mazxaki-Tovi S, Gomez R, Edwin S, Chaiworapongsa T, Levine RJ and Karumanchi A (2008): A longitudinal study of angiogenic (placental growth factor) and antiangiogenic (soluble endoglin and soluble vascular endothelial growth factor receptor-1) factors in normal pregnancy and patients destined to develop preeclampsia and deliver a small for gestational age neonate. *J Maternal-Fetal & Neonatal Med* 21;1:9-23.

Rosario D, et al. Plasma adiponectin concentration in early pregnancy and subsequent risk of hypertensive disorder. American College of Obstetricians and Gynecologists - Augustus 2005, Vol 106, No 2.

Rolv Skjaerven: Norway British Medical Journal edisi online 2 November, 2007.
Rukiyah, Yulianti (ed.). *Asuhan kebidanan* ed IV. Trans info media. Jakarta. 2010 hal: 172- 190.

Rozikhan R. faktor-faktor risiko terjadinya preeklampsia berat di rumah sakit dr. H. Soewondo kendal. tesis program magister epidemiologi universitas diponegoro semarang. 2007.

Rukiyah, Yulianti. 2010. *Asuhan Kebidanan IV (Patologi Kebidanan)*, Jakarta : TIM

Saifuddin AB. 2009. *Buku Panduan Praktis Pelayanan Kesehatan Maternal Neonatal*. Jakarta : Yayasan Bina Pustaka Sarwono Prawirohardjo.

Salimi S, Farzaneh, Farajian-Mashhadi, Anoosh Naghavi, Mojgan Mokhtari, Mahnaz

Shahrakipour, Mohsen Saravani, and Minoo Yaghmaei : Different serum of leptin between Early Onset and Late Onset Preeclampsia. Hindawi Publishing Corporation Disease Markers Volume 2014, Article ID 628476, P.1-7.

Sampson, M.J., Gopaul, N., Davies, I.R., Hughes, D.A., Carrier, M.J. 2002. Plasma F2 Isoprostanes : Direct Evidence of Increased Free Radical Damage During Acute Hyperglycemia in Type 2 Diabetes. *Diabetes care*, 25(3):537-541.

Sari Mayang, 2014 :Faktor – faktor yang berhubungan dengan kejadian pre eklamsi pada ibu hamil di poli kebidanan RS Kesdam Banda Aceh, Banda Aceh : Stmikubudiyah.

Sattar, N. Bendomir, A. Lipoprotein Subfraction Concentrations in Preeclampsia : Pathogenic Parallels to Atherosclerosis. *AJOG* 1997;89:403-8.

Sattar N, Bedomir A, Berry C, Shepherd J, Greer IA, Packard CJ. Lipoprotein subfraction concentration in preeclampsia: pathogenic parallels to atherosclerosis. *Obstet Gynecol*. 1997;83:p.403-408.

Scholten RR; Maria T.E. Hopman; Fred C.G.J. Sweep; Maureen J. van de Vlugt; Arie P. van Dijk; Wim J.G. Oyen; Fred K. Lotgering; Marc E.A. Spaanderman. Co-occurrence of cardiovascular and prothrombotic risk factors in women with a history of preeclampsia. *Obstet Gynecol*. 2013; 121(1): 97-105.

Shore VH, Wang TH, Wang CL, Torry RJ, Claude MR and Torry DS. Vascular endothelial growth factor ,placental growth factor and their receptor in isolated human trophoblast. *placenta* 1997.18:657-665.

Sibai, B.M. Diagnosis, Prevention, and Management of Eclampsia. *American Journal Obstetrics Gynaecology*. 2005 Vol: 105:405-410.

Sibai B. Hypertensive Disorders in Pregnancy. In: SG SG, Niebyl J, Simpson J, Landon M, HL HG, Jauniaux E, editors. *Obstetrics Normal and Problem Pregnancies*. Philadelphia: Elsevier; 2012. p. 780-284.

Sibiude J, Guibourdenche J, Dionne MD, Le Ray C, Anselem O, Serrau R Goffinet F, Tsatsaris. Placental growth factor for the prediction of adverse outcomes in patients with suspected preeclampsia or intrauterine growth restriction. *PLoS One*. 2012;7(11):e50208. doi:10.1371/journal.pone.0050208. Epub 2012 Nov 28.

Sitti Nur Djannah, Ika Sukma Arianti. Gambaran epidemiologi kejadian preeklamsi eklamsi di rsu pku muhammadiyah yogyakarta tahun 2007–2009

Sofoewan.H.M: Preeklamsi-Eklamsi di beberapa Rumah Sakit di Indonesia, patogenesis dan kemungkinan pencegahannya. Pidato Pengukuhan Jabatan Guru Besar pada Fakultas Kedokteran Universitas Gadjah Mada Diucapkan di depan Rapat

Terbuka Majelis Guru Besar Universitas Gadjah Mada pada tanggal 3 Februari 2003 di Yogyakarta.

Steegers EA, von Dadelszen P, Duvekot JJ, Pijnenborg R. Pre-eclampsia. Lancet. 2010 Aug 21; 376(9741):631-44. Epub 2010 Jul 2.

Stern.M : Diabetes and Cardiovascular Disease The “common soil” Hypothesis. Diabetes. 1995;44. P.369-74.

Sulistiyowati S, Abadi A. Low Class Ib (HLA-G/Qa-2) MHC Protein Expression against Hsp-70 and VCAM-1 Profile on Preeclampsia. Indones J Obstet Gynecol 2010;34(1):103-7.

Sutopo H, Surya IGP. Characteristics of Patient with Hypertension in Pregnancy at Sanglah Hospital. Indonesia Journal Obstetrics and Gynecology. 2011;35(3):97-9.

Sujiyatini, dkk. 2009: Asuhan patologikebidanan Yogyakarta: Nuh Medika. Sulaiman,sastrawinata. 2005. Jakarta : PT. Bina sarwono prawirohardjo.

Suwaki Naoko. Hypoadiponectinemia and circulating angiogenic factors in overweight patients complicated with pre-eclampsia , American Journal of Obstetrics and Gynecology, 2006; 92-195.

Takacs P, Kauma SW, Sholley MM, Walsh SW, Dinsmoor MJ, Green K. 2001.Increased circulating lipid peroxides in severe preeclampsia activate NF- κ B and upregulate ICAM-1 in vascular endothelial cells. The FASEB Journal: 15: 279-281.

Takamura Y, Osuga Y, Koga K, Tajima T, Hirota Y, Hirata T, et al. Selective increase in high molecular weight adiponectin concentration in serum of women with preeclampsia. J Reprod Immunol 2007;73:60–65.

Tan K H KK, H YGS. Epidemiology of pre-eclampsia and eclampsia at the KK Women’s and Children’s Hospital, Singapore. Singapore Med J. 2006;41(1):48-53.

Tanto, S.S. 2008. Pengaruh Konsumsi Ekstrak Buah Mengkudu (Morinda Citrifolia Fructus) pada Penanganan Preeklamsi Ringan secara Konvensional Melalui Penilaian Status Nitrik Oksida, F2-Isoprostan, Tekanan Darah, dan Daya Pencegahan Kejadian Preeklamsi Berat dan Eklamsi. (*disertasi*). Bandung : Universitas Padjadjaran,

Taylor.RN,Grimwood J,Taylor RS,McMaster MT,Fisher SJ,North RA. Longitudinal serum concentrations of placental growth factor:evidence for abnormal placental angiogenesis in pathologic pregnancies.Am J Obstet Gynecol 2003;188: 177-82.

Thadhani R, Mutter WP, Wolf M, Levine RJ, Taylor RN, Sukhatme VP, Ecker J, Karumanchi SA (2004): First trimester placental growth factor and soluble fms-like tyrosine kinase 1 and risk for preeclampsia. J Clin Endocrinol Metab. 89(2):770-775.

Trijatmo : Peeklamsi dan Eklamsi.Ilmu Kebidanan.Edisi III.Bina Pustaka SarwonoPrawirohardjo.2005.p.281-301.

Tug N, Celik H, Cikim G, Ozcelik O, Ayar A. 2003. The correlation between plasma homocysteine and malondialdehyde levels in preeclampsia. Neuroendocrinology Letters; 24: 445-448.

Vatten LJ, Eskild A, Nilsen TI, Jeansson S, Jenum PA, Staff AC (2007): Changes in circulating level of angiogenic factors from the first to second trimester as predictors of preeclampsia. Am J Obstet Gynecol. 196(3):239. e1-6.

WHO Buku saku pelayanan kesehatan ibu di fasilitas kesehatan dasar dan rujukan. Jakarta2013.

Wibisono. Kematian perinatal pada *preeklampsia-eklampsia*. Fak.Ked. Undip Semarang, 1997; 6-12.

Wibowo N. Peran Radikal bebas pada preeldanpsia. Seminar Konsep Mutakhir preeklamsia. Jakarta.2001.

Wikipedia.com.Hytest.2009.Finland : Adiponectin.

Winknjosastro, H. Preeklampsia Berat. Ilmu kandungan edisi ketiga. Yayasan Bina Pustaka Sarwono Prawirohardjo, Jakarta 1999; 281-308.

Winkjosastro, Hanifa. 2007. *Ilmu Kebidanan*. Jakarta : Yayasan Bina Pustaka Sarwono Prawirohardjo

Wijaya, A. Adiponectin: A promising biomarker for MetS, & risk factors for T2D, hypertension, CHD, CVD, NASH & cancer. Prodia. 2006.

Wright D, Syngelaki A, Akolekar R, Poon LC, Nicolaides KH. Competing risks model in screening for preeclampsia by maternal characteristics and medical history. Am J Obstet Gynecol. 2015;62:1-10.

Zhonghua F, Chan Ki, Study of adiponectin expression in placenta and itscorrelation with preeclampsia, Pubmed. U.S National Library of Medicine National Institutes of Health. February 2008 ; 43(2):90