

DAFTAR PUSTAKA

- Abler LL, Keil KP, Mehta V, Joshi PS, Schmitz CT, Vezina CM. 2011. A high resolution molecular atlas of the fetal mouse lower urogenital tract. *Dev Dyn* 240(10):2364-77
- Adamovic T, Nordenskjöld A. 2012. The CAG repeat polymorphism in the androgen receptor gene modifies the risk for hypospadias in caucasians. *BMC Medical Genetics* 13:1-4.
- Adamovic T, Thai HT, Liedén A, nordenskjöld A. 2013. Association of a tagging single nucleotide polymorphism in androgen receptor gene region with susceptibility to several hypospadias in a Caucasian population. *Sex Dev* 7(4):173-9.
- Alberts B, Bray D, Hopkin B, Johnson A. 2010. *Essential Cell Biology* 3 rd ed. Garland Science. New york: 272-81
- Andhika A, Gerhard RS, Wahyudi I. 2015. *Current practice in Paediatric hypospadias surgery in Indonesia; Survey in Indonesian Urologist*. Jakarta.
- Angaji SA. 2009. Single nucleotide polymorphism genotyping its application on mapping and marker-assisted plant breeding. *African J Biotechnolog* 8(6):908-14.
- Aschim EL, Nordenskjold A, Giwercman A, Lundin KB, Ruhayel Y, Haugen TB. et al. 2004. Linkage between cryptorchidism, hypospadias, and GGN repeat length in the androgen receptor gene. *J Clin Endocrinol Metab* 89:5105-5109.
- Ban S, Sata F, Kurahashi N, Kasai S, Moriya K, Kakizaki, et al. 2008. Genetics polymorphisms of ESR1 and ESR2 that may influence estrogen activity and the risk of hypospadias. *Human Reproduction* 23(6):1466-71.
- Baskin LS, Himes K, Colborn T. 2001. Hypospadias and endocrine disruption: is there a connection?. *Environmental Health Perspectives* 109:1175-83.
- Baskin LS, 2008. Can we prevent hypospadias? *Fertil Steril* 89(2 Suppl): e39.
- Bay K, Asklund C, Skakkebaek NE, Andersson AM. 2006. Testicular dysgenesis syndrome: possible role of endocrine disruption. *Best Pract Res ClinEndocrinolMetab* 20(1):77-90.

- Bayu F, Doddy E, Alvarino, Yevri Z, Etriyel M.2014. Distribusi frekuensi dan tatalaksana kasus hipospadia di RSUP dr M Djamil Padang tahun 2013. Sub divisi urologi. Bagian Bedah FK Unand
- Beleza-Miereles A, Barbaro M, Wedell A, Töhönen V, Nordenskjöld A. 2007. Studies of a co-chaperone of the androgen receptor, FKBP52, as candidate for hypospadias. *Reprod Biol Endocrinol* 5:1-7.
- Beleza-Miereles A, Lunberg F, Lagerstedt K, Zhou X, Omrani D, Frisen L et al. 2007. FGF2, FGF8, FGF10 and BMP7 as candidate genes for hypospadias. *European JHumGenet*15:405-10.
- Beleza-Miereles A, Kockum I, Lundberg F, Söderhäll C, Nordenskjöld A. 2007. Risk factors for hypospadias in the estrogen receptor 2 gene. *JClinEndocrinol Metab* 92(9):3712-18.
- Beleza-Miereles A, Töhönen V, Söderhäll C, Schwentner C, Radmayr C, Kockum I et al, 2008. Activating transcription factor 3: a hormone responsive gene in the etiology of hypospadias. *European JEndocrinol*158:729-39.
- Bieche I, Parfait B, Tozlu S, Lidereau R, Vidaud M. 2001. Quantitation of androgen receptor gene expression in sporadic breast tumors by real time RT-PCR: evidence that MYC is an AR-regulated gene. *Carcinogenesis* vol 22 no 9, 1521-26.
- Blaschko SD, Cunha GR, Baskin LS. 2012. Molecular mechanisms of external genitalia development. *Differentiation* 84(3):261-8.
- Borhani N, Novin MG, Manochehri M, Rouzrokh M, Kazemi B, Koochaki A et al, 2014. New single nucleotide variation in the promoter region of androgen receptor (AR) gene in hypospadiac patients. *Iran J Reprod Med* 12(3) : 217-220.
- Botta S, Cunha GR, Baskin S. 2014. Do endocrine disruptors cause hypospadias?. *TranslAndrolUrol* 3(4):330-339.
- Carmichael SL, Shaw GM, Nelson V, Selvin S, Torfs CP, Curry CJ. 2003. Hypospadias in California: trends and descriptive epidemiology. *Epidemiol* 14(6):701-6.
- Carmichael SL, Shaw GM, Laurent C, Olney RS, Lammer EJ. 2007. Maternal reproductive and demographic characteristics as risk factors for hypospadias. *Paediatr Perinat Epidemiol.* 21:210–218
- Carmichael SL, Shaw GM, Lammer EJ. 2012. Environmental and genetic contributors to hypospadias: a review of the epidemiologic evidence. *Birth Defects Res A Clin Mol Teratol* 94(7):499-510.

- Carmichael SL, Ma C, Choudhry S, Lammer EJ, Witte JS, Shaw GM. 2013. Hypospadias and genes related to genital tubercle and early urethral development. *J. Urol* 190(5):1884-1892.
- Carmichael SL, Yang W, Roberts EM, Kegley E, Wolff C, Guo L, et al. 2013. Hypospadias and residential proximity to pesticide application. *Pediatrics* 132: e1216-e1226.
- Carmichael SL, Shaw GM, Laurent C, Croughan MS, Olney MS, Lammer EJ. 2005. Maternal progestin intake and risk of hypospadias. *Arch Pediatr Adolesc Med* 159(10): 957-962
- Carmichael SL, Witte JS, Ma C, Lammer J, Shaw M. 2014. Hypospadias and variants in genes related to sex hormone biosynthesis and metabolism. *Androl* 2:130-7.
- Chan SC, Dehm SM. 2014. Constitutive activity of the androgen receptor. *Adv Pharmacol* 70: 327-366.
- Cunningham RL, Lumia AR, McGinnis MY. 2012. Androgen receptors, sex behavior, and aggression. *Neuroendocrinology* 96(2):131-140
- Dart DA, Waxman J, Aboagye EO, Bevan CL. 2013. Visualising androgen receptor activity in male and female mice. *Plos one* 8(8):1-16
- Drysdale CM, McGraw DW, Stack CB et al. 2000. Complex promoter and coding region *b₂*-adrenergic receptor haplotypes alter receptor expression and predict in vivo responsiveness. *PNAS* 97:10483-8.
- Dussor GO, Price TJ, Flores CM. 2003. Activating transcription factor 3 mRNA is upregulated in primary cultures of trigeminal ganglion neurons. *Mol Brain Research* 118:156-9.
- Eisermann K, Wang D, Jing Y, Pascal LE, Wang Z. 2013. Androgen receptor mutation, rearrangement, polymorphism. *Transl Androl Urol* 2(3):137-147
- Fariz M, Rodjani A, Wahyudi I. 2010. Komplikasi pasca operasi hipospadia dan penatalaksanaannya. Departemen Urologi, RSUPN Cipto Mangunkusumo, FK UI
- Fariz M. 2009. Faktor-faktor risiko timbulnya distula uretrokutan pasca operasi hipospadia satu tahap. Departemen Urologi, RSUPN Cipto Mangunkusumo, FK UI
- Fisch H, Golden RJ, Gary L, Libersen, Hyun GS, Madsen P. 2001. Maternal Age as a risk factor for hypospadias. *J Urol* 165:934-936.

Fisch H, Lambert Sm, Hensel TW, Hyun G. 2009. Hypospadias rates in New York State are not increasing. *J Urol* 181(5):2291-4.

Frisen L. 2002. Genetic Studies of Hypospadias. Department of Molecular Medicine Karolinska Institute. Stockholm.

Genecard. ATF3. <http://www.genecards.org/cgi-bin/carddisp.pl?gene=ATF3> diakses Jan 10, 2015

Genecard. AR. <http://www.genecards.org/cgi-bin/carddisp.pl?gene=AR> diakses Feb 22, 2015

Ghirri P, Scaramuzza RT, Bertelloni S, Pardi D, Celandroni A, Cocchi G et al. 2009. Prevalence of hypospadias in Italy according to severity, gestational age and birthweight: an epidemiological study. *Italian J Pediat* 35:1-7.

Gurbuz C, Demir S, Zemheri E, Canat L, Kilic M, Caskurlu T. 2010. Is activating transcription factor 3 up regulated in patients with hypospadias?. *Korean J Urol* 51:561-4.

Hackl C et al. 2010. Activating transcription factor-3 (ATF3) functions as a tumor suppressor in colon cancer and is up-regulated upon heat-shock protein 90 (Hsp90) inhibition. *BMC cancer*, 10:668

Hai T, Jalgaonkar S, Wolford CC, Yin X. 2011. Immunohistochemical detection of activating transcription factor 3, a hub of the cellular adaptive-response network. *Methods Enzymol* 490:175-94.

Hoetzenecker W, Echtenacher B, Guenova E, Hoetzenecker K, Woelbing F, Brug J et al. 2013. ROS-Induced ATF3 causes susceptibility to secondary infections during sepsis-associated immunosuppression. *Nat Med* 18(1):128-34.

Jamsari. 2007. Bioteknologi pemula, prinsip dasar dan aplikasi analisis molekuler. UNRI Press. Pekanbaru.

Jamsari. 2013. Rekayasa genetika, untuk analisis genom dan produksi organisme transgenik. UR Press. Pekanbaru

Kalfa N, Liu B, Klein O, Wang MH, Cao M, Baskin LS. 2008. Genomic variants of ATF3 in patients with hypospadias. *J Urol* 180:2183-8.

Karabulut R, Turkyilmaz Z, Sonmez K, Kumas G, Ergun SG, Ergun MA et al. 2013. Twenty four genes are upregulated in patients with hypospadias. *BJMG* 16(2); 39-44.

Kim EY, Shin HY, Kim JY, Kim DG, Choi YM, Kwon HK et al. 2010. ATF3 plays a key role in Kdo₂-lipid a-induced TLR4-dependent gene expression via NF- κ B activation. *PLoS ONE* 5:1-9.

- King MW. Control of gene expression. [homepage on the Internet]. [cited 2017 May 25]. Available from: <http://themedicalbiochemistrypage.org/gene-regulation.php>
- Kojima Y, Kohri K, Hayashi Y. 2010. Genetic pathway of external genitalia formation and molecular etiology of hypospadias. *JPediatUrol* 6:346-54.
- Kon M, Suzuki E, Dung VC, Hasegawa Y, Mitsui T, Muroya K et al. 2015. Molecular basis of non syndromic hypospadias: systemic mutation, screening and genome-wide copy-number analysis of 62 patients. *Human reproduction* 0(0): 1-8.
- Limatahu N, Oley MH, Monoarfa A. 2012. Angka kejadian hipospadia di RSUP Prof. Dr. R. D. Kandou Manado periode januari 2009-oktober 2012. *Bagian Bedah Universitas Sam Ratulangi Manado*.
- Liu B, Wang Z, Lin G, Agras K, Ebbers M, Willingham E et al. 2005. Activating transcription factor 3 is up-regulated in patients with hypospadias. *Pediatr Res* 58:1280-3.
- Lu Dan, Chen J, Hal T. 2007. The Regulation of ATF3 gene Expression by mitogen-activated protein kinases. *Biochemical Society*: 559-67
- Manson JM, Carr MC. 2003. Molecular epidemiology of hypospadias: review of genetic and enviromental risk factors. *Birth Defects Research (Part A)* 67: 825-36.
- McLaughlin DT, Donahoe PK. 2004. Sex determinan and differentiatin. *N Engl J Med* 350:367-78.
- Miyagawa S, Matsumaru D, Murashima A, Omori A, Satoh Y, Haraguchi R et al. 2011. The Role of sonic hedgehog-gli2 pathway in the masculinization of external genetalia. *Endocrinol* 152(7):2894-903.
- Muhammadzadeh A, Farhat A, Esmaili H, Shiranzaei S. 2011. Prevalence and risk factors of hypospadias in a private hospital in northeast iran. *Iranian journal of pediatrics* 21(4):497-501.
- Nassar N, Bower C, Baker A. 2007. Increasing prevalence of hypospadias in Western Australia, 1980-2000. *Arch Dis Child* 92(7): 580-4.
- Palmer JR, Wise LA, Robboy SJ, Titus-Ernstoff, Noller KL, Herbst Al et al. 2005. Hypospadias in sons of woman exposed to diethylstilbestrol in uretro. *Epidemiology* 16(4):583-6.

- Pande Made WT, Yevri Z, Alvarino. 2015. The outcomes of urethroplasty for hypospadias repair in M Djamil Hospital. *Journal of Advances in Medical and Pharmaceutical Sciences*.4:1-5.
- Pichler R et al. 2013. Quantitative measurement of the androgen receptor in prepuces of boys with and without hypospadias. *BJU int* Jul;112(2): 265-70
- Pierik FH, Burdorf A, Nijman JMR, Keizer-Schrama SM, Juttman RE, Weber RFA. 2002. A high hypospadias rate in the Netherlands. *Hum Reprod*17(4):1112-15.
- Pirmohamed and Park. 2001. Genetic susceptibility to adverse drug reactions. *Trends PharmacolSci* 22:298-305.
- Porter MP, Faizan MK, Grady RW, Mueller BA. 2004. Hypospadias in Washington State: maternal risk factors and prevalence trends. *Pediatrics* 115(4):e495-9.
- Qiao L, Tasian GE, Zhang H, Cao M, Ferretti M, Cunha GR et al. 2012. Androgen receptor is overexpressed in boys with severe hypospadias, and ZEB1 regulates androgen receptor expression in human foreskin cells. *Pediatr Res*. 71(4 Pt 1):393-8.
- Qin XY, Kojima Y, Mizuno K, Ueoka K, Muroya K, Miyado M et al. 2012. Identification of novel low-dose bisphenol A targets in human foreskin fibroblast cells derived from hypospadias patients. *Plos ONE* 7:1-9.
- Qin XY, Kojima Y, Mizuno K, Ueoka K, Massart F, Spinelli C.2012. Association of variants in genes involved in environment chemical metabolism and risk of cryptorchidism and hypospadias. *Journal of human Genetic* 00: 1-8
- Radpour R, Rezaee M, Tavasolu A, Solati S, Saleki A. 2007. Association of long polyglycine tracts (GGN repeats) in exon 1 of the androgen receptor gene with cryptorchidism and penile hypospadias in Iranian patients. *J Androl* ; 28: 164-169
- Rignell-Hydbom A, Lindh CH, Dillner J, Jönsson Bo AG, Rylander L. 2012. A nested case control study of intrauterine exposure to persistent organochlorine pollutants and the risk of hypospadias. *Plos ONE* 7:1-5.
- Risa A, Zulfiqar Y, Etriyel MYH, Erkadius, Rodjani A, Wahyudi I. 2016. Uroflowmetri parameter evaluation of post urethroplasty hypospadias patients in Padang. *Journal of advances in Medical and Pharmaceutical Sciences* 5(4): 1-6
- Risch N, Merikangas K. 1996. The future of genetic studies of complex human diseases. *Science* 273:1516-7.
- Rocheleau CM. 2009. Pesticide exposure and risk of hypospadias: assesment and the adequacy of exposure measurements. Dissertation. University of Iowa.

- Rynes J, Donohoe CD, Frommolt P, Brodesser S, Jindra M, Uhlirova M. 2012. Activating transcription factor 3 immune and metabolic homeostasis. *Mol Cell Biol* 32:3949-62.
- Samtani R, Bajpai M, Vashisht K, P. K. Ghosh, K. N. Saraswathy. 2011. Hypospadias Risk and Polymorphism in SRD5A2 and CYP17 Genes: Case-Control Study Among Indian Children. In *Journal of Urology*, Vol. 185, 2334-2339
- Sastroasmoro S, Madiyono B, Moeslichan S, Budiman I, Purwanto, SH. 2008. Perkiraan besar sampel. Dalam Sastroasmoro S & Ismael S. Edisi 3, *Dasar-Dasar Metodologi Penelitian Klinis*. Sagung Seto Jakarta.
- Sastrasupena RH, Utomo T. Penggunaan kontrasepsi oral saat kehamilan dan resiko terjadinya hipospadia pada anak di SMF urologi RSUP Dr Sardjito. Jakarta, 2013
- Sathyanarayana S, Swan SH, Farin FM, Wilkerson HW, Bamshad M, Grady R et al. 2012. A pilot study of the association between genetic polymorphism involved in estrogen signaling and infant male genital phenotypes. *Asian J Androl* 14:766-72.
- Schnack TH, Poulsen G, Myrup C, Wohlfachrt J, Melbye M. 2010. Familial coaggregation of chryptorchidism an hypospadia. *Epidemiology* 21(1):109-113
- Schnack TH, Poulsen G, Myrup C, Wohlfachrt J, Melbye M. 2009. Familial coaggregation of chryptorchidism an hypospadia, and testicular germ cell cancer: a nationwide cohort study. *J Natl Cancer Inst* 102:1-6.
- Sharma N, Bajpal M, Panda SS, Singh A, Pandey RK, Ali A. 2013. Family based genetic study in proximal penile hypospadias with undescended testis. *Journal of Progres in Paediatric Urology* 16(2):42-45.
- Snodgrass WT, 2012. Hypospadias. In: Wein AJ, Kavoussi LR, Novick AC, Partin AW, Peter CA, eds. *Campbell-Walsh Urology*, 10th ed. Philadelphia: Elsevier Saunders, 3503-9.
- Stein R. 2012. Hypospadias. *European Urology Supplements* 11:33-45.
- Steinhardt GF. 2004. Endocrine disruption and hypospadias. *AdvExp Med Biol* 545: 203-15.
- Stoll C, Alembik Y, Roth MP, Dott B. 1990. Genetic and enviromental factors in hypospadias. *J Med Genet* 27:559-63.
- Storgaard L, Bonde JP, Olsen J. 2006. Male reproductive disorders in human and prenatal indicators of estrogen exposure, a review of published epidemiological studies. *ReprodToxicol* 21(1):4-15.

- Suzuki K, Haraguchi R, Ogata T, Barbieri O, Alegria O, Vieux-Rochas M et al. 2008. Abnormal urethra formation in mouse models of split-hand/split-foot malformation type 1 and type 4. *European J Genet* 16:36-44
- Takahashi T, Shimotakahara A, Miyahara K, Lane GJ, Yamataka A. 2013. Activating transcription factor 3 is not up-regulated in hypospadias patients in Japan. *African J Paediatric Surgery* 10:371-3.
- Takahashi T et al. 2013. Activating transcription factor 3 is not up-regulated in hypospadias patients in Japan. In *African Journal of Paediatric Surgery*. Vol 10: 371-373
- Thai HT, Kalbasi M, Lagerstedt K, Frisén L, Nordenskjöld A. 2005. The valine of the V89L polymorphism in the 5-alpha-reductase gene confers a reduced risk for hypospadias. *J Clin Endocrinol Metab* 90:6695-8.
- Traish AM, Morgentaler A. 2009. Epidermal Growth Factor receptor expression escapes androgen regulation in prostate cancer : a potential molecular switch for tumour growth. *British Journal of Cancer*: 1949-56.
- Trinh Thi. 2009. Hypospadias: Gene Mapping and Candidate gene studies. Department of Women's and Children's health Karolinska Institute. Stockholm
- Van der Zanden LFM, van Rooij IA, Feitz WF, Franke B, Knoers NV, Roeleveld N. 2012. Aetiology of hypospadias: a systematic review of genes and environment. *Hum Reprod Update* 18(3):260-83.
- Van der Zanden LFM, Galesloot TE, Feitz WFJ, Brouwers MM, Shi M, Knoers NV et al. 2012. Exploration of gene-environment interactions, maternal effects, and parent-of-origin effects in the etiology of hypospadias. *J Urol* 188(6): 2354-60.
- Van der Zanden LFM, van Rooij IAL, Feitz WFJ, Vermeulen SHH, Kiemeneys LA et al. 2010. Genetics of hypospadias: are single-nucleotide polymorphisms in SRD5A2, ESR1, ESR2, and ATF3 really associated with the malformation?. *J Clin Endocrinol Metab* 95(5):2384-90.
- Viddal KO, Aksnes G. 2008. Hypospadias-not only misplaced urinary tract opening. *Tidsskr Nor Lægeforen* 128:586-8.
- Wan S, Wang Y, Gu S. 2014. Epidemiology of male genital abnormalities: a population study. *Pediatrics* 133: e624-e627.
- Wang H, Jiang M, Cui H et al. 2012. The stress response mediator ATF3 represses androgen signaling by binding the androgen receptor. *Mol Cell Biol* 32:3190-202.

Wang MH, Baskin L.2008. Endocrine disruptors, genital review development, and hypospadia.J Androl 29(5) :499-505.

Withlock NC, Bahn JH, Lee SH, Eling TE, Seung JB. 2011. Resveratrol-induced apoptosis is mediated by early growth response-1, Krüppel-like factor 4, and activating transcription factor 3. Cancer Prev Resv(Phila) 4(1):116-27.

Xiang Pan Y, Chen H, Kilberg MS. 2005. Interaction of RNA-binding proteins HuR and AUF1 with the human ATF3 mRNA 3'-untranslated region regulates its amino acid limitation-induced stabilization. J Biol Chem 280:34609-16

Xu LF, Liang CZ, Lipianskaya J, Chen XG, Fan S, Zhang L et al. 2014. Risk factors for hypospadias in China. Asian J Andro 16: 778-81.

