

DAFTAR PUSTAKA

- Adrie, C., Azoulay, E., Francais, A., Clec'h, C., Darques, L., Schwebel, C., Nakache, D., Jamali, S., Goldgran-Toledano, D., dan Garrouste-Orgeas, M. (2007). Influence of gender on the outcome of severe sepsis: a reappraisal. *Chest*, 132 (6), 1786-1793.
- Almasdy, D., Deswinar, dan Helen. (2013). Evaluasi penggunaan antibiotika pada suatu rumah sakit pemerintah di kota Padang. *Prosiding Seminar nasional Perkembangan Terkini Sains Farmasi & Klinik III*. ISSN: 2339-2592, 7-15.
- Angus, D.C. dan Poll, T.V.D. (2013). Severe sepsis and septic shock. *N. Engl. J. Med*, 369 (9), 840-851.
- Ashley, C., Currie, A. (2009). *Renal Drug Handbook 3rd ed*. New York: Radcliffe Publishing.
- Baktygul, K., Martan, B., Ashirali, Z.M.D, Rashid, H., & Sakamoto, J. (2011). An assessment of antibiotics prescribed at the secondary health-care level in the Kyrgyz Republic. *Nagoya J. Med. Sci.*, 73 (3-4), 157-168
- Bauer, L. A. (2006). *Clinical Pharmacokinetics Handbook*. Washington: McGram Hill
- Bhattacharjee, P., Edelson, D.P., Churpek, M.M. (2017). Identifying patients with sepsis on the hospital wards. *CHEST*, 151(4), 898-907
- Bloch, K.C. (2003). Infectious Diseases. In McPhee SJ (ed.):*Pathophysiology of disease: An introduction to clinical medicine. 4th edition*. Lange Medical Books: McGraw-Hill.
- Brooks, G.F., Butel, J.S. dan Morse S.A., 2001. *Mycobacteriaceae in Jawetz medical microbiologi, 22ed*, McGraw-Hill Companies Inc:453-65
- Brun-Buisson, C., Doyon, F., Carlet, J., Dellamonica, P., Gouin, F., Lepoutre, A., Mercier, J.C., Offenstadt, G., dan Regnier, B. (1995). Incidence, risk factor, and outcome of severe sepsis and septic shock in adults: multicenter prospective study in Intensive Care Unit. *JAMA*, 274 (12), 968-974.
- Brunto, Laurence I., Lazo, John S., Parker, Keith L. (2006). *Goodman & Gilman the pharmacological basis of therapeutic 11th edition*. New York: McGraw-Hill
- Chidiac, C., Cavallo, J.D., Cohen, R., Dupon, M., Galperinee, T., Garraffof, R., Garog, B., Gauzith, R., Gavazzii, G., Kouzanj, S., Varonk, E., Lecomptel, T., Leroy, O., Sennevillen, E., Tattevino, P., Thiebault-Bertrandp, A., dan Voiriot, P. (2015). Update on a proper use of systemic fluoroquinolones

- in adult patients (ciprofloxacin, levofloxacin, moxifloxacin, norfloxacin, ofloxacin, pefloxacin). *Med Mal Infect*, 45(9), 348-373.
- Danai, P., Martin, G. (2005). Epidemiology of sepsis: Recent advances. *Current Infectious Disease Reports*, 7 (5), 329-334.
- Dellinger, R.P., Levy M.M., Rhodes A., Rhodes, A., Annane, D., Gerlach, H., Opal, S.M., Sevransky, J.E., Sprung, C.L., Douglas, I.S., Jaeschke, R., Osborn, T.M., Nunnally, M.E., Townsend, S.R., Reinhart, K., Kleinpell, R.M., Angus, D.C., Deutschman, C.S., Machado, F.R., Rubenfeld, G.D., Webb, S.A., Beale, R.J., Vincent, J.L., dan Moreno, R. (2013). Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock. *Crit Care Med*, 41 (2), 580-637.
- Elixhauser, A., Friedman, B., & Stranges, E. (2009). *Septicemia in U.S. Hospital*. Agency for Healthcare Research and Quality, Rockville, MD.
- Engel, C., Brunkhorst, F.M., Bone, H.G., Brunkhorst, R., Gerlach, H., Grond, S., Gruendling, M., Huhle, G., Jaschinski, U., John, S., Mayer, K., Oppert, M., Olthoff, D., Quintel, M., Ragaller, M., Rossaint, R., Stuber, F., Weiler, N., Welte, T., Bogatsch, H., Hartog, C., Loeffler, M., dan Reinhart, K. (2006). Epidemiology of sepsis in Germany: results from a national prospective multicenter study. *Intensive Care Med*, 33 (4), 606-618.
- Fatimah, S. (2006). *Buku ajar geriatri*. Jakarta: Fakultas Kedokteran Universitas Indonesia
- Ferrer, R., Martin-Loeches, I., Phillips, G., Osborn, T.M., Townsend, S., Dellinger, R.P., Artigas, A., Schorr, C., dan Levy, M.M. (2014). Empiric antibiotic treatment reduces mortality in severe sepsis and septic shock from the first hour: Results from a guideline-based performance improvement program. *Crit Care Med*, 42 (8), 1749-1755.
- Frizzel, J.P. (2001). *Handbook of Pathophysiology*. Philadelphia: Springhouse Corporation.
- Garmacho, M.J, Garcia, G.J., Barrero, A.A., Jimenez, F.J., dan Perez, P.C. (2003). Impact of adequate empiric antibiotic therapy on the outcome of patients admitted to the intensive care unit with sepsis. *Crit Care Med*, 31 (12), 2742-2751
- Guntur dan Hermawan, A. (2009). *Sepsis*. Dalam : Sudoyo, A.W., Setiyohadi, B., Alwi, I., Simadibrata, M., Setiati, S. (eds). Buku Ajar Ilmu Penyakit Dalam, Jilid III, Edisi IV. Jakarta: Pusat Penerbitan Departemen Ilmu Penyakit Dalam FKUI
- Gushka, H. (2015). Evaluasi penggunaan antibiotika pada pasien sepsis di ruang ICU Rumah Sakit Umum Daerah Serang. (*Thesis*). Yogyakarta: Universitas Gajah Mada

- Gyssens, I.C. (2005). *Audit for monitoring the quality of antimicrobial prescription*. New York: Kluwer Academic Publishers,
- Halawiyah, A. (2015). Evaluasi kualitatif penggunaan antibiotik meropenem pada pasien sepsis BPJS di RUMKITAL Dr. Mintohardjo tahun 2014. (*Skripsi*). Jakarta: UIN Syarif Hidayatullah Jakarta
- Hidayati. (2014). Kajian penggunaan antibiotik pada pasien sepsis dengan gangguan ginjal. (*Thesis*). Padang: Universitas Andalas
- Jawetz, E. (1997). *Principle of antimicrobial drug action, basic and clinical pharmacology, Third edition*. Appleton and Lange
- Katzung, G.B. (2008). *Farmakologi dasar dan klinik, edisi 8*. Alih Bahasa oleh Bagian Farmakologi Fakultas Kedokteran Universitas Airlangga. Jakarta: Salemba Medika
- Kementrian Kesehatan Republik Indonesia. (2011). *Peraturan menteri Kesehatan Republik Indonesia No. 246/MENKES/PER/XII/2011*. Jakarta: Kementrian Kesehatan Republik Indonesia
- Lagu T, Rothberg MB, Shieh MS, Pekow PS, Steingrub JS, Lindenauer PK. (2012). Hospitalizations, costs, and outcomes of severe sepsis in the United States 2003 to 2007. *Crit Care Med*, 40 (3), 754-761.
- Lestari, W. (2011). Studi penggunaan antibiotik berdasarkan sistem ATC/DDD dan Kriteria Gyssens di bangsal penyakit dalam RSUP Dr. M. Djamil Padang. (*Thesis*). Padang: Universitas Andalas
- Lomaestro, B.M. 2000. Fluoroquinoloneinduced renal failure. *Drug Safety*, 22 (6), 479–485
- Lucida, H., Trisnawati, R., dan Suardi, M. (2011). Analisis aspek farmakokinetika klinik pasien gagal ginjal pada irna penyakit dalam RSUP Dr. M. Djamil Padang. *Jurnal Sains dan Teknologi Farmasi*, 16 (2), 144-155.
- McEvoy & Gerald. (2008). *AHFS drugs information*. USA: American Society of health system pharmacists.
- Mutschler, E. (1991). *Dinamika Obat, Farmakologi dan Toksikologi*. Edisi V. Bandung: Institut Teknologi Bandung.
- Pamela, Sintia D. (2011). Evaluasi kualitatif penggunaan antibiotika dengan metode Gyssens di Ruang Kelas 3 Infeksi Departemen Ilmu Kesehatan Anak RSCM secara retrospektif. (*Skripsi*). Depok: Universitas Indonesia
- Rhodes, A., Evans, L.E., Alhazzani, W., Levy, M.M., Antonelli, M., Ferrer, R., Kumar, A., Sevransky, J.E., Sprung, C.L., Nunnally, M.E., Rochwerg, B., Rubenfeld, G.D., Angus, D.C., Dellinger, R.P. (2017). Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock: 2016. *Critical Care Medicine*, 45 (3), 486-552.

- Sacce, K.C., Nauenberg, E., Long, A., Anton, B., Tucker, H.J., & Hu, T.W. (1995). Long term survival after intensive care unit admission with sepsis. *Crit Care Med*, 23 (6), 1040-1047.
- Sagy, M., Al-Qaaqa, Y., dan Kim, P. (2013). Definition and pathophysiology of sepsis. *Curr Probl Pediatric Adolesc Health Care*, 43 (10), 260-263.
- Sattler, F.R., Colao, D.J., Caputo, G.M., dan Schoolwerth, A.C. (1985). Cefoperazone for Empiric Therapy in Patients with Impaired Renal Function. *The American Journal of Medicine*, 81 (12), 229-236.
- Seckel, M.A., & Ahrens, T. (2016). Challenges in sepsis care new sepsis definitions and fluid resuscitation beyond the central venous pressure. *Crit Care Nurs Clin*, 28 (4), 513-532.
- Setiabudi, R. (2007). *Farmakologi terapi dan klinik edisi lima*. Jakarta: Departemen Farmakologi dan Terapeutik FKUI
- Seymour, C.W., Liu, V.X., Iwashyna, T.J., Brunkhorst, F.M., Rea, T.D., Scherag, A., Rubenfeld, G., Kahn, J.M., Shankar-Hari, M., Singer, M., Deutschman, C.S., Escobar, G.J., dan Angus, D.C. (2016). Assessment of Clinical Criteria for Sepsis For the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, 315 (8), 762-774.
- Shargel, L., Wu-Pong, S., Yu, A. B. C. (2005). *Applied Biopharmaceutics and Pharmacokinetics. Fifth edition*. United States : The McGraw-Hill Companies.
- Soong, J. dan Soni, N. (2012). Sepsis: recognition and treatment. *Clin Med*, 12 (3), 276-280.
- Sugiyono. (2011). *Metode penelitian kuantitatif, kualitatif dan R&D*. Bandung: Alfabeta.
- Sweetman, S.C. (2009). *Martindale: the complete drug reference thirty-sixth edition*. London: Pharmaceutical Press.
- Tenover, F. C. (2006). Mechanisms of antimicrobial resistance in bacteria. *The American Journal of Medicine*, 119 (6), 3-10.
- Tjay T.H., Rahardja K. (2010). *Obat-obat penting, khasiat, penggunaan dan efek-efek sampingnya*. Jakarta: PT. Elex Media Komputindo
- Tripathi, K.D. (2003). *Antimicrobial drug: general consideration essential of medical pharmacology, fifth edition*. New Delhi: Jaypee Brothers Medical Publishers
- Tupchong, K., Koifman, A., dan Foran, M. (2015). Sepsis, severe sepsis, and septic shock: A review of the literature. *African Journal of Emergency Medicine*, 5 (3), 127-135.

Wells, B.G., Dipiro, J.T., Schwinghammer, T.L., dan Dipiro, C.V. (2015). *Pharmacotherapy handbook ninth edition*. New York: McGraw-Hill

World Health Organization. (2012). *The role of education in the rational use of medicine*. New Delhi: WHO

