

DAFTAR PUSTAKA

- Abdou, H. M. (1989). *Dissolution, Bioavailability and Bioeqivalency*. Pennsylvania : Mack Publishing Company.
- Aiache, J.M. (1982). *Farmasetika 2. Biofarmasi*. Edisi Kedua. Penerjemah: Widji Soeratri. Surabaya: Penerbit Airlangga University Press.
- Akbuga, J. Gursoy, A. & Kendi, E. (1988). The Preparation and Stability of Fast Release Furosemide-PVP Solid Dispersion. *Drug-Dev. Ind Pharm*, 14(10), P, 1439-1464.
- Anonim. (1982). *The Extra Pharmacopeia 28th ed.* The Pharmaceutical Press. London.
- Anonim. (1986). *Hand Book of Pharmaceutical Excipien*. Pharmaceutical Association. Washington.
- Anonim. (1987). *Farmakope Indonesia*, Ed III. Direktorat Jendral Pengawasan Obat dan Makanan, Departemen Kesehatan RI, Jakarta.
- Anonim. (1995). *Farmakope Indonesia*, Ed IV. Direktorat Jendral Pengawasan Obat dan Makanan, Departemen Kesehatan RI, Jakarta.
- Anonim. (2014). *Farmakope Indonesia*, Ed V. Direktorat Jendral Pengawasan Obat dan Makanan, Departemen Kesehatan RI, Jakarta.
- Ansel, H.C. (1989). *Pengantar Bentuk Sediaan Farmasi* (Edisi IV). Penerjemah :F. Ibarahim. Jakarta : UI Press.
- Banakar, Umesh V.(1991). *Pharmaceutical Dissolution Testing*, Marcell Dekker Inc. United State America.
- Bazegar-Jalali, M., Valizadeh, H., & Adibkia, K. (2007). Enhancing Dissolution Rate of Carbamazepine Via Co-Grinding with Crosspovidone and Hydroxypropylmethylcellulose. *Iranian Journal of Pharmaceutical Research*, 6(3), 159-165.
- Chiou, W.L., & S. Riegelman. (1971). Pharmaceutical Application of Solid Dispersion System. *J. Pharm. Sci.*, 60, 9: 128 – 1302.
- Chivate, N., Anuradha, C., Shubhangi, S., Jagdish, S. (2013). Improvement of bioavailability and solubility of telmisartan by solid dispersion technique using various carriers. *Int. J. Pharm. Sci. Rev. Res.*, 19 (2) : 36-41.
- Chono, S., Takeda, E., Seki, T., dan Marimoto, K. (2008). Enhanchement of Dissolution Rate and Gastrointestinal Absorbtion of Prankulast as a Model

- Poorly Water Soluble Drug by Grinding with Gelatin, *Int J 78 of Pharm*, 347, 71.
- David, B., Remington. (2005). *The Science and Practice of Pharmacy*, ed 21. Philadelphia : Lippincott William.
- Dachriyanus. (2004). *Analisis struktur Senyawa organik secara spektroskopi*. Padang: Universitas Andalas Press.
- Dollery, S. C. (1991). Therapeutic Drugs (Volume 1). London: *Churchill Livingstone*.
- Erizal, Goeswin, A., Sasanti, T. D. (2003). Studi Sistim Dispersi Padat Glibenklamid dalam Polivinil Pirolidon K-30. *J. Sains, Tek. Far*, 8(1), 1-5.
- Friedrich, H., Nada, A., & Bodmeir, R. (2005). Solid State and Dissolution Rate Characterization of Co-ground Mixture of Nifedipine and Hydrophilic Carriers. *Drug Development Industrial Pharmacy*, 31, 719-728.
- Garg, A., Singh, S., Rao, V. U., Bindu, K., & Balasubramaniam, J. (2009). Solid state interaction of raloxifene HCL with different hydrophilic carriers during co-grinding and its effect on dissolution rate, *Drug Development Industrial Pharmacy*, 35, 455-470.
- Goodman and Gillman. (2001). *The Pharmacological Basic of Theurapeutic*. 10th ed.
- Goldberg, A.H., Gibaldi, M., and Kanig, J.L. 1996. Increasing dissolution rates and gastrointestinal absorprtion of drugs via solid solutions and autectis mixture III. *J. Pham. Sci*, 55, 5, 487-492.
- Gue, E., et.al. (2013). Accelerated Ketoprofen Release from Polymeric Matrices : Importance of the Homogeneity / Heterogeneity of Excipient Distribution. *Int. J. Pharm.*, 457 (1): 298-307.
- Handoko, T, Soeharto, B. (1995). *Insulin Glukagon Dan Antidiabetik Oral*. Farmakologi dan Terapi, edisi IV. FKUI Jakarta.
- Harmita. (2007). *Pemanfaatan Kromatografi Cair Kinerja Tinggi Untuk Analisis Obat dalam Bentuk Sediaan Farmasi dan Bioanalisis*. Jakarta: Universitas Indonesia.
- Kamalakkannan, V., Puratchicody, A., Masilamani, K., and Senthilnathan, B. (2010). Solubility Enhacement of Poorly Soluble Drugs by Solid Dispersion Technique. *J. Pham. Reseach*.
- Khaleel, N. Y., Abdulrasool, A. A., Ghareeb, M. M. And Hussain S. A., (2011). Solubility and Dissolution Improvement of Ketoprofen by Solid Dipersion in Polymerand Surfactant Using Solven Evaporation Method. *Int. J. Of Pharmacy And Pharm Sci*, 3, Issue 4.

Khopkar, S. M. (2003). *Konsep Dasar Kimia Analitik*. Jakarta : Universitas Indonesia Press.

Klein, S., Wempe, M. F., Zoellera, T., Buchanan, N. L., Lambertb, J. L., Ramsey, M. G., Edgard K. J., & Buchanan, C. M. (2009). Improving glyburide solubility and dissolution by complexation with hydroxybutenyl-b-cyclodextrin. *JPP*, 61, 23-30.

Jagadeesan, R., & Adhakrishnan, M. (2013). Novel approaches in the preparation of solid dispersion on solubility: a review. *Int. J. Pharm. Pharm. Sci.*, 5 (3) : 1000-1004

Lachman, L., H.A. Lieberman & J.L. Kanig, 1989, *Teori dan Praktek Industri Farmasi*, terjemahan Siti Suyatmi, edisi III , UI Press, Jakarta.

Leuner, C., and Dressman, J. (2000). *Improving Drug Solubility for Oral Delivery Using Solid Dispersion*. Eur. J. Pharm. Biopharm.

Levy, G., (1963). Effect of Particle Size on Dissolution and Gastrointestinal Absorption Rates of Pharmaceutical. *Am. J. Pharm. Sci.*

Martin, A., J. Swarbrick., and A. Cammarata.(1990). *Farmasi Fisika*, Jilid 2, edisi-3, diterjemahkan oleh Yoshita, Jakarta : UI Press.

Mudgal, S.S., & Pancholi, S. (2012). Formulation of Glibenklamide Solid Dispersion by Solvent Evaporation Technique. *JOCPR*, 4(1):353-359.

Neha, Preeti, C., Atin, K., Rajan,, P., Kumar, M. R., Santanu, M., Pardeep, K., Munsab, A. (2012). Approaches To Improve The Solubility And Bioavaibility Of Poorly Soluble Drugs And Different Parameter To Screen Them. *Novel Science International Journal Of Pharmaceutical Science*, 1(4), 498-502.

Rajesh, Nawale, Bharat, N., & Mehta. (2013). Glibenclamide Loaded Self-Microemulsifying Drug Delivery System. *Int. J. of Pharm. Sc.*, 5, (2), 325-330.

Rheimer, L. (1998). *Scanning electron microscopy: Physics of image formation and microanalysis* (edisi 2). London: Springer.

Sareen, S., Mathew, G., & Joseph., L. (2014). Improvement in solubility of poor water-soluble drugs by solid dispersion: Review Article. *Int. J. Pharm. Invest.*, 2 (1) : 12-17.

Shargel, L., & Yu, A. (2005). *Biofarmasetika & farmakokinetika terapan*. Terjemahan oleh Fasich dan Budi Suprapti, Edisi kedua, Surabaya: Airlangga University Press.

Shefter, E. & Cheng, K.C. (1980). Drug Polyvinilpyrolidone Dispersion a Differential Scanning Calorimetric Study. *J. Pharm Sci.* 6., 179-182.

- Singh, S., Baghel, R. S., & Yadav, L. (2011). A Review on solid dispersion. *Int. J. of Pharm & Life Sci.* 2 (9): 1078-1095.
- Sekiguchi, K. and Obi, N., (1961). *Studies on Absorption of Eutectic Mixtures II*. Chem. Pharm. Bull.
- Sweetman, S. C. (2009). *Martindale The Complete Drug Reference* (36th ed). London: Pharmaceutical Press.
- Vander, A. H., Parikh, J. R., Parikh, R, H., &Solanki, A. B. (2009). Preparation and Characterization of Co-grinded Mixture of Aceclofenac and Neusilin Us₂ for enhancement of Aceclofenac, *AAPS Pharmaceutical Sciences And Technology*, 10(2).
- Voigh, R. (1994). *Buku pelajaran teknologi farmasi*. (edisi V). Penerjemah: Noerono, S. Yogyakarta: Gajah Mada University Press.
- Vogt M., Kunath K., Dressman J. B. (2008). Co-grinding Enhances the Oral Bioavaibility of EMD 57033, a Poorly Water Soluble Drug in Dogs. *Eur J Pharm Biopharm*, 68, 338-345.
- Watson, D.G (1999). *Pharmaceutical Analysis*. New York: Churchill Livingstone.
- Wismogroho, A.S. & Widayatno, W.B. (2012). Pengembangan Alat Differential Thermal Analysis untuk Analisa Termal Material Ca(OH)₂. *Telaah Jurnal Ilmu Pengetahuan dan Teknologi*, 30 (1): 7-12.
- Yadav, P.S., Kumar, V., Singh, U. P., Bhat, H. R., & Mazumder. (2013). Physicochemical characterization and in vitro dissolution studies of solid dispersion of ketoprofen with PVP K30 and d-mannitol. *Saudi Pharmaceutical Journal*. 21:77-84.
- Zaini, E., Witarsah, A., S., & Agustin, R. (2014). Enhancement of Dissolution Rate of Meloxicam by Co-Grinding Technique Using Hydroxypropylmethylcellulose. *J. Chem. Pharm. Res.*, 6(11):263-267.
- Zaini, E., Halim, A., Soewandi S. N., dan Sumirtapura, Y. C. (2010). Identifikasi Fisika antara Trimetoprim dab Sulfametoksazol dengan metode Kontak Kofler dan Reaksi Rekristalisasi. *Majalah Farmasi Indonesia.*, 21(1):263-267.
- Zaini, E., Halim, A., Soewandi S. N., dan Setiawan, Y. (2011). Peningkatan Laju Pelarutan Trimetoprim Melalui Metode Ko-kristalisasi dengan Nikotinamida. *Jurnal Farmasi Indonesia.*, 5:263-267.
- Zaini, E., Yeyet, C.S., Soewandhi, S.N.,&Halim, A. (2008). Identifikasi interaksi fisika antara trimetoprim dan sulfametoksazol dengan metode kontak Kofler dan reaksi kristalisasi. *Majalah Farmasi Indonesia.*, 21(1) : 32-39.