

## DAFTAR PUSTAKA

- Agoes G. Teknologi Bahan Alam. Bandung: ITB Press; 2007.
- Ahmadjian V. The Lichens. New York & London: Academic Press; 1993.
- Arifa N, Nofitasari Y, Putri WZ, Yuldi MG. Laporan Akhir Penelitian Program Kreativitas Mahasiswa : Formulasi dan Evaluasi Sediaan Gel dari Isolat Murni dan Ekstrak *Stereocaulon graminosum* yang Memiliki Aktivitas Antibakteri terhadap Methicillin-Resistant *Staphylococcus aureus* (MRSA) pada Luka Infeksi Kaki Diabetik; 2016.
- Backorova M, Jendzelovsky R, Kello M. Lichen secondary metabolites are responsible for induction apoptosis in HT-29 and A2780 human cancer cell line: Toxicol In Vitro. 2012; 26: 462-468.
- Betnina, V.. Bioutography in paper and thin layer chromatography and its scope in the antibiotic field: J. Chromatography. 1973; 78: 31-34.
- Brander GC, Pugh RJ, Bywater WL. Veterinary applied pharmacology and therapeutics.5th ed : Bailliere Tindall ELBS. 1991; 436: 467-473.
- Branislav R Ranković, Marijana M Kosanić, dan Tatjana P Stanojković. Antioxidant, Antimicrobial and Anticancer activity of the lichens *Cladonia furcata*, *Lecanora atra* and *Lecanora muralis*.*Complementary and Alternative Medicine* 11:97.2011
- Brooks GF, Butel JS, dan Morse SA. Jawetz, Melnick, & Adelberg's Medical Microbiology, 21st ed: Prentice Hall International Inc. 1998; 21: 145 –176.
- Candan M, Yilmaz T, Tay M, Erdem AÖ, Türk. Antimicrobial activity of extracts of the lichen *Parmelia sulcata*and its salazinic acid constituent: NatuRforsch C. 2007; 62(7/8): 619-621.
- Casselman KO, Casselman KL. Lichen Dyes. United States of America; 2001.
- Control for Disease and Prevention (CDC): diakses 16 Januari 2017 dari <https://www.cdc.gov/drugresistance/index.html>
- Crawford S . Ethnolichenology of *Bryoria fremontii* : wisdom of elders, population ecology, and nutritional chemistry. M.Sc. thesis University of Victoria Canada; 2007.
- Dachriyanus. Analisis Struktur Senyawa Organik Secara Spektroskopi. Padang: Andalas University Press; 2004.
- Darmadi. Infeksi Nosokomial:Problematika Dan Pengendaliannya. Jakarta :Penerbit Salemba Medika; 2008.

Davis WW, Stout TR. Disc Plate Methods of Microbiological Antibiotic Assay: Microbiology. 1971; 22(4): 659-665.

Departemen Kesehatan Republik Indonesia. Parameter Standar Umum Ekstrak Tumbuhan Obat: Direktorat Pengawasan Obat Tradisional; 2000.

Dévéhat, Tomasi FLS, Elix JA, Bernard A, Rouaud I, Uriac P, Boustie J. Stictic acid derivatives from the lichen *Usnea articulate* and their antioxidant activities: J. Nat. Prod. 2007; 70(7): 1218-1220.

Dobson FS. Lichens An Illustrated The Guide to British and Irish Species. Singapore: Stamford Press; 1992.

Elix J A. Biochemistry and secondary metabolites: Lichen Biology. T. H. Nash. Cambridge, Cambridge University Press: 154-180. 1996.

Engel KU, Schmidt J, Reuter S, Weckesser BH, Simno CM, Schempp. *Usnea barbata* extract prevents ultraviolet-b induced prostaglandin e2 synthesis and COX-2 expression in hacat keratinocytes. J. of Photochemistry and Photobiology. 2007; 84 : 9-14.

Esimone CO, Ofokansi MU, Adikwu EC, Ibezim DO, Abonyi GN, Odaibo DO, Olaleye. In vitro evaluation of the antiviral activity of extracts from the lichen *Parmelia perlata*(L.) Ach. againts three RNA viruses; J. Infect. Dev. Count. 2007; 1(3): 315-320.

Frengki. Isolasi, Elusidasi Struktur Dan Uji Bioaktivitas Kandungan Kimia Fraksi Etil Asetat Kulit Batang Tanaman *Calophyllum Macrophyllum Scheff* : (Tesis) Program Studi Magister Ilmu Kefarmasian. Depok: Universitas Indonesia; 2010.

Galun M, Shomer-Ilan A. Secondary Metabolic Products. In Galun M (ed) CRC Handbook of Lichenology Vol III. CRC: Boca Raton; 1988.

Gülçin, Oktay IM, Küfrevoğlu OI, Aslan A. Determination of antioxidant activity of lichen *Cetraria islandica* (L) Ach: J. Ethnopharmacol. 2002; 79(3): 325-329.

Gupta VK, Darokar MP, Saikia A, Pal A, Fatima A, Khanuja SPS. Antimycobacterial activity of lichens: Pharm.Biol. 2007; 45(3): 200-204.

Hale ME. Growth in The Lichen. New York and London :Academic Press; 1973.

Harborne JB. Metode Fitokimia: Penuntun Cara Modern Menganalisis Tumbuhan. Terbitan Kedua. Diterjemahkan oleh K. Padmawinata dn I. Soediro. Penerbit ITB. Bandung; 1987

Harmita. Buku Ajar Analisis Fisikokimia. Depok: Departemen Farmasi FMIPA UI, 205-211; 2006.

Hesse M, Meier H, Zeeh B. In spektroskopische methoden der organischen chemie, 4.ü berarbeitete auflage. New York : Georg Thieme Verlag Stuttgart; 1991.

Hickey BJ, Lumsden AJ, Cole ALJ, Walker JRJ. Antibiotic compounds from new zealand plant : methyl haematomate. an antifungal agent from *Stereocaulon Ramulosum*: New Zealand Natural Sciences 17; 1990.

Huneck S, Yoshimura I. Identification of lichen substances. Verlag Berlin Heidelberg New York : Springer; 1996.

Ingolfsdottir Hylands KP, Solberg Y. Structure of vesuvianic acid from *Stereocaulon* species. Phytochemistry. 1985; 25(2): 550-553.

Ismed F, Dévéhat FLLH, Delalande O, Sinbandhit S, Bakhtiar A, Boustie J. Lobarin from the sumatran lichen, *Stereocaulon halei*: Fitoterapia. 2012; 83: 1693–1698.

Jawetz E, Melnick JL, Adelberg EA. Review of medical microbiology ed. 10. University of California: San Francisco; 2001.

Kinoshita KT, Togawa A, Hiraishi Y, Nakajima K, Koyama T, Narui L, Wang K, Takahashi. Antioxidant activity of red pigments from the lichens *Lethariella sernanderi*, *L. cashmeriana*, and *L. sinensis*: J. Nat. Med. 2010; 64(1): 85-88.

Kristmundsdóttir T, Aradóttir RAE, Ingólfssdóttir K, Ogmundsdóttir RM. Solubilization of the lichen metabolite (+)-usnic acid for testing in tissue culture: J. Pharm. Pharmacol. 2002; 54(11): 1447-1452.

Lamb M. A conspectus of lichen genus *Stereocaulon* (Schreb.) Hoffm: Journ. Hattori Bot. Lab. 1977; 43: 191-335 (Nov. 1977).

Luo H, Yamamoto Y, Kim JA, Jung JS, Koh YH, Hur JH. Lecanoric acid, a secondary lichen substance with anti-oxidant properties from umbilicaria antarcticain maritime antarctica (King George Island): Polar. Biol. 2009; 32(7): 1033-104.

Manojlovic NT, Vasiljevic P, Juskovic M, Najman S, Jankovic S, Andjelkovic AM. HPLC analysis and cytotoxic potential of extracts from the lichen *Thamnolia vermicularisvar. Subuliformis*: J. Med. Plant. Res. 2010; 4(9): 817-823.

Morita H, Tsuchiya T. Antimitotic activity of lobanic acid and a new benzofuran, sakisacaulon a from *Stereocaulon sasakii*: Bioorganic and Medicinal Chemistry Letters. 2009; 19: 3679-3681.

Muggia L, Schmitt I, Grube M. Lichen as treasure chest of natural products: SIM News. 2009.

Muller K. 2011. Pharmaceutically relevant metabolite from lichen: Appl. Microbiol Biotechnol. 2011; 56: 9-16.

Nash TH. Lichen Biology 2nd ed. Cambridge :Cambridge University Press; 2008.

Nash, TH. Lichen Biology. Cambridge : Cambridge University Press; 1996.

Okuyama E, Umeyama K, Yamazaki M, Kinoshita Y, Yamamoto Y. Usnic acid and diffractive acid as analgesic and antipyretic components of *Usnea diffracta*: Plant. Med. 1995; 61(2): 113-115.

Paudel BH, Bhattacharai K, Chim H, Lee R, Sofronov L, Ivanova L, Poryadina JH, Yim. Estimation of antioxidant, antimicrobial activity and brine shrimp toxicity of plants collected from oymyakon region of the republic of sakha (yakutia): Biological Research. 2014; 47: 10.

Pratiwi ST. Mikrobiologi Farmasi, 176. Yogyakarta: Penerbit Erlangga; 2008.

Radji M. Buku Ajar Mikrobiologi Panduan Mahasiswa Farmasi dan Kedokteran, 107, 118, 201-207, 295. Jakarta : Buku Kedokteran EGC; 2011.

Ranković B (Editor). Lichen Secondary Metabolite Bioactive Properties and Pharmaceutical Potential. Serbia : Springer; 2015.

Rankovic B. Lichen Secondary Metabolites Bioactive Propesties and pharmaceutical Potential. New York : Springer; 2015.

Ranković B, Misić M, Sukdolak S. Antimicrobial activity of extracts of the lichens *Cladonia furcata*, *Parmelia caperata*, *Parmelia pertusa*, *Hypogymnia physodes* and *Umbilicaria polyphylla*. Br: J. Biomed. Sci. 2007; 64(4): 143-148.

Rao PS, Sarma KG, Seshadari. The Ultraviolet and Infrared Spectra Of Some Lichen Depsides and Depsidones. Departement Of Chemistry of New Delhi: Delhi; 1966.

Reeves DS, Philips I, William JD. Laboratory Methods in Antimicrobial Chemotherapy. Churchill Livingstone: New York; 1978.

Richardson, DHS. Lichens and Man. In Hawksworth DL, ed., Frontiers in Mycology; 1991.

Russo A, Piovano M, Lombardo L, Garbarino J, Cardile V. Lichen metabolites prevent UV light and nitric oxide mediated plasmid DNA damage and induce apoptosis in human melanoma cells: Life. Sci. 2008; 83(13-14): 468-474.

Sarker SD, Latif Z, Gray AI. Natural products isolation. In: Sarker SD, Latif Z, Gray AI, editors. Natural Products Isolation. 2nd ed:Totowa (New Jersey). Human Press. 2006;18 :6-10.

Seidel V. Initial and bulk extraction. In: Sarker SD, Latif Z, Gray AI, editors. Natural Products Isolation. 2nd ed: Totowa (New Jersey) : Humana Press Inc. 2006; 31-5.

Setiabudy R. Farmakologi dan Terapi. Jakarta : Balai Penerbit FKUI; 2009.

Silverstein RM, Webster FX, Kiemle DJ. Spectrometric Identification Of Organic Compounds. State University of New York: United State of Amerika; 2005.

Solhaug KA, Lind M, Nybakken L. Possible Functional roles of cortical depsides and medullary depsidones in the foliose hypogymnia physodes: Flora. 2009.

Stahl E. Apparatus and general techniques in TLC. Dalam : Stahl, E. (ed). Thin layer chromatography a laboratory handbook. Terj. dari Dunnschicht chromatographie, oleh Ashworth, M.R.F. Berlin: Springer-Verlag. 1969; 61-77.

Stocker – Worgotter E. Metabolic diversity of lichen-forming ascomycetous fungi : cultulring polyketide and shikimate metabolite production and PKS genes: Nat Prod Rep. 2008; 25:188-20.

Suleyman, H., F. Odabasoglu, A. Aslan, A. Cakir, Y. Karagoz, F. Gocer, M. Halici, Y. Bayir. Anti-inflammatory and Antiulcerogenic Effects of the Aqueous Extract of *Lobaria pulmonaria*(L.) Hoffm: Phytomedicine. 2003; 10(6-7): 552-557.

Tanas S, Odabasoglu F, Halici Z, Cakir A, Aygun H, Aslan A, Suleyman H. Evaluation of anti-inflammatory and antioxidant activities of *Peltigera rufescens* lichen species in acute and chronic inflammation models: J. Nat. Med. 2010; 64(1): 42-49.

Thadhani VM, Naaz Q, Choudhary MI, Mesaik MA, Karunaratne V. Enzyme inhibitory and immunomodulatory activities of the depsidone lobaric acid extracted from the lichen *Heterodermia* sp: J.Natn.Sci.Foundation Sri Lanka. 2014; 42 (2): 193-196.

Tokiwano T, Satoh H, Obara T, Hirota H, Yoshizawa Y, Yamamoto Y. A lichen substance as an antiproliferative compound against HL-60 human leukemia cells: 16-OAcetyl-leucotyllic acid isolated from *Myelochroa aurulenta*(Note): Biosci. Biotechnol. Biochem. 2009; 73(11): 2525-2527.

Touchstone, J.C. ,Dobbins, M.F. *Practice of thin layer chromatography*. Canada: John Wiley & Sons, 2-12. 1983

Turk A, Yilmaz OM, Kivanc M, Turk H. The antimicrobial activity of extracts of the lichen *Cetraria aculeate* and its protolichesterinic acid constituent. Z: NatuRforsch. C. 2003; 58(11/12): 850-854.

Vartia KO, Antibiotics in lichens. In: Ahmadjian V, Hale ME Jr (eds) The lichens. Academic, New York. 1973.

Vijayakumar C S, Viswanathan S, Reddy MK, Parvathavarthini S, Kundu SB, Sukumar E. Anti-inflammatory activity of (+) usnic acid: Fitoterapia. 2000; 71(5): 564-566.

Wagner HS, Blast. Plant drug analysis, a thin layer chromatography atlas: Springer. 1996.

Waksmundzka-Hajnos, Sherma MJ, Kowalska T. Overview of the field of TLC in phytochemistry and the structure of the book. Dalam: Waksmundzka-Hajnos, M., J. Sherma & T. Kowalska (eds): Thin layer chromatography in phytochemistry. Boca Raton: CRC. 2008; 1 -6.

WHO. Top 10 Causes of Death. [www.who.int/mediacentre/factsheets/fs310/en/](http://www.who.int/mediacentre/factsheets/fs310/en/) diakses 16 Januari 2017. 2011.

Yim JH, Kim C, Kim DC, Han SJ. Pharmaceutical and Food Compositions For Preventing Or Treating Diabetes Or Obesity. United States Patent Application Publication 2014. US; 2014.

Yurnaliza. Lichens (Karakterisasi, Klasifikasi dan Kegunaan). USU Digitalis Library. Medan; 2002.

