

## DAFTAR PUSTAKA

- Abraham, E. M., Kyriazopoulos, A., Parissi, Z. M., Sklavou, P. and Tsiovaras, C. N. 2010. Defoliation frequency effects on winter forage production and nutritive value of different entries of *Agropyron cristatum* (L.) Gaertn. *Spanish Journal of Agricultural Research*, 8(3), 703–712.
- Abraham, E. M., Parissi, Z. M., Sklavou, P., Kyriazopoulos, A. and Tsiovaras, C. N. 2009. Defoliation frequency effects on winter forage production and nutritive value of different entries of *Dactylis glomerata* L. *New Zealand Journal of Agricultural Research*, 52(3), 229–237. <https://doi.org/10.1080/00288230909510508>
- Adiningsih, S. dan Mulyadi. 1992. Alternatif teknik rehabilitasi dan pemanfaatan lahan alang-alang. Dalam S. Sukmana, - Suwardjo, J. S. Adiningsih, H. Subagjo, H. Suhardjo, dan Y. Prawirasumantri (Eds.), *Pemanfaatan lahan alang-alang untuk usaha tani berkelanjutan*. Prosiding Seminar Lahan Alang-alang, Bogor, Desember 1992 (pp. 29–50). Pusat Penelitian Tanah dan Agroklimat. Badan Litbang Pertanian.
- Akinrinde, A. A. 2006. Strategies for improving crops` use efficiencies of fertilizer nutrients in sustainable Agricultural systems. *Pakistan Journal of Nutrition*, 5, 185–193.
- Akobundu, I., and Agyakwa, C. W. (998. A *Handbook of West African Weeds* (2ed., revised and expanded)). International Institute of Tropical Agriculture, Oyo Road, PMB 5320, Ibadan, Nigeria.
- Albayrak, S., Türk, M. and Yüksel, O. 2011. Effect of row spacing and seeding rate on Hungarian Vetch yield and quality. *Turkish Journal of Field Crops*, 16(1), 54–58.
- Allen, B. L. and Hajek, B. F. 1989. Mineral occurrence in soil environment. In J. B. Dixon and S. B. Weed (Eds.), *Mineral in Soil Environments* (2nd ed., pp. 199–278). Soil Sci. Soc. Am. Madison, Wisconsin.
- Angonyissa, D. and Sinsin, B. 1998. Productivity and carrying capacity of natural grassland in Benin. *Revue d'Élevage et de Médecine Vétérinaire Des Pays Tropicaux*, 51(3), 239–246.
- Ansah, T., Osafo, E. L. K. and Hansen, H. H. 2010. Herbage yield and chemical composition of four varieties of Napier (*Pennisetum purpureum*) grass harvested at three different days after planting. *Agric. Biol. J. Am.*, 1(5), 923–929. <https://doi.org/10.5251/abjna.2010.1.5.923.929>
- Asiegbu, J. E. and Onyeonagu, C. C. 2008. Effect of cutting frequency and nitrogen application on herbage yield and nitrogen content of a degraded *Panicum maximum* pasture. *Nigerian Journal of Animal Production*, 35(1), 114–127.

- Assefa, G. 2013. Effect of plant height at cutting and previous frequency of defoliations on DM yield and nutritive value of hay made from final regrowth of *Hyparrhenia rufa* (Nees) grass. *Middle-East Journal of Scientific Reserach*, 17(8), 1055–1060. <https://doi.org/10.5829/idosi.mejsr.2013.17.08.11920>
- Ayub, M., Nadeem, M. A., Tahir, M., Ibrahim, M. and Aslam, M. N. 2009. Effect of nitrogen application and harvesting intervals on forage yield and quality of Pearl Millet (*Pennisetum americanum* L.). *Pak. J. Life Soc. Sci*, 7(2), 185–189.
- Bänziger, M., Edmeades, G. O., Beck, D. and Bellon, M. 2000. *Breeding for Drought and Nitrogen Stress Tolerance in Maize; From Theory to Practice*. Mexico, D.F.: CIMMYT.
- Barker, A. V. and Bryson, G. M. 2006. Nitrogen. In A. V. Barker and D. J. Pilbeam (Eds.), *Handbook of plant nutrition*. Taylor & Francis.
- Bates, L. S., Waldren, R. and Teare, I. . D. 1973. Rapid determination of free proline for water stress studies. *Plant and Soil*, 39(1), 205–207. <https://doi.org/10.1007/BF00018060>
- Bayble, T., Melaku, S., and Prasad, N. K. 2007. Effects of cutting dates on nutritive value of Napier (*Pennisetum purpureum*) grass planted sole and in association with Desmodium (*Desmodium intortum*) or Lablab (*Lablab purpureus*). *Livestock Research for Rural Development, Volume 19, Article #11. Retrieved July 21, 2017*. Retrieved from <http://www.lrrd.org/lrrd19/1/bayb19011.htm>
- Belaygue, C., Wery, J., Cowan, A. and Tardieu, F. 1996. Contribution of leaf expansion, rate of leaf appearance, and stolon branching to growth of plant leaf area under water deficit in White Clover. *Crop Science*, 36, 1240–1246. <https://doi.org/10.2135/cropsci1996.0011183X003600050028x>
- Bhatti, B. M., Mohammad, D., Sartaj, and Sultani, M.(1985. Effects of different inter-and intra-row spacing on forage yield and quality in elephant grass. *Pakistan J. Agri. Res*, 6(2), 107–112.
- Bogdan, A. (1977). *Tropical pasture and fodder plants (grasses and legumes)*. Longman Inc. New York.
- Bolat, I., Dikilitas, M., Ercisli, S., Ikinci, A. and Tonkaz, T. 2014. The effect of water stress on some morphological, physiological, and biochemical characteristics and bud success on apple and quince rootstocks. *The Scientific World Journal*, 2014. <https://doi.org/10.1155/2014/769732>
- Bos, H. J. 1999. *Plant morphology, environment, and leaf area growth in wheat and maize*. Landbouwuniversiteit te Wageningen.
- Boschini, C. 2015. Nutritional duality of mulberry cultivated and harvested for

- ruminant feeding. FAO animal production and health paper, Rome, Italy.
- Braga, G. J., Portela, J. N., Pedreira, C. G. S., Leite, V. B. O. and Oliveira, E. A. 2009. Herbage yield in Signalgrass pastures as affected by grazing management. *South African Journal of Animal Sciences*, 39(SUPPL. 1), 130–132. <https://doi.org/10.4314/sajas.v39i1.61168>
- Bray, E. A. 2007. Plant Response to Water-deficit Stress. In *Encyclopedia of Life Sciences* (pp. 1–5). Nature Publishing Group. <https://doi.org/10.1002/9780470015902.a0001298.pub2>
- Brima, F. I. A. 2007. *Effect of seed rate and NPK fertilization on growth, yield and forage quality of Rhodes Grass (Chloris L. kunth.)*. University of Khartoum, Sudan.
- Brink, G. E., Casler, M. D. and Jackson, R. D. 2014. Response of four temperate grasses to defoliation height and interval. *Communications in Biometry and Crop Science*, 9(1), 15–25.
- Brink, G. E., Casler, M. D. and Martin, N. P. (2010). Meadow fescue, tall fescue, and orchardgrass response to defoliation management. *Agronomy Journal*, 102(2), 667–674. <https://doi.org/10.2134/agronj2009.0376>
- Brink, G. E., Jackson, R. D. and Alber, N. B. 2013. Residual sward height effects on growth and nutritive value of grazed temperate perennial grasses. *Crop Science*, 53(5), 2264–2274. <https://doi.org/10.2135/cropsci2013.01.0068>
- Briske, D. D., and Richards, J. H. 1993. Physiology of plants recovering from defoliation. In *Proceedings of the XVII International Grassland Congress* (pp. 85–94).
- Budiman, B. 2013. Pengaruh pemupukan nitroden dan stress air terhadap bukaan stomata, kandungan klorofil dan akumulasi prolin tanaman rumput gajah (*Pennisetum purpureum* Schum.). *JITP*, 2(3), 159–166.
- Buitink, J., Claessens, M. M. A. E., Hemminga, M. A. and Hoekstra, F. A. 1998. Influence of watercontent and temperature on molecular mobility and intracellular glasses in seeds and pollen. *Plant Physiol*, 118(2), 531–541.
- Bumane, S. 2010. The influence of NPK fertilization on *Lolium perenne* L . forage quality. *Agronomy Research*, 8(2002), 531–536.
- Carow, R. N., Johnson, B. J. and Burns, R. E. 1987. Thatch and quality of Tifway bermudagrass turf in relation to fertility and cultivation. *Agronomy Journal*, v. 79(3), 524-530–1987 v.79 no.3. <https://doi.org/10.2134/agronj1987.00021962007900030025x>
- Casal, J. J., Sanchez, R. A. and Dereibus, V. A. 1986. The Effect of plant-density on tillering - the Involvement of R/Fr ratio and the proportion of radiation intercepted per plant. *Environmental and Experimental Botany*, 26(3), 365–

371.

- Chaves, M. M. and Oliveira, M. M. 2004. Mechanisms underlying plant resilience to water deficits: Prospects for water-saving agriculture. *Journal of Experimental Botany*, 55(407), 2365–2384. <https://doi.org/10.1093/jxb/erh269>
- Chen, C.-S., Hwa, Y.S., Wang, S. M. and Chang, Y.-K. 1999. The relationship between climatic factors and Acid-Detergent Fiber, Neutral-Detergent Fiber and crude protein contents in digitgrass. *Taiwan Livestock Res.*, 32(3), 155–265.
- Chen, C. S., Wang, S. M. and Hsu, J. T. 2006. Factors affecting in vitro true digestibility of Napiergrass. *Asian-Australasian Journal of Animal Sciences*, 19(4), 507–513.
- Christianse, S. and Svejcar, T. 1988. Grazing effects on shoot and root dynamics and above - and below-ground non structural carbohyfrat in Caucasian blustem. *Grass and Forage Science*, 43, 111–119.
- Ciríaco, E., Silva, D., Rejane, Mansur, J., Nogueira, C., Almeida, M. and Bandeira De Albuquerque, M. 2011. Drought stress and plant nutrition. *Plant Stress*, 5(Special issue 1), 32–41.
- Clavelo, T. and Razz, R. 1997. Tiller dynamics of guinea grass (*Panicum maximum*) under defoliation. In F. . O'Mara, F. . Wilkins, L. 't Mannetje, P. A. . Lovett, D.K. Rogers, and T. . Boland (Eds.), *Proceedings XVIII IGC 1997 Winnepeg, Manitoba*. Wageningen Academic Publisher, Netherland.
- Close, W. H. and Menke, K. H. 1986. *Selected Topics in Animal Nutrition. A Manual Prepared for the 3rd Hoheinheim Course on animal nutrition in the tropics and semi-tropics* (2nd ed.). The Institute of Animal Nutrition, University of Hoheinheim.
- Coi, G. J., Jung, E. S., Rim, Y. W., Lim, Y. C., Kim, K. Y., Sung, B. R. and Park, G. J. 2002. Effects of drill widths and nitrogen application levels in early spring on the growth characteristics and seed productivity of Italian ryegrass (*Lolium multiflorum* Lam.). *J. Korean Soc. GRassland Sci.*, 3, 221–226.
- Cortes, P. M. and Sinclair, T. R. 1986. Water relations of field-grown soybean under drought. *Crop Science*, 26. <https://doi.org/10.2135/cropsci1986.0011183X002600050031x>
- Cox, J. W., Cherney, D. R. and Hanchar, J. J. 1998. Row spacing, hybrid, and plant density effects on corn silage yield and quality. *Journal of Production Agriculture*, 11, 128–134. <https://doi.org/10.2134/jpa1998.0128>
- Crasta, O. R. and Cox, W. J. 1996. Temperature and soil water effects on maize growth, development yield, and forage quality. *Crop Science*, 36, 341–348. <https://doi.org/10.2135/cropsci1996.0011183X003600020022x>

- Cuomo, G. J., Blouin, D. C., Corkern, D. L., McCoy, J. E. and Walz, R. 1996. Plant morphology and forage nutritive value of three Bahiagrasses as affected by harvest frequency. *Agronomy Journal*, 88, 85–89. <https://doi.org/10.2134/agronj1996.00021962008800010018x>
- Cusicanqui, J. A. and Lauer, J. G. 1999. Plant density and hybrid influence on corn forage yield and quality. *Agronomy Journal*, 91(6), 911–915. <https://doi.org/10.2134/agronj1999.916911x>
- Danalatos, N. G., Archontoulis, S. V. and Mitsios, I. 2007. Potential growth and biomass productivity of *Miscanthus giganteus* as affected by plant density and N-fertilization in central Greece. *Biomass and Bioenergy*, 31(2–3), 145–152. <https://doi.org/10.1016/j.biombioe.2006.07.004>
- Daur, I. 2016. Feed value of blue panic (*Panicum antidotale* retz.) grass at different growth stages and under varying levels of humic acid in saline conditions. *Turkish Journal of Field Crops*, 21(2), 210–217. <https://doi.org/10.17557/tjfc.18296>
- de Freitas, F. P., da Fonseca, D. M., dos Santos Braz, T. G., Martuscello, J. A. and Santos, M. E. R. 2012. Forage yield and nutritive value of Tanzania grass under nitrogen supplies and plant densities. *Revista Brasileira de Zootecnia*, 41(4), 864–872. <https://doi.org/10.1590/S1516-35982012000400006>
- Delaauney, A. J. and Verma, D. P. S. 1993. Proline biosynthesis and osmoregulation in plants. *The Plant Journal*, 4, 215–223.
- Demir, Y. 2000. Growth and proline content of germinating wheat genotypes under ultraviolet light. *Turk J Bot*, 24, 67–70.
- Dhanda, S. S., Sethi, G. S. and Behl, R. K. 2004. Indices of drought tolerance in wheat genotypes at early stages of plant growth. *Journal of Agronomy and Crop Science*, 190(1), 6–12. <https://doi.org/10.1111/j.1439-037X.2004.00592.x>
- Dong, X. and Patton, J. 2011. Biomass Allocation in Four Prairie Grasses under Drought Stress. *CGREC Annual Report*, 1–4.
- Donkor, N. T., Bork, E. W., and Hudson, R. J. 2003. Defoliation regime effects on accumulated season-long herbage yield and quality in boreal grassland. *Journal of Agronomy and Crop Science*, 189(1), 39–46. <https://doi.org/10.1046/j.1439-037X.2003.00007.x>
- Dougall, H. W. and Bogdan, A. V. 1965. The Chemical Composition of the Grasses of Kenya—III. *East African Agricultural and Forestry Journal*, 30(4), 314–319. <https://doi.org/10.1080/00128325.1965.11662002>
- Douglas, M. M., Setterfield, S. A., Rossiter, N., Barratt, J. and Hutley, L. B. 2004. Effects of mission grass (*Pennisetum polystachyon* (L.) Schult.) invasion on fuel loads and nitrogen availability in a northern Australia tropical savanna.

- In* B. M. Sindel and S. B. Johnson (Eds.), *Weed management: balancing people, planet, profit. 14th Australian Weeds Conference, Wagga Wagga, New South Wales, Australia, 6-9 September 2004* (pp. 179–181). Weed Society of New South Wales, Sydney, Australia.
- du Toit, J. C. O. 2014. Growth and tiller production of *Themeda triandra* as affected by NPK fertilisation. *African Journal of Range & Forage Science*, 31(3), 229–232. <https://doi.org/10.2989/10220119.2014.899272>
- Earl, H. J. and Davis, R. F. 2003. Effect of drought stress on leaf and whole canopy radiation use efficiency and yield of maize. *Agron. Agron J.*, 24(1), 688–698.
- El-Gengaihi, S. and Abdallah, N. 1978. The effect of date of sowing and plant spacing on yield of seed and volatile oil of Fennel (*Foeniculum vulgare* Mill.). *Pharmazie*, 33(9), 605–606.
- Esrita, Ichwan, B. dan Irianto. 2011. Pertumbuhan dan hasil tomat pada berbagai bahan organik dan dosis Trichoderma. *Jurnal Penelitian Universitas Jambi Seri Sains*, 13(2), 37–42. <https://doi.org/ISSN 0852-8349>
- Fageria, N. K. 2009. *The use of nutrients in crop plants. Annals of Botany* (Vol. 105). CRC Press, Taylor & Francis Group. <https://doi.org/10.1093/aob/mcp227>
- Fahej, M. A. S. 2012. *Screening Switchgrass (Panicum virgatum L.) for Water Stress Tolerance*. Oklahoma State University.
- [FAO] Food and Agriculture Organization of the United Nations. 2009. *Pennisetum polystachyon* (L.) Schult. Retrieved March 15, 2013, from <http://www.fao.org/ag/AGP/AGPC/doc/GBASE/data/pf000300.htm>
- Farooq, M., Wahid, A., Fujita, N. K. D. and Basra, S. M. A. 2009. Plant drought stress : effects, mechanisms and management To cite this version : Review article. *Agronomy for Sustainable Development, Springer Verlag (Germany)*, 29(1), 185–212. <https://doi.org/10.1051/agro:2008021>
- Fitriatin, B. N., Yuniarti, A., Turmuktini, T. and Ruswandi, F. K. 2014. The effect of phosphate solubilizing microbe producing growth regulators on soil phosphate, growth and yield of maize and fertilizer efficiency on ultisol. *Eurasian J. of Soil Sci. Indonesia*, 101–107.
- Flagella, Z., Rotunno, T., Tarantino, E., Di Caterina, R. and De Caro, A. 2002. Changes in seed yield and oil fatty acid composition of high oleic sunflower (*Helianthus annuus* L.) hybrids in relation to the sowing date and the water regime. *European Journal of Agronomy*, 17(3), 221–230. [https://doi.org/10.1016/S1161-0301\(02\)00012-6](https://doi.org/10.1016/S1161-0301(02)00012-6)
- Foth, H. D. 1990. *Fundamentals of Soil Science* (8th ed.). John Wiley & Sons.

- Gajbhiye, B. R., Momin, Y. D. and Puri, A. N. 2013. Effect of FYM and NPK fertilization on growth and quality parameters of Lemongrass (*Cymbopogon flexuosus*). *Agriculture Science Research Journal*, 3(April), 115–120.
- Garg, B. K., Burman, U. and Kathju, S. 2004. The influence of phosphorus nutrition on the physiological response of moth bean genotypes to drought. *Journal of Plant Nutrition and Soil Science*, 167(4), 503–508. <https://doi.org/10.1002/jpln.200320368>
- Gates, R. N., Mislevy, P. and Martin, F. G. 2001. Herbage accumulation of three Bahiagrass populations during the cool season. *Agronomy Journal*, 93. <https://doi.org/10.2134/agronj2001.931112x>
- Geren, H., and Kavut, Y. T. 2015. Effect of different plant densities on the yield and some silage quality characteristics of giant king grass (*Pennisetum hybridum*) under mediterranean climatic conditions. *Turkish Journal of Field Crops*, 20(1), 85–91.
- Girousse, C., Bournoville, R. and Bonnemain, J. L. 1996. Water deficit induced changes in concentration in proline and some other amino acids in phloem sap of Alfalfa. *Plant Physiol*, 111, 109–113.
- Gittins, C. and Busso, C. 2010. Defoliation frequency affects morphophysiological traits in the bunchgrass *Poa ligularis*. *Revista Internacional de Botanica Experimental*, 79, 55–68. Retrieved from <http://www.scielo.org.ar/pdf/phyton/v79n1/v79n1a09.pdf>
- Göksoy, A. T., Demir, A. O., Turan, Z. M. and Dağıstü, N. 2004. Responses of sunflower (*Helianthus annuus* L.) to full and limited irrigation at different growth stages. *Field Crops Research*, 87(2–3), 167–178. <https://doi.org/10.1016/j.fcr.2003.11.004>
- Gomez, A. A and Gomez, K. A. 1984. *Statistical Procedures for Agricultural Research* (2nd ed., Vol. 6), John Wiley & Sons.
- Grant, C., Bittman, S., Montreal, M., Plenchette, C. and Morel, C. 2005. Soil and fertilizer phosphorus: Effects on plant P supply and mycorrhizal development. *Canadian Journal of Plant Science*, 85(1), 3–14. <https://doi.org/10.4141/P03-182>
- Graybill, J. S., Cox, W. J. and Otis, D. J. 1991. Yield and quality of forage maize as influenced by hybrid, planting date, and plant density. *Agronomy Journal*, 83, 559–564. <https://doi.org/10.2134/agronj1991.00021962008300030008x>
- Grusak, M. A., Broadley, M. R. and White, P. J. 2016. Plant Macro- and Micronutrient Minerals. *Ecyclopedia of Life Science (eLS)*, (2001), 1–6. <https://doi.org/10.1002/9780470015902.a0001306.pub2>
- Gunamanta, P. G. Winten, K. T. I. dan Sukasana, I. W. 2014. Pengaruh pupuk NPK dan pupuk kandang sapi terhadap produktivitas rumput *Setaria*

- splendida* Stapf. *Majalah Ilmiah Universitas Tabanan*, 11(8).
- Hajibabaee, M., Azizi, F. and Zargari, K. 2012. Effect of drought stress on some morphological, physiological and agronomic traits in various foliage corn hybrids. *American-Eurasian Journal of Agriculture & Environmental Science.*, 12(7), 890–896. <https://doi.org/10.5829/idosi.aejaes.2012.12.07.1751>
- Hakim, N., Nyakpa, Y., Lubis, A. M., Rusdi Saul, M., Diha, A., Ban Hong, G. dan Bailey, H. H. 1986. *Dasar-dasar Ilmu Tanah*. Universitas Lampung.
- Hamim, H. 2005. Photosynthesis of C3 and C4 species in response to increased CO<sub>2</sub> concentration and drought stress. *Hayati: Journal of Biosciences*, 12(4), 131–138. [https://doi.org/10.1016/S1978-3019\(16\)30340-0](https://doi.org/10.1016/S1978-3019(16)30340-0)
- Hanson, A. P. and Hitz, W. 1982. Metabolic response of mesophytes to plant water deficits. *Annual Review of Plant Physiology and Molecular Biology*, 33, 163–203.
- Hare, P. D. and Cress, W. A. 1997. Metabolic implications of stress-induced proline accumulation in plants. *Plant Growth Reg.*, 21, 79–102.
- Hassan, A., Zewdu, T., Urge, M. and Fikru, S. 2015. Effect of nitrogen fertilizer application on nutritive value of *Cenchrus ciliaris* and *Panicum maximum* grown under irrigation at Gode, Somali Region. *J. Nutr Food Sci*, 11(5), 1–6. <https://doi.org/10.4172/2155-9600.1000S11005>
- Henzell, E. 1971. Recovery of nitrogen from four fertilizers applied to Rhodes grass in small plots. *Aust. J. Exp. Agr.*, 11(51), 420–430.
- Heuzé, V. and Tran, G. 2011. Mission grass (*Pennisetum polystachyon*). Retrieved from <http://www.feedipedia.org/node/400>
- Hidayat, I. 2002. *Penggunaan bakteri Azospirillum sp. pada tanah podsolik merah kuning terhadap pertumbuhan, produksi dan kualitas rumput Setaria splendida* stapf. dan *Chloris gayana* kunth. Fakultas Peternakan, Institut Pertanian Bogor, Bogor.
- Hintz, H. F., Schryver, H. F. and Williams, J. 1985. Correlation coefficients between nutrients in forages. *Can. J. Anim. Sci.*, 65, 251–253.
- Hodgson, J. 1990. *Grazing management: Science into practice*. Harlow: Longman Group UK Ltd.
- Hoekstra, F. A., Golovina, E. A. and Buitink, J. 2001. Mechanisms of plant desiccation tolerance. *Trends in Plant Science*, 6, 431–438.
- Holechek, J. L. 1982. Fertilizer Effects on Above- and Below- ground Biomass of Four Species. *Journal of Range Management*, 35(1), 39–42.

- Humphreys, L. R. 1980. *A Guide to Better Pastures for The Tropics and Sub-tropics* (4th ed.). Wright Stephenson and Co., Australia.
- Iftikhar, A., Aslam Khan, M. and Qasim, M. 2003. Growth and Development of Different Turfgrasses as Influenced by Nitrogen Application and Leaf Nitrogen Contents. *International Journal of Agriculture & Biology*, 5(2), 175–178.
- Jaleel, C. A., Manivannan, P., Lakshmanan, G. M. A., Gomathinayagam, M. and Panneerselvam, R. 2008. Alterations in morphological parameters and photosynthetic pigment responses of *Catharanthus roseus* under soil water deficits. *Colloids and Surfaces B: Biointerfaces*, 61(2), 298–303. <https://doi.org/http://dx.doi.org/10.1016/j.colsurfb.2007.09.008>
- Jaleel, C. A., Maniyannan, P., Sankar, B., Kishorekumar, A. and Gopi, R. 2007. Water deficit stress mitigation by calcium chloride in *Catharanthus roseus*: Effects on oxidative stress, proline metabolism and indole alkaloid accumulation. *Colloids Surf. B. Biointerf.*, 60, 110–116.
- Jawad, M. M., Al-shahwany, A. W. and Khudhair, S. H. 2015. Effect of biochemical fertilizer on proline accumulation, catalase and peroxidase enzymes activity in leaves of two wheat cultivars (Ipa99 and Rabyaa) under water deficit stress. *Iraqi Journal of Science*, 56(2), 1350–1358.
- Jones, R. M., Tothill, J. C. and Jones, R. J. 1984. *Pastures and pasture management in the tropics and sub-tropics*. Tropical Grassland Society of Australia.
- Jupp, A. P., and Newman, E. I. 1987. Morphological and anatomical effects of severe drought on the roots of *Lolium perenne* L. *New Phytologist*, 105, 393–402. <https://doi.org/10.1111/j.1469-8137.1987.tb00876.x>
- Kamel, M. S., Abdel-Raouf, M. S., El-Din, S. A. and Abbas, T. 1983. Effects of cutting height and frequency and nitrogen application rate on growth and forage yield of Napier grass, *Pennisetum purpureum* Schum. *Annals of Agricultural Science, University of Ain Shams (Egypt)*, 28(607–625).
- Karagić, D., Mihailović, V., Katić, S., Mikić, A., Milić, D., Vasiljević, S. and Milošević, B. 2011. Effect of row spacing on seed yield of hairy, common and Hungarian vetches. *Romanian Agricultural Research*, (28), 143–150.
- Khan, A. ., McNeilly, T. and Azhar, F. 2001. Stress tolerance in crop plants. *Int. J. Agric. Biol.*, 3, 250–255.
- Khan, R. I., Alam, M. R. and Amin, M. R. 1999. Effect of season and fertilizer on species composition and nutritive value of native grasses. *Asian-Australasian Journal of Animal Sciences*, 12(8), 1222–1227. <https://doi.org/10.5713/ajas.1999.1222>
- Khorshidi, J., Fakhr Tabatabaie, M., Omidbaigi, R. and Sefidkon, F. 2009. The

- Effect of Different Densities of Planting on Morphological Characters, Yield, and Yield Components of Fennel (*Foeniculum vulgare* Mill cv. Soroksary). *Journal of Agricultural Science*, 1(2), 66–73. <https://doi.org/10.5539/jas.v1n2p66>
- Kilcher, M. 1981. Plant Development, Stage of Maturity and Nutrient Composition. *Journal of Range Management*, 34(5), 363–364.
- Kitaba, A. and Tamir, B. 2007. Effect of harvesting stage and nutrient levels on nutritive values of natural pasture in central highlands of Ethiopia. *Agricultura Tropica et Subtropica*, 40(November 2001), 7–13. Retrieved from [http://www.projects.its.cz/ats/pdf\\_files/vol\\_40\\_1\\_pdf/KITABA.pdf](http://www.projects.its.cz/ats/pdf_files/vol_40_1_pdf/KITABA.pdf)
- Kitessa, S., Flinn, P. C. and Irish, G. G. 1999. Comparison of methods used to predict the in vivo digestibility of feeds in ruminants. *Australian Journal of Agricultural Research*, 50(5), 825–841. <https://doi.org/10.1071/AR98169>
- Kramer, P. J. 1969. *Plant and water relationships. A modern synthesis*. McGraw-Hill, New York.
- Kristyowantari, R. 1992. *Pengaruh interval dan tinggi pemotongan terhadap produksi dan beberapa aspek kualitas rumput Raja*. Institut Pertanian Bogor, Bogor.
- Kumar, D. and Nikhil, K. 2016. Effect of FYM , NPK and Algal fertilizers on the Growth and Biomass of Vetiver Grass (*Vetiveria nass*, L). *International Journal of Engineering and Applied Sciences (IJEAS)*, 3(3), 85–89.
- Kumar, G. S., Madhusudhan, K. V., Sreenivasulu, N., and Sudhakar, C. 2000. Stress responses in two genotypes of mulberry (*Morus alba* L.) under NaCl salinity. *Indian Journal of Experimental Biology*, 38(2), 192–195.
- Kuoppala, K., Rinne, M., Nousiainen, J. and Huhtanen, P. 2008. The effect of cutting time of grass silage in primary growth and regrowth and the interactions between silage quality and concentrate level on milk production of dairy cows. *Livestock Science*, 116(1), 171–182. <https://doi.org/http://dx.doi.org/10.1016/j.livsci.2007.10.002>
- Kusuma, M. E. (2014). Respon rumput gajah (*Pennisetum purpureum*) terhadap pemberian pupuk majemuk. *Jurnal Ilmu Hewani Tropika*, 3(1), 6–11.
- Kusumawati, I. N., Witariadi, N. M., Budiasa, I. K., Suranjaya, I. G. dan Roni, N. G. K. 2016. Pengaruh jarak tanam dan dosis bio-urin terhadap pertumbuhan dan hasil rumput *Panicum maximum* pada pemotongan ketiga. *Dalam Seminar Nasional V HITPI: Intensifikasi sistem produksi hijauan pakan untuk penguatan ketahanan pangan* (pp. 15–20).
- Kusvuran, A., Kaplan, M. and Nazli, R. I. 2014. Effects of mixture ratio and row spacing in Hungarian Vetch (*Vicia pannonica* Crantz.) and annual Ryegrass (*Lolium multiflorum* Lam.) intercropping on yield and quality under semiarid

- climate conditon. *Turkish Journal of Field Crops*, 19(1), 118–128.
- Kuwahara, F. A., Souza, G. M., Guidorizi, K. A., Costa, C. and Meirelles, P. R. de L. 2016. Phosphorus as a mitigator of the effects of water stress on the growth and photosynthetic capacity of tropical C4 grasses. *Acta Scientiarum. Agronomy*, 38(3), 363. <https://doi.org/10.4025/actasciagron.v38i3.28454>
- Lee, S. 1988. *Weed to watch: Pennisetum polystachion*. MACA Newsletter vol 1 (2): 11-12.
- Li, Y. Z., Wang, F. X. and Liu, L. H. 1999. Use and management of soil water and nitrogen resource. I. Soil water and nitrogen conditions and root development. *Plant Nutr. Fert. Sci.*, 5, 206–313.
- Liang, J. C. 1982. The response of Napier grass (*Pennisetum purpureum*) to animal manure and chemical fertiliser. 1. Effects on dry matter yield and quality. *Journal of the Agricultural Association of China (Taiwan)*, 119(64–74).
- Lingga, P. and Marsono, M. 2007. *Petunjuk Penggunaan Pupuk*. Penebar Swadaya, Jakarta.
- Lounglawan, P., Lounglawan, W. and Suksombat, W. 2014. Effect of cutting interval and cutting height on yield and chemical composition of King Napier Grass (*Pennisetum Purpureum x Pennisetum Americanum*). *APCBEE Procedia*, 8(Caas 2013), 27–31. <https://doi.org/10.1016/j.apcbee.2014.01.075>
- Lugiy, L. dan Sumarto, S. 2000. Teknik budidaya Rumput Gajah cv. Hawaii (*Pennisetum purpureum*). In *Temu Teknis non Peneliti* (pp. 120–125). Pusat Penelitian dan Pengembangan Pertanian. Departemen Pertanian.
- Lyon, D. J. 2009. How Do Plant Populations Affect Yield? *Press Releases from Panhandle Research and Extension Center*, 19, 1–4. Retrieved from <http://digitalcommons.unl.edu/panpressrel%5Cnhttp://digitalcommons.unl.edu/panpressrel/19%5Cnhttp://digitalcommons.unl.edu/panpressrel%5Cnhttp://digitalcommons.unl.edu/panpressrel/19>
- Mahajan, S. dan Tuteja, N. 2005. Cold, salinity and drought stress. *Archives of Biochemistry and Biophysics*, 444, 139–158. <https://doi.org/10.1002/9783527628964.ch7>
- Mahmood, A., Ullah, H., Shahzad, A. N., Ali, H., Ahmad, S., Zia-Ul-Haq, M. Hasanuzzaman, M. 2013. Dry matter yield and chemical composition of sorghum cultivars with varying planting density and sowing date. *Sains Malaysiana*, 42(10), 1529–1538.
- Malhi, S. dan Gill, K. 2004. Fertilizer management of forage crops in the Canadian Great Plains. *Recent Res Dev Crop* 1, 1–50. Retrieved from [http://www1.foragebeef.ca/\\$foragebeef/frgebeef.nsf/all/frg90/\\$FILE/fertilizermanagementofforagecropscanadiangreatplains.pdf](http://www1.foragebeef.ca/$foragebeef/frgebeef.nsf/all/frg90/$FILE/fertilizermanagementofforagecropscanadiangreatplains.pdf)

- Man, D., Bao, Y. X., Han, L. B. dan Zhang, X. 2011. Drought tolerance associated with proline and hormone metabolism in two tall fescue cultivars. *HortScience*, 46(7), 1027–1032.
- Maralian, H. dan Ebadi, A. 2010. Influence of water deficit stress on wheat grain yield and proline accumulation rate. *African Journal of Agricultural Research*, 5(4), 286–289. Retrieved from <http://www.academicjournals.org/journal/AJAR/article-abstract/A9B8F3837406>
- May, A., de Souza, V. F., Gravina, G. D. and Fernandes, P. G. 2016. Plant population and row spacing on biomass sorghum yield performance. *Ciencia Rural, Santa Maria*, 46(3), 434–439. <https://doi.org/10.1590/0103-8478cr20141133>
- McCue, K. F. dan Hanson, A. D. 1990. Drought and salt tolerance: towards understanding and application. *Trends in Biotechnology*, 8, 358–362.
- McIlroy, R. J. 1976. *Pengantar budidaya padang rumput tropika ; diterjemahkan Subadio, I., Susetyo, S. dan Soedarmadi, H.* Pradnya Paramita, Jakarta.
- Meier, H., and Reid, J. S. G. 1982. Reserve polysaccharides other than starch in higher plants. In F. A. Loewus dan W. Tanner (Eds.), *Encyclopedia of Plant Physiology* (pp. 418–471). Springer-Verlag, Berlin.
- Mengel, K. dan Kirby, E. A. 2001. *Principles of Plant Nutrition* (5th ed.). Kluwer Academic Publishers.
- Meyer, R. F., dan Boyer, J. S. 1981. Osmoregulation, solute distribution, and growth in soybean seedlings having low water potentials. *Planta*, 151(5), 482–489. <https://doi.org/10.1007/BF00386543>
- Miller, I. 2006. Management of Mission Grass (*Pennisetum polystachyon*) Agnote. No: F38. Biosecurity and Product Integrity, Department of Primary Industry, Fisheries and Mines, Northern Territory Government, Darwin, Northern Territory.
- Mishra, I. N. dan Sandhya, J. 1996. Nutritive profile of some grasses of Darbhanga. *Environment and Ecology*, 14(1), 93–95.
- Mislevy, P., Burton, G. W. dan Busey, Y. 1990. Bahiagrass Response to Grazing Frequency. *Soil Crop Sci. Soc. Fla.*, 50, 58–64.
- Mounika, B., Chellamuthu, V. and Sridevi, V. 2015. Plant Spacing influence on the Relative Productivity of Bajra Napier Hybrid Grass. *National Academy of Agricultural Science (NAAS)*, 33(2), 875–878.
- Mukhtar, M. 2006. Dry matter productivity of the dwarf and normal Elephantgrasses as affected by the planting density and cutting frequency. *Jurnal Ilmu Ternak Dan Veteriner*, 11(3), 198–205.

- Mulyani, A., Rachman, A. dan Daira, A. 2010. Penyebaran lahan masam, potensi dan ketersediaannya untuk pengembangan pertanian. *Dalam Prosiding Simposium Nasional Pendayagunaan Tanah Masam*. Pusat Penelitian dan Pengembangan Tanah dan Agroklimat. Bogor.
- Murphy, L. 1996. *Effects of row spacing on dryland forage grass quality Abstract*: Montana State University.
- National Research Council. 2000. *Nutrient Requirements of Beef Cattle* (Seventh Re). The National Academy Press.
- Navie, S. C. and Adkins, S. W. 2007. Environmental Weeds of Australia. Centre for Biological Information and Technology, The University of Queensland, Brisbane, Queensland. Brisbane, Queensland: Centre for Biological Information and Technology.
- Nevens, F. and Rehuel, D. 2003. Effects of cutting or grazing grass swards on herbage yield, nitrogen uptake and residual soil nitrate at different levels of N fertilization. *Grass and Forage Science*, 58(4), 431–449. <https://doi.org/10.1111/j.1365-2494.2003.00396.x>
- Ningalo, R. R., Kaligis, D. A. dan Bawole, N. 2017. Pengaruh defoliasi dan level pupuknitrogen terhadap performans rumput *Brachiaria humidicola* (Rendle) Schweick cv. Tully. *Jurnal Zootek*, 37(1), 25–32.
- Nohong, B. and N. S. 2015. Effect of water stress on growth , yield , proline and soluble sugars contents of Signal grass and Napier grass species. *American-Eurasian Journal of Sustainable Agriculture*, 9(5), 14–21.
- Nonami, H. 1998. Plant water relation and control of cell elongation at low water potentials. *Journal of Plant Reseach*, 111(373–382).
- Noogle, G. R. dan Fritz, G. J. 1983. *Introductory Plant Physiology*. Prentice-Hall, Inc. Englewood Cliffs. New Jersey.
- Nopriani, U., Karti, P. dan Prihantoro, I. 2014. Produktivitas Duckweed (*Lemna minor*) sebagai hijauan pakan alternatif ternak pada intensitas cahaya yang berbeda. *JITV*, 19(4), 272–286.
- Okwori, A. I. and Magani, I. E. 2010. Influence of nitrogen sources and cutting interval on the digestibility of four (4) grass species in the southern guinea savanna of Nigeria. *Agriculture and Biology Journal of North America*, 1(4), 526–533. Retrieved from <http://scihub.org/ABJNA/PDF/2010/4/1-4-526-533.pdf>
- Olanite, J. A., Anele, U. Y., Arigbede, O. M., Jolaosho, A. O. and Onifade, O. S. 2010. Effect of plant spacing and nitrogen fertilizer levels on the growth, dry-matter yield and nutritive quality of Columbus grass (*Sorghum almum* stapf) in southwest Nigeria. *Grass and Forage Science*, 65(4), 369–375. <https://doi.org/10.1111/j.1365-2494.2010.00755.x>

- Onyeonagu, C. and Asiegbu, J. . 2005. Effects of cutting management and N-fertilizer application on plant height, tiller production and percentage dry matter in a run-down *Panicum Maximum* pasture. *Journal of Agriculture, Food, Environment and Extention*, 4(2), 28–33.
- Osório, J., Osório, M. L., Chaves, M. M. and Pereira, J. S. 1998. Water deficits are more important in delaying growth than in changing patterns of carbon allocation in *Eucalyptus globulus*. *Tree Physiology*, 18, 363–373. <https://doi.org/10.1093/treephys/18.6.363>
- Overman, A. R. and Scholtz, R. V. 2003. Dry matter production and cutting interval for perennial grasses. *Communications in Soil Science and Plant Analysis*, 34(June 2011), 225–229. <https://doi.org/10.1081/CSS-120017427>
- Pandey, R. 2015. Mineral nutrition of plants. In B. Bahadur, M. V. Rajam, L. Sahijram, and K. V. Krishnamurthy (Eds.), *Plant Biology and Biotechnology*. Springer India. <https://doi.org/10.1007/978-81-322-2286-6>
- Partridge, I. J. 1986. Effect of stocking rate and superphosphate level on an oversown fire climax grassland of mission grass (*Pennisetum polystachyon*) in Fiji. *Tropical Grasslands*, 20(4), 166–173.
- Patidar, M. and Rajora, M. P. 2009. Forage yield and quality of different genotypes of Buffel grass (*Cenchrus ciliaris*) at various cutting intervals. *Annals of Arid Zone*, 48(1), 51–55.
- Patil, V. C., Al-Gaadi, K. A., Madugundu, R., Tola, E., Marey, S., Mulla, D. and Upadhyaya, S. (2016). Response of Rhodes grass to variable rate application of irrigation water and fertilizer nitrogen. *Pakistan Journal of Agricultural Sciences*, 53(3), 599–607. <https://doi.org/10.21162/PAKJAS/16.3491>
- Pebrriansyah, A. 2012. Effect of drought stress and addition Arbuscula Mychorizal Fungi (AMF) on growth and productivity of tropical grasses (*Chloris gayana*, *Paspalum dilatatum*, and *Paspalum notatum*). *Pastura*, 2(1), 41–48.
- Peiretti, P. G. 2009. Effects of growth stage on chemical composition, organic matter digestibility, gross energy and fatty acid content of safflower (*Carthamus tinctorius* L.). *Ivestock Research for Rural Development*. Volume 22, Article #206. Retrieved July 21, 2017, from <http://www.lrrd.org/lrrd21/12/peir21206.htm>.
- Peng, Z., Lu, Q. and Verma, D. P. S. 1996. Reciprocal regulation of Delta(1)-pyrroline-5-carboxylate synthetase and proline dehydrogenase genes controls proline levels during and after osmotic stress in plants. *Mol Gen Genet*, 253(3), 334–41.

- Plensicar, M. and Kustori, R. 2005. Corn yield and water use as influenced by irrigation level, N rate and planting populations. *Trans. Kansan Acad. Science*, 53(4), 121–7.
- Pontes, L. S., Carrère, P., Andueza, D., Louault, F. and Soussana, J. F. 2007. Seasonal productivity and nutritive value of temperate grasses found in semi-natural pastures in Europe: Responses to cutting frequency and N supply. *Grass and Forage Science*, 62(4), 485–496. <https://doi.org/10.1111/j.1365-2494.2007.00604.x>
- Prado, F. E., Boero, C., Gallarodo, M. and Gonzales, J. 2000. Effect of NaCl on germination, growth and soluble sugar content in *Chenopodium quinoa* Willd. seeds. *Bot Bull Acad Sin*, 41, 27–34.
- Prajapati, K. and Modi, H. A. 2012. The importance of potassium in plant growth - a review. *Indian Journal of Plant Sciences*, 1(2–3), 177–186.
- Prasetyo, B. H., Subardja, D. dan Siswanto, A. B. 2000). Karakteristik, potensi, dan teknologi Pengelolaan tanah ultisol untuk Pengembangan pertanian lahan Kering di Indonesia. *Litbang Pertanian*, 23, 1–12.
- Prasetyo, B. H., Suharta, N. and Subagyo, H. 2000). Chemical and mineralogical properties of ultisols of Sasamba area, East Kalimantan.
- Priyadarshani, N. D. N., Amarasinghe, M. K. T. K., Subasinghe, S. and Kumarasinghe, I. R. P. and H. K. M. 2013. Effect of organic and inorganic fertilizers on biomass production, oil yield and quality Vetiver (*Vetiveria zizanioides* L.). *The Journal of Agricultural Sciences*, 8(1), 28–35.
- Purbajanti, E. D., Anwar, S. and Kusmiyati, F. 2012. Drought stress effect on morphology characters , water use efficiency , growth and yield of guinea and napier grasses. *International Research Journal of Plant Science*, 3(4), 47–53.
- Purbajanti, E. D., Soetrisno, R. R., Hanudin, E. dan Budhi, S. P. S. 2011. Produksi, kualitas, dan kecernaan in vitro tanaman rumput benggala (*Panicum maximum*) pada lahan salin. *Buletin Peternakan*, 35(1), 30–37.
- Rafiei, M. 2009. Influence of tillage and plant density on mungbean. *American-Eurasian Journal of Sustainable Agriculture*, 3(4), 877–880. Retrieved from <http://www.aensiweb.net/AENSIWEB/aejsa/aejsa/2009/877-880.pdf>
- Riaz, A., Younis, A., Hameed, M. and Kiran, S. 2010. Morphological and biochemical responses of turf grasses to water deficit conditions. *Pakistan Journal of Botany*, 42(5), 3441–3448.
- Ruggia Chiesa, A. P., Kozloski, G. V., Bonnecarrère Sanchez, L. M., Lima, L. D., Oliveira, L., Härtter, C. J. and Cadorn, R. L. 2008. Age of regrowth as a factor affecting the nutritive value of hay of kikuyu grass (*Pennisetum clandestinum*) offered to lambs. *Grass and Forage Science*, 63(2), 193–201.

<https://doi.org/10.1111/j.1365-2494.2007.00624.x>

- Rusdy, M. 2010. Dry matter production, carbohydrtae reserve content and nitrogen utilization in some tropical grasses as influenced by nitrogen fertilization and age of plants. *JITP*, 1(1), 28–34.
- Saddam, S., Bibi, A., Sadaqat, H. A. and Usman, B. F. 2014. Comparison of 10 Sorghum (*Sorghum bicolor* L) Genotypes under various Water stress Regimes. *The Journal of Animal & Plant Sciences*, 24(6), 1811–1820.
- Sajimin, Kompiang, I., Supriyati, dan Suratmin, M. 2001. Penggunaan biofertilizer untuk peningkatan produktivitas hijauan pakan rumput gajah (*Pennisetum purpureum* cv. Africa) pada lahan marjinal di Subang Jawa Barat. *Media Peternakan*, 24(2), 46–50.
- Salehi, A., Tasdighi, H. and Gholamhoseini, M. 2016. Evaluation of proline, chlorophyll, soluble sugar content and uptake of nutrients in the German chamomile (*Matricaria chamomilla* L.) under drought stress and organic fertilizer treatments. *Asian Pacific Journal of Tropical Biomedicine*, 6(10), 886–891. <https://doi.org/10.1016/j.apjtb.2016.08.009>
- Samaras, Y., Bressan, R. A., Csonka, L., Garcia-Rios, M. G., Paino, D. and Rhodes, D. 1995. Proline accumulation during water deficit. In N. Smirnoff (Ed.), *Environment and plant metabolism. Flexibility and acclimation* (pp. 161–187). BIOS Scientific Publishers.
- Sanderson, M. A., Voigt, P. and Jones, R. M. 2014. Yield and quality of warm-season grasses in central Texas. *J. Range Management*, 52(2), 145–150.
- Sarieff, S. 1985. *Kesuburan dan Pemupukan Tanah Pertanian*. CV. Pustaka Buana, Bandung.
- Särkijärvi, S., Sormunen-Cristian, R., Heikkilä, T., Komppa, J., Rinne, M. and Saastamoinen, M. 2008. Effect of grass species and time of cutting on in vivo digestibility in horses and sheep. In M. Saastamoinen (Ed.), *Nutrition of the exercising horse : 4th European Workshop on Equine Nutrition (EWEN)* (p. 26). MTT Agrifood Research Finland.
- Sarwar, M., Mahr-un-Nisa, Ajmal Khan, M. and Mushtaque, M. 2006. Chemical composition, herbage yield and nutritive value of *Panicum antidotale* and *Pennisetum orientale* for Nili buffaloes at different clipping intervals. *Asian-Australasian Journal of Animal Sciences*, 19(2), 176–180.
- Sawwan, J., Shibli, R. a., Swaidat, I. and Tahat, M. 2000. Phosphorus regulates osmotic potential and growth of African violet under in vitro-induced water deficit. *Journal of Plant Nutrition*, 23(6), 759–771. <https://doi.org/10.1080/01904160009382057>
- Serrano, L. J. P. and Pugnaire, F. I. 1999. Constraints by Water Stress on Plant Growth. In M. Pessarakli (Ed.), *Handbook of Plant and Crop Stress* (pp.

- 271–283). CRC Press.
- Seseray, D. Y., Santoso, B. dan Lekitoo, N. 2013. Produksi rumput gajah (*Pennisetum purpureum*) yang diberi pupuk N, P dan K dengan dosis 0 , 50 dan 100 % pada devoliasi hari ke-45. *Sains Peternakan*, 11(1), 49–55.
- Setyamidjaja, D. 1986. *Pupuk dan Pemupukan*. CV. Simplex, Jakarta.
- Shwarpshakka, S. Y., Dalakon, C. and Kapang, H. 2016. Effect of cutting height and frequency on the forage yield of signal grass (*Brachiaria decumbens*). *International Journal of Science and Applied Science*, 1(1), 47–50.
- Siddique, K. H. M., Belford, R. K. and Tennant, D. 1990. Root:shoot ratios of old and modern, tall and semi-dwarf wheats in a mediterranean environment. *Plant and Soil*, 121(1), 89–98. <https://doi.org/10.1007/BF00013101>
- Sinaga, R. 2008. Air pada rumput gajah dan rumput raja akibat penurunan ketersediaan air. *Jurnal Biologi Sumatera*, 3(1), 29–35.
- Singer, J. W. 2002. Species and Nitrogen Effect on Growth Rate, Tiller Density, and Botanical Composition in Grass Hay Production. *Crop Science*, 42, 208–214. <https://doi.org/10.2135/cropsci2002.2080>
- Singh, N. P. and Singh, R. A. 2002. *Scientific Crop Production*. X Press Graphics, Delhi-28.
- Sinthika, K., Sinniah, J., Sivaneson, S. and Sarmini, N. 2014. Effects of plant spacing on yields and nutritive values of hybrid Napier grass CO-3 in dry zone of Sri Lanka. Retrieved from <http://repo.lib.jfn.ac.lk/ujrr/handle/123456789/693>
- Sirait, J. 2005. *Pertumbuhan dan Serapan Nitrogen Rumput pada Naungan dan Pemupukan Berbeda*. Sekolah Pascasarjana Institut Pertanian Bogor. Retrieved from <http://repository.ipb.ac.id/handle/123456789/8951>
- Sirait, J., Tarigan, A. dan Simanihuruk, K. 2015. Karakteristik morfologi rumput gajah kerdil (*Pennisetum purpureum* cv Mott) pada jarak tanam berbeda di dua agroekosistem di Sumatera Utara. In *Prosiding Seminar Nasional Teknologi Peternakan dan Veteriner* (pp. 643–649).
- Sitompul, S. M. dan Guritno, B. 1995. *Analisis Pertumbuhan Tanaman*. Gadjah Mada University Press, Yogyakarta.
- Smart, A. J., Schacht, W. H. and Moser, L. E. 2001. Predicting leaf/stem ratio and nutritive value in grazed and nongrazed big bluestem. *Agronomy Journal*, 93(6), 1243–1249. <https://doi.org/10.2134/agronj2001.1243>
- Smeekens, S. 2000. Sugar-induced signal transduction in plants. *Annual Review of Plant Physiology and Molecular Biology*, 51, 49–81.

- Smirnoff, N. and Cumbes, Q. J. 1989. Hydroxylradical scavenging activity of compatible solutes. *Phytochemistry*, 28, 1057–1060.
- Soekardi, M., Retno, M. W. dan Hikmatullah, H. 1993. Inventarisasi dan karakterisasi lahan alang-alang. Dalam S. Sukmana, S. Suwardjo, J. Sri Adiningsi, H. Subagyo, H. Suhardho, dan Y. Prawirasumantri (Eds.), *Pemanfaatan Lahan Alang-alang untuk Usaha Tani Berkelanjutan. Prosiding Seminar Lahan Alang-alang* (pp. 1–8). Pusat Penelitian Tanah dan Agroklimat. Badan Litbang Pertanian.
- Soepraptohardjo, M. 1961. “*Tanah Merah*” di Indonesia. Balai Besar Penjelidikan Pertanian, Bogor, Indonesia.
- Somasegaran, P. and Hoben, H. J. (1994). *Handbook for rhizobia: Methods in Legume-Rhizobium Technology*. Springer-Verlag.
- Stanisljević, R., Marković, J., Dinić, B., Lazarević, D., Milenković, J., and An, B. 2009. Yield and chemical composition of Orchard grass harvest remains-straw (*Dactylis glomerata* L.) depending on the vegetation space and application of mineral fertilizers. *Biotechnology in Animal Husbandry*, 25(5–6), 1233–1239.
- Stanley, R. L. and Rhoads, F. M. 2000. Bahiagrass production, nutrient uptake, and soil-test P and K. *Proceedings - Soil and Crop Science Society of Florida*, 59, 159–163.
- Stanton, D., Grombacher, A. W., Pinnisch, R., Mason, H. and Spaner, D. 2007. Hybrid and population density affect yield and quality of silage maize in central Alberta. *Canadian Journal of Plant Science*, 87(4), 867–871. <https://doi.org/10.4141/CJPS06024>
- Stoskopf, N. 1981. *Understanding Crop Production*. Reston Publishing Company, Inc. Virginia.
- Stür, W. ., Shelton, H. and Gutteridge, R. 1994. Defoliation management of forage tree legumes. In R. . Gutteridge and H. . Shelton (Eds.), *Forage tree legumes in tropical agriculture* (p. 168=167). CAB International, UK.
- Subagyo, H., Suharta, N. dan Siswanto, A. B. 2004. Tanah-tanah pertanian di Indonesia. In A. Adimihardja, L. I. Amen, F. Agus, dan D. Djaenudin (Eds.), *Sumberdaya Lahan Indonesia dan Pengelolaannya* (pp. 21–66). Bogor: Pusat Penelitian dan Pengembangan Tanah dan Agroklimat.
- Sudirman. 2013. *Evaluasi Pakan Tropis: Dari Konsep ke Aplikasi (Metode in-vitro Feses)*. Pustaka Reka Cipta, Bandung.
- Susanti, S. 2007. Produksi dan kecernaan in vitro rumput gajah pada berbagaiimbangan pupuk nitrogen dan sulfur. *Buana Sains*, 7(2), 151–156.
- Syahputra, E., Fauzi, F. dan Razali, R. 2015. Karakteristik sifat kimia sub grup

- tanah ultisol di beberapa wilayah Sumatera Utara. *Jurnal Agroekoteknologi*, 4(1), 1796–1803.
- Taiz, L. and Zeiger, E. 2002. *Plant Physiology*. Sinauer Associates Inc., Publishers Sunderland, Massachusetts U.S.A. <https://doi.org/10.1104/pp.900074>
- Temu, V., Rude, B. and Baldwin, B. 2014. Nutritive value response of native warm-season forage grasses to harvest intervals and durations in mixed stands. *Plants*, 3(2), 266–283. <https://doi.org/10.3390/plants3020266>
- Tezara, W., Mitchell, V., P., D. S. and Lawlor, D. W. 2002. Effects of water deficit and its interaction with CO<sub>2</sub> supply on the biochemistry and physiology of photosynthesis in sunflower. *J. Exp. Bot.*, 375(53), 1781–1791.
- Thorvaldsson, G., Tremblay, G. F. and Tapani Kunelius, H. 2007. The effects of growth temperature on digestibility and fibre concentration of seven temperate grass species. *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science*, 57(4), 322–328. <https://doi.org/10.1080/09064710600984221>
- Tilley, J. M. A. and Terry, R. 1963. A two-stage technique for the in vitro digestion of forage crops. *J. Brit. Grassl. Soc.*, 18(104–111).
- Tillman, D. 1988. On the meaning of competition and the mechanisms of competitive superiority. *Functional Ecology*, 1, 304–315.
- Timpong-Jones, E. C., Adjorlolo, L. K. and Ayizanga, R. A. 2015. The impact of harvest frequency on herbage yield and quality of cynodon nlemfuensis. *West African Journal of Applied Ecology*, 23(2), 7–15.
- Tjitrosoedirjo, S. 1990. *Pennisetum polystachyon* (L.) Schult. Weed Info Sheet. SEAWIC SEAMEO BIOTROP. Bogor, Indonesia.
- Tomlinson, K. W. and O'Connor, T. G. 2004. Control of tiller recruitment in bunchgrasses: Uniting physiology and ecology. *Functional Ecology*, 18(4), 489–496. <https://doi.org/10.1111/j.0269-8463.2004.00873.x>
- Torales, A. T. A., Acosta, G. L., Dereibus, V. A. and Moauro, P. M. 2000. Effects of grazing frequency on the production, nutritive value, herbage utilisation, and structure of a *Paspalum dilatatum* sward. *New Zealand Journal of Agricultural Research*, 43(March), 467–472. <https://doi.org/10.1080/00288233.2000.9513443>
- Tremmel, D. C. and Bazzaz, F. A. 1993. How Neighbor Canopy Architecture Affects Target Plant Performance. *Ecology*, 74(7), 2114–2124. <https://doi.org/10.2307/1940856>
- Trenholm, L. E., Dudeck, A. E., Sartain, J. B. and Cisar, J. L. 1998. Bermudagrass growth, total nonstructural carbohydrate concentration, and quality as influenced by nitrogen and potassium. *Crop Science*, 38(1), 168–174.

<https://doi.org/10.2135/cropsci1998.0011183X003800010028x>

- Trenton, F. S. and Joseph, G. L. 2005. Corn stalk response to plant population and the Bt–European corn borer trait. *Agron J.*, 97, 1129–35.
- Troelsen, J. E. 1969. Quality of hay and roughage. In *Proc. Can. Forage Crops Symp.* (pp. 1–18).
- Turner, L. R., Donaghy, D. J., Lane, P. A. and Rawnsley, R. P. 2006. Effect of defoliation interval on water-soluble carbohydrate and nitrogen energy reserves, regrowth of leaves and roots, and tiller number of cocksfoot (*Dactylis glomerata* L.) plants. *Australian Journal of Agricultural Research*, 57(2), 243–249. <https://doi.org/10.1071/AR05130>
- Twidwell, E. K., Johnson, K. D., Cherney, J. H. and Volenec, J. J. 1988. Forage quality and digestion kinetics of Switchgrass herbage and morphological components. *Crop Science*, 28, 778–782. <https://doi.org/10.2135/cropsci1988.0011183X002800050011x>
- Ullah, M. A., Anwar, M. and Rana, A. S. 2010. Effect of nitrogen fertilization and harvesting intervals on the yield and forage quality of elephant grass (*Pennisetum purpureum*) under mesic climate of Pothowar plateau. *Pak. J. Agri. Sci.*, 47(3), 231–234.
- Umami N., M. P. Dewi , B. Suhartanto, C. T., Noviandi, B. Suwignyo, N. Suseno, G. Ishigaki, R. and Akashi. 2015. Growth and productivity of *Brachiaria brizantha* cv MG 5 under the effect of different dose of NPK fertilization. In *The 6th ISTAP International Seminar on Tropical Animal Production* (pp. 978–979).
- Valentin, K. M., Aliou, S. and Augustin, S. B. 2014. Response to fertilizer of native grasses (*Pennisetum polystachion* and *Setaria sphacelata*) and legume (*Tephrosia pedicellata*) of savannah in Sudanian Benin. *Agriculture, Forestry and Fisheries*, 13(3), 142–146. <https://doi.org/10.11648/j.aff.20140303.11>
- Van Man, N. and Wiktorsson, H. 2003. Forage yield , nutritive value , feed intake and digestibility of three grass species as affected by harvest frequency. *Tropical Grasslands*, 37, 101–110.
- van Soest, P. J. 1977. Plant Fiber and Its Role in Herbivore Nutrition. *The Cornell Veterinarian*, 67(3), 307–326.
- van Soest, P. J. 1994. *Nutritional Ecology of the Ruminant*. Cornell University Press.
- Venuto, B. C., Redfearn, D. D., Pitman, W. D. and Alison, M. W. 2004. Impact of seeding rate on annual ryegrass performance. *Grass and Forage Science*, 59(1), 8–14. <https://doi.org/10.1111/j.1365-2494.2004.00397.x>

- Verma, K. C. 2009. *Influence of cutting interval and intercropping on growth, yield and quality of bajra napier hibryd (Pennisetum purpureum x Pennisetum americanum)*. rcharya N.G. Rangan Agricultural University, Hyderabad.
- Vicente-Chandler, J. 1964. *The Intensive Management of Tropical Forages in Puerto Rico*. University of Puerto Rico.
- Villiers, C. J. De. 2007. *The effect of Phosphorus on the growth, plant mineral content and essential oil composition of Buchu (Agathosma betulina)*. Stellenbosch University, South Africa.
- Virkajarvi, P., Pakarinen, K., Hyrkas, M., Seppanen, M. and Bélanger, G. 2012. Tiller characteristics of timothy and tall fescue in relation to herbage mass accumulation. *Crop Science*, 52(2), 970–980. <https://doi.org/10.2135/cropsci2011.01.0039>
- Volenec, J. J. and Nelson, C. J. 1983. Responses of tall fescue leaf meristems to N fertilization and harvest frequency. *Crop Science*, 23(4), 720. <https://doi.org/10.2135/cropsci1983.0011183X002300040028x>
- Vučković, S., A. Simić, B. Ćupina, I. S. and Stanisavljević, R. 2003. The effect of vegetation area size on grass seed yield. *Journal of Agricultural Sciences*, 48(1), 125–134.
- Wang, X., Zhao, L., Yan, B., Shi, L., Liu, G. and He, Y. 2016. Morphological and physiological responses of Heteropogon contortus to drought stress in a dry - hot valley. *Botanical Studies*, 57(17). <https://doi.org/10.1186/s40529-016-0131-0>
- Wangchuk, K., Rai, K., Nirola, H., Dendup, C. and Mongar, D. 2015. Forage growth, yield and quality responses of Napier hybrid grass cultivars to three cutting intervals in the Himalayan foothills. *Tropical Grasslands - Forrajes Tropicales*, 3(3), 142. [https://doi.org/10.17138/TGFT\(3\)142-150](https://doi.org/10.17138/TGFT(3)142-150)
- West, L. T., Beinroth, F. H., Sumner, M. E. and Kang, B. T. 1997. Ultisols: Characteristics and Impacts on Society. In D. Sparks (Ed.), *Advances in Agronomy, Volume 63* (1st ed.). Academic Press.
- Wijitphan, S., Lorwilai, P. and Arkaseang, C. 2009. Effect of cutting heights on productivity and quality of King Grass (*Pennisetum purpureum* cv. King Grass) under irrigation. *Pakistan Journal of Nutrition*, 8(8), 1244–1250.
- Wijitphan, S., Porncha, L. and Chutipongi, A. 2009. Effects of plants spacing on yields and nutritive values of Napier grass (*Pennisetum purpureum* Schum.) under intensive management of nitrogen fertilizer and irrigation. *Pakistan Journal of Nutrition*, 8(8), 1240–1243.
- Wilman, D. and Asiegbu, J. 1982. The effects of variety, cutting interval and nitrogen application on the morphology and development of stolons and

- leaves of white clover. *Grass and Forage Science*, 37, 15–27. <https://doi.org/doi:10.1111/j.1365-2494.1982.tb01572.x>
- Wilson, J. B. 1988. Shoot competition and root competition. *Journal of Applied Ecology*, 25(1), 279–296. <https://doi.org/10.2307/2403626>
- Wilson, J. R., Anderson, K. L. and Hacker, J. B. 1989. Dry matter digestibility in vitro of leaf and stem of buffel grass (*Cenchrus ciliaris*) and related species and its relation to plant morphology and anatomy. *Australian Journal of Agricultural Research*, 40(2), 281–291. Retrieved from <https://doi.org/10.1071/AR9890281>
- Wilson, S. D. and Tilman, D. 1993. Plant competition and resource availability in response to disturbance and fertilization. *Ecology*, 74(2), 599–611. <https://doi.org/10.2307/1939319>
- Xiong, L., Wang, R.-G., Mao, G. and Koczan, J. M. 2006. Identification of drought tolerance determinants by genetic analysis of root response to drought stress and abscisic Acid. *Plant Physiology*, 142(3), 1065–74. <https://doi.org/10.1104/pp.106.084632>
- Yancey, P. H., Clark, M. E., Hand, S. C., Bowlus, R. D. and Somero, G. N. 1982. Living with water stress: evolution of osmolyte systems. *Science*, 217(4566), 1214–1222. Retrieved from <http://science.sciencemag.org/content/217/4566/1214.abstract>
- Yang, C. W. and Kao, C. 1999. Importance of ornithine- $\alpha$ -aminotransferase to proline accumulation caused by water stress in detached rice leaves. *Plant Growth Reg.*, 27, 189–192.
- Yasin, M., Malik, M. A. and Nazir, M. S. 2003. Effects of different spatial arrangements on forage yield, yield components and quality of Mott Elephantgrass. *Pak. J. Agri. Sci*, 2(1), 52–58.
- Yoshiba, Y., Kiyosue, T., Nakashima, K. and Yamaguchi-Shinozaki, K. Shinozaki, K. 1997. Regulation of levels of proline as an osmolyte in plants under water stress. *Plant Cell Physiol.*, 38, 1095–1102.
- Young, J. W. H., Ng, Y. F., Tan, S. N. and Chew, A. Y. L. 2010. Effect of fertilizer application on photosynthesis and oil yield of *Jatropha curcas* L. *Photosynthetica*, 48(2), 208–218.
- Zewdu, T. 2008. Effect of plant density on morphological characteristics, yield and chemical composition of Napier grass (*Pennisetum purpureum* Schumach.). *East African Journal of Sciences*, 2(1), 55–61. <https://doi.org/10.4314/eajsci.v2i1.40365>
- Zewdu, T., Baars, R. and Yami, A. 2003. Effect of plant height at cutting and fertilizer on growth of Napier grass (*Pennisetum purpureum*). *Tropical Science*, 43(1), 57–61. <https://doi.org/10.1002/ts.90>

Zhao, T. J., Sun, S., Liu, Y., Liu, J. M., Liu, Q., Yan, Y. Bin and Zhou, H. M. 2006. Regulating the drought-responsive element (DRE)-mediated signaling pathway by synergic functions of trans-active and trans-inactive DRE binding factors in *Brassica napus*. *Journal of Biological Chemistry*, 281(16), 10752–10759. <https://doi.org/10.1074/jbc.M510535200>

