

DAFTAR PUSTAKA

1. Kawasaki T. Update on pediatric sepsis : a review. *J Intensive Care*. 2017;5:1-12.
2. Ruth A, McCracken CE, Fortenberry JD, et al. Pediatric severe sepsis : current trends and outcomes from the pediatric health information systems database. *Pediatr Crit Care Med*. 2014;15:828-38.
3. Weiss SL, Fitzgerald JC, Pappachan J, et al. Global epidemiology of pediatric severe sepsis : the sepsis prevalence, outcomes, and therapies study. *Am J Respir Crit Care Med*. 2015;191:1147-57.
4. Shime N, Kawasaki T, Saito O, et al. Incidence and risk factors for mortality in paediatric severe sepsis : results from the national paediatric intensive care registry in Japan. *Intensive Care Med*. 2012;38:1191-7.
5. Dewi R. Sepsis pada anak : pola kuman dan uji kepekaan. *Maj Kedokt Indon*. 2011;61:101-6. Yuniar I, Dewi R, Pudjiadi A. Epidemiology of pediatric sepsis in pediatric intensive care unit FKUI-RSCM. *Pediatr Indon*. 2010;50:1-8.
6. Somasetia DH. Tatalaksana awal sepsis berat dan syok sepsis anak. Dalam : Tatalaksana awal kegawatan pada bayi dan anak, PKB IDAI cabang Sumatera Utara/ Departemen IKA fakultas kedokteran USU. 2012;38-61.
7. Venet F, Lepape A, Monneret G. Clinical review : flow cytometry perspective in the ICU – from diagnosis of infection to monitoring of injury-induced immune dysfunctions. *Crit Care*. 2011;231:1-9.
8. Allen E, Bakke AC, Purtzer MZ, Deodhar A. Neutrophil CD64 expression : distinguishing acute inflammatory autoimmune disease from systemic infections. *Ann Rheum Dis*. 2002;61:522-25.
9. Chan T, Gu F. Early diagnosis of sepsis using serum biomarkers. *Expert Rev Mol Diagn*. 2011;11:487-96.
10. Khilnani P, Deopujari S, Carcillo J. Recent advances in sepsis and septic shock. *Indian J of Pediatr*. 2008;75:821-30.
11. Marraro GA. Pediatric sepsis : actions to decrease sepsis in children. *Expert Rev Anti Infect Ther*. 2009;7:941-4.
12. Gibot S, Bene MC, Noel R. Combination biomarkers to diagnose sepsis in the critically ill patient. *Am J of Resp and Crit Care Med*. 2012;186:65-71.
13. Pierrekos C, Vincent JL. Sepsis biomarkers : a review. *Crit Care*. 2010;14:1-18.
14. Icardi M, Erickson Y, Kilborn S, Stewart B, Grief B, Schamweber G. CD64 provides simple and predictive testing for detection and monitoring of sepsis and bacterial infection in hospital patients. *J Clin Microbiol*. 2009;47:3914-9.
15. Standage SW, Wong HR. Biomarkers for pediatric sepsis and septic shock. *Expert rev Anti Infect Ther*. 2011;9:71-9.
16. Schiff DE, Rae J, Martin TR, Davis BH, Curnutte JT. Increased phagocyte FcγRI expression and improved Fcγ-receptor-mediated phagocytosis after in vivo recombinant human interferon-γ treatment of normal human subjects. *Blood*. 1997;90:3187-94.
17. Davis BH. Improved diagnostic approaches to infection/sepsis detection. *Expert Rev Mol Diagn*. 2005;5:193-207.
18. Livaditi O, Kotanidou A, Psarra A. Neutrophil CD64 expression and serum IL-8: sensitive early markers of severity and outcome in sepsis. *Cytokine*. 2006;36:283-90.
19. Davis B. Improved diagnostic approaches to infection/sepsis detection. *Expert Rev Mol Diagn*. 2005;5:193-207.
20. Hoffman JJ. Neutrophil CD64: a diagnostic marker for infection and sepsis. *Clin Chem Lab Med*. 2009;47:903-16.
21. Groselj-Grenj M, Ihan A, Derganc M. Neutrophil and monocyte CD64 and CD163 expression in critically ill neonates and children with sepsis : comparison of fluorescence intensities and calculated indexes. *Hindawi Publishing Corp Mediat of Inflamm*. 2008;1:1-6.
22. Stubljar D, Skvarc M. Expression of CD64 on neutrophils can be used to predict the severity of bloodstream infection before broad range 16S rRNA PCR. *Folia Microbiol*. 2014;1:1-8.

23. Cid J, Aguinaco R, Sanchez R, Garcia-Pardo G, Liorente A. Neutrophil CD64 expression as marker of bacterial infection : a systematic review and meta analysis. *J of Infect.* 2010;60(5):313-9.
24. Li S, Huang X, Chen Z. Neutrophil CD64 expression as a biomarker in the early diagnosis of bacterial infection: a meta-analysis. *Int J of Infect Dis.* 2013;17:e12-23.
25. De Jong E, De Lange DW, Beishuizen A, Van De Ven PM, Girbes ARJ, Huisman A. Neutrophil CD64 expression as a longitudinal biomarker for severe disease and acute infection in critically ill patients. *Int Jnl Lab Hem.* 2016;1:1-9.
26. Djordjevic D, Pejovic J, Surbatovic M, Jevdjic J, Radakovic S, Veljovic M, et al. Prognostic value and daily trend of interleukin-6, neutrophil CD64 expression, C-reactive protein and lipopolysaccharide-binding protein in critically ill patients : reliable predictors of outcome or not? *J Med Biochem.* 2015;34:431-9.
27. Goldstein B, Giroir, Randolph A. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 2005;6:1-7.
28. Enrione MA, Powell KR. Sepsis, septic shock, and systemic inflammatory response syndrome. Dalam : Kliegman RM, Behrman RE, Jenson HB, Stanton BF, penyunting. *Nelson textbook of pediatrics.* Edisi ke-18. Philadelphia: WB Saunders; 2007. p. 1094-9.
29. Latief A, Chairulfatah A, Alam A, Pudjiadi AH, Malisie RF, Hadinegoro SR. Diagnosis dan tata laksana sepsis pada anak. *UKK ERIA dan UKK Infeksi dan Penyakit Tropik IDAI.* 2016
30. Singer M, Deutschman CS, Seymour CW, Hari MS, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016;315(8):801-810.
31. Bryce J, Boschi-Pinto C, Shibuya K, Black RE; WHO Child Health Epidemiology Reference Group. WHO estimates of the causes of death in children. *Lancet.* 2005;365:1147-52.
32. Yuniar I, Dewi R, Pudjiadi A. Epidemiology of pediatric sepsis in pediatric intensive care unit FKUI-RSCM. *Pediatr Indon.* 2010;50(2):1-8.
33. Wolfler A, Silvani P, Musicco M, Antonelli M, Salvo I. Incidence of and mortality due to sepsis, severe sepsis and septic shock in Italian Pediatric Intensive Care Units: a prospective national survey. *Intensive Care Med.* 2008;34:1690-7.
34. Souza D, Shieh H, Barreira E, Ventura A, Bousso A, Troster E. Epidemiology of sepsis in children admitted to PICUs in south America. *Pediatr Crit Care Med.* 2016;17:727-34.
35. Thomas NJ, Tamburro RF, Hall MW, Rajasekaran S, Venglarcik JS. Bacterial sepsis and mechanisms of microbial pathogenesis. Dalam : Nichols DG, penyunting. *Roger's Textbook of Pediatric Intensive Care.* Edisi ke-4. Baltimore: Lippincott Williams & Wilkins; 2008. p. 477-88.
36. Bhan C, Dipankar P, Chakraborty P, Sarangi P. Role of cellular events in the pathophysiology of sepsis. *Inflamm Res.* 2016;1:1-17.
37. Bone RC. A Continuing evolution in our understanding of the Systemic Inflammatory Response Syndromes (SIRS) and the Multiple Organ Dysfunction Syndromes (MODS). *Ann Intern Med.* 1996;125:80-7.
38. Wynn JL, Wong HR. Pathophysiology and treatment of septic shock in neonates. *Clin Perinatol.* 2010;37:439-79.
39. Carcillo JA. Pediatric septic shock and Multiple Organ Failure. *Crit Care Clin.* 2003;19:413-40.
40. Adib-Conquy M, Cavaillon JM. Stress molecules in sepsis and SIRS. *FEBS Letters.* 2007;581:3723-33.
41. Castellheim A, Brekke O-L, Espevik T, Harboe M, Mollnes T. Innate immune responses to danger signals in Systemic Inflammatory Response Syndrome and sepsis. *J of Immunol.* 2009;69:479-91.
42. Short MA. Linking the sepsis triad of inflammation, coagulation, and suppressed fibrinolysis to infants. *Adv Neonatal Care.* 2004;5:258-73.

43. Wang KS, Ford HR, Upperman JS. Metabolic response to stress in the neonate who has surgery. *Neoreviews*. 2006;7:e410-7.
44. Russell JA. Management of sepsis. *NEJM*. 2006;355:1699-713.
45. Glauser MP. Pathophysiology basis of sepsis: Considerations for future strategies of intervention. *J Crit Care Med*. 2000;28:4-8.
46. Wong HR, Cvijanovich N, Wheeler DS, et al. Interleukin-8 as a Stratification Tool for Interventional Trials Involving Pediatric Septic Shock. *Am J Respir Crit Care Med*. 2008;1-3.
47. Estenssoro E, Gonzalez F, Laffaire E, et al. Shock on admission day is the best predictor of prolonged mechanical ventilation in the ICU. *Chest*. 2005;127:598-603.
48. Millar BC, Jiru X, More JE, Earle JAP. A simple and sensitive method to extract bacterial, yeast and fungal DNA from blood culture material. *J of microbiol method* 2000;42:139-47.
49. BATTERY JP. Blood cultures in newborns and children: Optimizing an every day test. *Arct Dist Child Fetal Neonatal*. 2002;87:25-8.
50. Gopi A, Ravikumar KL, Ambarish MG, Shwethalatha NN, Shree SK, Ashwini KV, et al. Time to positivity of microorganism with BACTEC 9050 : an 18 month study among children of 28 days to 60 months in an South Indian tertiary hospital. *Int J of Microbiol Res*. 2011;2(1):12-7.
51. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign : international guidelines for management of sepsis and septic shock : 2016. *Crit Care Med*. 2017;3:00-00.
52. Mejorada GS, Rosales C. Signal transduction by immunoglobulin Fc receptors. *J Leukoc Biol*. 1998;63:521-33.
53. Wen R, Jou ST, Chen Y, Hoffmeyer A, Wang D. Phospholipase C γ 2 is essential for specific function of Fc ϵ R and Fc γ R. *J Immunol*. 2002;169:6743-6752.
54. Lu J, Ellsworth JL, Hamacher N, Oak SW, Sun PD. Crystal structure of Fc γ RI and its implication in high affinity γ -immunoglobulin binding. *The J of Biol Chemistry*. 2011;286:40608-13.
55. Lilius EM, Nuutila J. Bacterial infection, DNA virus infection, and RNA virus infections manifest differently in neutrophil receptor expression. *The Scient World J*. 2012;1-7.
56. Boumazos S, Woolf JM, Dransfield I. Functional and clinical consequences of Fc receptor polymorphic and copy number variants. *Clin and Exper Immunol*. 2009;157:244-54.
57. Lee WL, Harrison RE, Grinstein S. Phagocytosis by neutrophils. *Microbe and Infection*. 2003;5:1299-306.
58. Brown KA, Brain SD, Pearson JD, Edgeworth JD, Lewis SM, Treacher DF. Neutrophils in development of multiple organ failure in sepsis. *Lancet*. 2006;368:157-69.
59. Faix JD. Established and novel biomarkers of sepsis. *Biomarkers Med*. 2011;5:117-30.
60. Grey D, Sack U, Scholz M, Knaack H, Fricke S, Ooppel C, et al. Increased CD64 expression on polymorphonuclear neutrophils indicates infectious complications following solid organ transplantation. *Cytometry Part A*. 2011;79:446-60.
61. Dhlamini MB, Suchard MS, Wiggill TM, Fadahun OO, Ballot DE. Neutrophil CD64 has a high negative predictive value for exclusion of neonatal sepsis. *South African J of Child Health*. 2013;7:25-9.
62. Biron BM, Ayala A, Lomas-Neira JL. Biomarkers for sepsis : what is and what might be? *Biomarker Insights*. 2015;10(54):7-17.
63. Hajian-Tilaki K. Sample size estimation in diagnostic test studies of biomedical informatics. *J of Biomed Informatics*. 2014;48:193-204.
64. Buderer NM. Statistical methodology : incorporating the prevalence of disease into sample size calculation for sensitivity and specificity. *Acad Emerg Med*. 1996;3:895-900.
65. Obuchowski NA. Sample size calculation in studies of test accuracy. *Stat Methods Med Res*. 1998;7:371-92.
66. Souza D, Shieh HH, Troster EJ. Comparisons in the epidemiology and outcomes of pediatric septic patients admitted to public and private hospitals in Latin America. *Pediatr Crit Care Med*. 2014;15:18-19.

67. Lesur O, Roussy JF, Chagnon F, Gallo-Payet N, Dumaine R, Sarret P, et al. Proven infection-related sepsis induces differential stress response early after ICU admission. *Crit Care*. 2010;14(4):131.
68. Pavare J, Grope I, Gardovska D. Prevalence of systemic inflammatory response syndrome (SIRS) in hospitalized children : a point prevalence study. *BMC Pediatr*. 2009;9:25
69. Carvalho PRA, Feldens L, Seitz EE, Rocha T, Soledade M, Trotta A. Prevalence of systemic inflammatory syndromes at a tertiary pediatric intensive care unit. *J Pediatr*. 2005;81:143-148.
70. Mirrett S, Reller LB, Weinstein MP. Controlled clinical comparison of BacT/ALERT standard aerobic medium with BACTEC standard aerobic medium for culturing blood. *J Clin Microbiol*. 2003;41(6):2391-4.
71. Horvath LL, George BJ, Hospenthal DR. Direct comparison of the BACTEC 9240 and BacT/ALERT 3D automated blood culture systems for candida growth detection. *J Clin Microbiol*. 2004;42(1):115-8.
72. Velasquez S, Matute JD, Gamez LY, Enriquez LE, Gomez ID, Toro F, et al. Characterization of nCD64 expression in neutrophils and levels of s-TREM-1 and HMGB-1 in patients with suspected infection admitted in an emergency department. *Biomedica*. 2013;33:643-52.
73. Shen XF, Cao K, Jiang JP, Guan WX, Du JF. Neutrophil dysregulation during sepsis: an overview and update. *J Cell Mol Med*. 2017;21: 1687-97.
74. Muller AC, Tulleken JE, Zijlstra JG, Sluiter W, Hermans J, Kallenberg CG, et al. Leukocyte activation in sepsis; correlations with disease state and mortality. *Intensive Care Med*. 2000;26:883-92.
75. Danikas DD, Karakantza M, Theodorou GL, Sakellaropoulos GC, Gogos CA. Prognostic value of phagocytic activity of neutrophils and monocytes in sepsis. Correlation to CD64 and CD14 antigen expression. *Clin Exp Immunol*. 2008;154:87-97.
76. Muzlovic I, Ihan A, Stubljur D. CD64 index on neutrophils can diagnose sepsis and predict 30-day survival in subjects after ventilator-associated pneumonia. *J Infect Dev Ctries*. 2016;10:260-8.

