

## DAFTAR PUSTAKA

- Andini P. Penentuan Konsentrasi Minyak Kelapa Sawit dan Waktu Fermentasi Bioplastik Poli (3-hidroksibutirat) menggunakan Bakteri *Bacillus* sp. UAAC 21501. [Skripsi]. Padang: Universitas Andalas; 2016.
- Artika IM, Safitri M. Struktur dan Fungs Subseluler. Bogor: IPB Press; 2010
- Asranudin, Putra SR. Efek penambahan PEG 400 pada plastik PHA yang diproduksi dari *Ralstonia picketti*. [Prosiding Seminar Nasional Kimia]. Padang: Institut teknologi Sepuluh November; 2014.
- Atkinson B, Mavituna, F. Biochemical Engineering And Biotechnology Handbook (2<sup>nd</sup> Ed). New York: M Stockton Press; 1991.
- Brodin. Lignocellulosics as sustainable resources for production of bioplastics-a review. Journal Of Cleaner Production. 2017;162:646-664.
- Borah B, Thakur PS, Nigam JN. The influence of nutritional and environmental conditions on the accumulation of poly-β-hydroxybutyrate in *Bacillusmycoides* RLJ B-017. Journal of Applied Microbiology. 2002;92: 776-783.
- Cappuccino JG, N Sherman. Microbiology A Laboratory Manual (7<sup>th</sup> Ed). Perason Education. San Fransisco: Publishing as Benjamin Cummings; 2005.
- Chee JY, Yoga SS, Lau, NS, Ling SC, Abed, RMM, Kumar S. Bacterially produced polyhydroxyalkanoate (PHA): converting renewable resources into bioplastics. Current Research Technology And Education Topics In Applied Microbiology And Microbial Biotechnology. Spain: Formatex; 2010.
- Chen G, Jiang X. Engineering bacteria for enhanced polyhydroxyalkanoate (PHA) biosynthesis. Synthetic and Systems Biotechnology. 2017;2:192-197.
- Czernieck A, Magon A, Schliesser J, Woodfield BF, Pyda M. Heat capacity of poly(3-Hydroxybutyrate). J Chem Thermodynamics. 2014;73:76-84.
- Dawes EA, Senior PL. The role and regulation of energy reserve polymers in microorganism. Adv. Microb. Physiol. 1973;10:135-267.
- Djamaan A, Majid MIA, Azizan MN. Fermentasi plastik mudah lopus poli (3-hidroksibutirat) dari glukosa sebagai sumber karbon tunggal menggunakan bakteri *Erwinia* sp. USMI-20. Jurnal Biosains. 2002:1 – 13.
- Djamaan A. Penghasilan dan Pencirian P(3HB) dan P(3HB-ko-3HV) dari Berbagai Sumber Karbon oleh *Erwinia* sp USMI-20. [Tesis Doktor Falsafah]. Penang: Universitas Sains Malaysia; 2004.

Djamaan A, Agustien A. Landasan Ilmiah Produksi P(3HB) dari Bahan Dasar Minyak Kelapa Sawit. Laporan penelitian fundamental. Dikti Depdiknas RI; 2006.

Djamaan A. Biosintesis Biopolimer Poli(3-hidroksibutirat) Campuran Minyak Kelapa Sawit dan 2-butanol Sebagai Sumber Karbon. Majalah Farmasi Indonesia. 2011;22(4):315 – 322.

Djamaan A, et al. Kajian Biodegradasi Film Plastik Campuran Polistiren dengan poli (3-hidroksibutirat-ko-3-hidroksivalerat) dalam tanah secara *in-vitro*. Jurnal Farmasi Andalas. 2013;1(1).

Djamaan A, Dewi AP. Metode Produksi Biopolimer dari Minyak Kelapa Sawit, Asam Oleat, dan Glukosa. Padang: Andalas University press; 2014.

Djamaan A. Konsep Produksi Biopolimer P(3HB) dan P(3HB-ko-3HV) secara Fermentasi. Padang: Andalas University Press; 2015.

Djamaan A, Fitriani L, Wangi QA, Asiska PD, Ismed F, Zaini E. Production and characterization of microbial polyesters poly (3-hydroxybutyrate) from oleic acid as sole carbon source. Research J. Of Pharmaceutical, Biological and Chemical Sciences. 2015a;6(1):814-822.

Ditjenbun. Statistik Perkebunan Indonesia. Jakarta: Direktorat Jendral Perkebunan; 2017.

Doi Y. Microbial Polyester. New York: UCH Publ. Inc; 1990.

Fardiaz S. Penuntun Praktek Mikrobiologi. Bogor: IPB; 1989.

Gahlawat G, Soni SK. Valorization of waste glycerol for the production of poly (3-hydroxybutyrate) and poly (3-hydroxybutyrate-co-3-hydroxyvalerate) copolymer by Cupriavidus necator and extraction in a sustainable manner. Bioresource Technology. 2017;243:492-501.

Gandjar IG, Rohman A. Kimia Farmasi Analisis. Yogyakarta: Pustaka Pelajar; 2007.

Gill M. Bioplastics: A better alternative to plastics. IMPACT: International Journal of Research in Applied, Natural and Social Sciences. 2014;6:113-117.

Gonzalez GY, Nungaray J, Córdova J, González R, Koller M, Atlic M, Braunegg G. Biosynthesis and characterization of polyhydroxyalkanoates in the polysaccharide-degrading marine bacterium *saccharophagus degradans* ATCC 43961. Journal Indian Microbiology Biotechnol. 2008;35(6):629-33.

Gouda MK, Swellam AE, Omar SH. Production of phb by a *Bacillus megaterium* strain using sugarcane molasses and corn steep liquor as sole carbon and nitrogen sources. Microbiological Research. 2001;156:201-207.

Hermawan A, Hana E, Wiwiek T. Pengaruh Ekstrak Daun Sirih (Piper Betle L.) Terhadap Pertumbuhan Staphylococcus Aureus Dan Escherichia Coli Dengan Metode Difusi Disk. Surabaya: Universitas Airlangga; 2007.

Hermiati E, Mangunwidjaja D, Sunarti TC, Suparno O, Prasetya B. Pemanfaatan biomassa lignoselulosa ampas tebu untuk produksi bioetanol. Jurnal Litbang Pertanian, IPB. 2010;29(4).

Hrabak O. Industrial production of poly-3-hydroxybutyrate. FEMS Microbiology Review. 1992;103:251-256.

Jay J. Modern Food Microbiology. New York: Springer; 2005.

Kumaravel S, Hema R, Lakshmi R. Production of polyhydroxybutyrate (bioplastic) and its biodegradation by *Pseudomonas lemoignei* and *Aspergillus niger*. E-Journal of Chemistry. 2010;7(1):536-542.

Kosseva MR, Rusbandi E. Review : Trends in the biomanufacture of polyhydroxyalkanoates with focus on downstream processing. International Journal of Biological Macromolecules. 2017.

Krisyanella, Djamaan A, Aulia W. Optimasi proses produksi bioplastik poli (3-Hidroksibutirat) dengan bakteri *Bacillus* sp FAAC 20801 menggunakan bahan dasar jerami padi secara fermentasi. J.I Sains dan Teknologi Farmasi. 2012;17(1):60-72.

Lay B. Analisis Mikroba di Laboratorium. Jakarta: R.T Raja Grafindo Persada; 1994.

Lab Bioteknologi LIPI. Hasil Pengujian 16S rRNA. Bogor: LIPI Pusat Penelitian Bioteknologi; 2016.

Lee SY. Bacterial Polyhydroxyalkanoate Biotechnol. Bio-eng, 1996;99:1-14.

Majid MIA, Djamaan A, Few LL, Agustien A, Toh MS, Samian MR, Najimudin N, Azizan MN. Production of poly(3-hydroxybutyrate) and its copolymer poly(3-hydroxybutyrate-co-3-hydroxyvalerat) by *Erwinia* sp. USMI-20. International Journal of Biological Macromolecules. 1999;25:95-104.

Martínez V, Garcia P, Garcia JL, Prieto MA. Controlled autolysis facilitates the polyhydroxyalkanoate recovery in *Pseudomonas putida* KT2440. Microb. Biotechnol. 2011;4(4):533–547.

Miller JN. Statistic and Chemometrics for Analytical Chemistry (4<sup>th</sup> Ed). Harlow: Prentice Hall; 2000

Mohapatra S, Maity S, Dash HR, Das S, Pattnaik S. *Bacillus* and biopolymer: Prospects and challenges. Biochemistry and Biophysics Reports. 2017;12:206-213.

Nubia M, Ivonne G, Dionisio M, Victoria G, Dolly R, Diego S, Juan G, Fabio A, Armando E, Dolly M. Bioprospecting and characterization of poly-hydroxyalkanoate (PHAs) producing bacteria isolated from colombian sugarcane producing areas. African J. Od Biotechnol. 2007;6(13):1536-1543.

Ostle GA, Holt JG. Nile blue A as a fluorescent stain for poly- $\beta$ -hydroxybutyrate. Applied and Environmental Microbiology. 1982;44(1):238-241.

Paju N, Paulina VYY, Novel K. Uji efektifitas salep ekstrak daun binahong (*Anredera cordifolia* Steenis) pada kelinci (*Oryctolagus cuniculus*) yang terinfeksi bakteri *Staphylococcus aureus*. Jurnal Ilmiah Farmasi-UNSRAT. 2013;2(1):51-61.

Pantazaki AA, Papaneophytou CP, Lambropoulou DA. Simultaneous polyhydroxyalkanoate and rhamnolipid production by *Thermus thermophilus* HBS. AMB Express. 2011;1(17):1-13

Pelczar MJ, Chan ECS. Dasar-Dasar Mikrobiologi. Jakarta: Penerbit Erlangga; 2007.

Pena T, Castillo A, Garcia M, Millan, D Segura. Biotechnological strategies to improve production of microbial poly-(3-Hydroxybutyrate): A Review of Recent Research Work. Microbial Biotechnology. 2014;7:278-293.

Petrasovits LA, Purnell MP, Nielsen LK, Brumbley SM. Production of polyhydroxybutyrate in sugarcane. Plant Biotechnology. 2007:162-172

Raza AR, Abid S, Banat IM. Polyhydroxyalkanoates: Characteristics, production, recent developments and applications. International Biodeterioration & Biodegradation. 2017;126:45-56.

Reddy CSK, Ghai R, Rashmi, Kalia VC. Polyhydroxyalkanoates: an overview. Bioresource Technology. 2003;87:137-146.

Rezwan K, Chen QZ, Blaker JJ, Boccaccini AR. Biodegradable and bioactive porous polymer inorganic composite scaffolds for bone tissue engineering. Biomaterials. 2006;27:3413-3431.

Ruslan R, Andini P, Usni FH, Yulisa N, Silvia A, Irwandi, Dewi AP, Lalfari RS, Djamaan A. Biosynthesis of a polyester poly(3-hydroxybutyrate) by using *Bacillus sp.* UAAC 21501. ICIND. 2017

Stanbury PF, Whitaker A, Hall SJ. Principles Of Fermentation Technology (2<sup>nd</sup> Ed). United Kingdom: Pergamon Ltd, Elsevier Science; 1995.

Stevens ES. Green plastic : an introduction to the new science of biodegradable plastic. New Jersey: Universitas Princeton; 2002.

Sumartono NW, Handayani F, Desiriana R, Novitasari W, Hulfa DS. Sintesis dan karakterisasi bioplastik berbasis alang-alang (*Imperata cylindrica* (L.) dengan penambahan kitosan, gliserol dan asam Oleat. Pelita. 2015;10(2):13-25.

Suprihatin. Teknologi Fermentasi. Surabaya: UNESA University Press; 2010.

Suriyamongkol P, Weslake R, Narine S, Monoley M, Shah S. Biotechnological approaches for the production of polyhydroxyalkanoates in microorganisms and plants-a review. Biotechnology Advance. 2007;25:148-175.

Sutton S. Determination of inoculum for Microbiological Testing. Summer. 2011;15:3

Wardani R. Bahaya Penggunaan Plastik. Pendidikan MIPA. Palangkaraya: Universitas Palangkaraya; 2009.

Wisuthiphaet N dan Napathorn SC. Optimisation of the use of products from the cane sugar industry for poly (3-hydroxybutyrate) production by *Azohydromonas lata* DSM 1123 in fed-batch cultivation. Process Biochemistry. 2016;51:352-361.

Yuli D, Chici A, Sri ID. Sintesa Bioplastik dari Pati Pisang dan Gelatin dengan Plasticizer Gliserol. Seminar Nasional Sains dan Teknologi-II. Lampung: Universitas Lampung; 2008.

Yulialdi Y. Isolasi dan karakterisasi bakteri termofilik penghasil bioplastik poli-(3-Hidroksibutirat) dari sumber air panas Bukit Gadang Kabupaten Solok, Sumatera Barat. [Skripsi]. Padang: Universitas Andalas; 2015.