

DAFTAR PUSTAKA

- Abdel Baset, et al. (2014). Validation of the Risk of Malignancy Index in primary evaluation of ovarian masses. *Middle East Fertility Society Journal* , 324–328.
- Ahmad S, et al. (2015). The characteristic ultrasound features of specific types of ovarian pathology (Review). *Int J Oncol.* 46(2): 445–458.
- Akker, P. A. V. D., et al. (2011). External validation of the adapted Risk of Malignancy Index incorporating tumor size in the preoperative evaluation of adnexal masses. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 159, 422-425.
- Akturk, E., et al. (2011). Comparison of four malignancy risk indices in the detection of malignant ovarian masses. *Journal of Gynaecologic Oncology*, 22, 177-182.
- Ali Yavuzcan, et al. (2013). Should Cut-Off Values of the Risk of Malignancy Index be Changed for Evaluation of Adnexal Masses in Asian and Pacific Populations?. *Asian Pac J Cancer Prev*, 14 (9), 5455-5459
- Alka Prakash, et al. (2004). The management of ovarian cysts in premenopausal women. *Royal College of Obstetricians and Gynaecologists*, 6:1.
- Ashrafgangooei, T. et al. (2011). Risk of Malignancy Index in Preoperative Evaluation of Pelvic Masses. *Asian Pasific Journal of Cancer Prevention*, 12, 1727-1730.
- Bernardin L, et al. (2012). Effectiveness of semi-quantitative multiphase dynamic contrast-enhanced MRI as a predictor of malignancy in complex adnexal masses: radiological and pathological correlation. *Eur Radiol.* 22(4):880-90.
- Bouzari, Z., et al. (2011). Comparison of three malignancy risk indices and CA-125 in the preoperative evaluation of patients with pelvic masses. *BMC Research Note*, 4,206.
- Brett M. Reid, et al. (2017). Epidemiology of ovarian cancer: a review. *Cancer Biol Med* 2017.

- Budiana, I. N. (2013). Tumor Ovarium: Prediksi Keganasan Prabedah. *Jurnal Ilmiah Kedokteran*, 44 (Divisi Onkologi-Ginekologi, Departemen Obstetri dan Ginekologi FK Unud/RSUP Sanglah Denpasar), 179-85.
- Charuwan Tantipalakor, et al. (2014). IOTA Simple Rules in Differentiating between Benign and Malignant Ovarian Tumor. *Asian Pacific Journal of Cancer Prevention*, Vol 15, 5123-5126
- Cristina Anton, et al. (2012). A comparison of CA125, HE4, risk ovarian malignancy algorithm (ROMA), and risk malignancy index (RMI) for the classification of ovarian masses. *Clinical Science* (Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Obstetrics and Gynecology, São Paulo/SP, Brazil), 437-41.
- Dahlan, M. Sopiudin. 2009. Penelitian Diagnostik, dasar-dasar teoritis dan aplikasi dengan program SPSS dan Stata. Seri evidence based medicine 5. Jakarta; Salemba medika.
- Dahlan, M. Sopiudin. 2014. Statistik untuk Kedokteran dan Kesehatan. . Seri 1. Edisi 6. Jakarta; Epidemiologi Indonesia
- Daniel W. Chan, et al. (2009). *Use of Tumor Markers in Testicular, Prostate, Colorectal, Breast, and Ovarian Cancers*. Washington, United State: the American Association for Clinical Chemistry, Inc.
- Danijela Jelovac and Deborah K. Armstrong. (2011). Recent Progress in the Diagnosis and Treatment of Ovarian Cancer. *CA Cancer J Clin*. 2011 ; 61(3): 183–203.
- Deepti Vijay Mankar, et al. (2015). *Histopathological profile of ovarian tumours: A twelve year institutional experience*. MJMSR, Vol 6 (2): 107-111.
- Dirk Timmerman, et al. (2000). Terms, definitions and measurements to describe the sonographic features of adnexal tumors: a consensus opinion from the International Ovarian Tumor Analysis (IOTA) group.. *Ultrasound Obstet Gynecol* 2000; 16: 500-505.
- Dirk Timmerman, et al. (2008). Simple ultrasound-based rules for the diagnosis of ovarian cancer. *Ultrasound Obstet Gynecology*; 31, 681–690
- Dirk Timmerman, et al. (2010). Simple Ultrasound Rules to Distinguish Between Benign and Malignant Adnexal Masses Before Surgery : Prospective Validation by IOTA Group. *British Medical Journal*, 1-8

- Dirk Timmerman, et al. (2010). Terms, definitions and measurements to describe the sonographic features of adnexal tumors: a consensus opinion from the International Ovarian Tumor Analysis (IOTA) group. *Ultrasound in Obstetrics and Gynecology*, 500-505.
- Dirk Timmerman, et al. (2016). Predicting the risk of malignancy in adnexal masses based on the Simple Rules from the International Ovarian Tumor Analysis group. *American Journal of Obstetrics*, 425-437
- Dominique Trudel, et al. (2012). Human epididymis protein 4 (HE4) and ovarian cancer prognosis. *Gynecologic Oncology* , 511–515.
- Fang Shen, et al. (2017). The prevalence of malignant and borderline ovarian cancer in pre- and post-menopausal Chinese women. *Oncotarget*. 2017 Oct 6; 8(46): 80589–80594.
- Fathallah K, et al. (2011). [External validation of simple ultrasound rules of Timmerman on 122 ovarian tumors]. *Gynecologie, Obstetrique & Fertilité*. 39(9):477-481
- Froyman W, et al. (2017). Performance of the RMI and IOTA ADNEX and Simple Rules risk model in the evaluation of adnexal masses not classifiable using the Easy Descriptors as first step. *Ultrasound in Obstetrics & Gynecology*; Volume 50, Issue S1
- Greg L. Perkins, et al. (2003). Serum Tumor Markers. *American Family Physician* , 1075-82.
- Hariyono Winarto, et al. (2014). Modification of Cutoff Values for HE4, CA125, the Risk of Malignancy Index, and the Risk of Malignancy Algorithm for Ovarian Cancer Detection in Jakarta, Indonesia. *Asian Pacific journal of cancer prevention: APJCP*.
- Ignace Vergote, et al. (2016). Ovarian Cancer: Diagnosis, Treatment and Follow-Up. *Good Clinical Practice. KCE Report*.
- Insin Putsarat and Prueksaritanond Nisa. (2013). Evaluation of Four Risk of Malignancy Indices (RMI) in the Preoperative Diagnosis of Ovarian Malignancy at Rajavithi Hospital. *Thai Journal of Obstetrics and Gynaecology* , 163-175.

Iwani Rahmah Rambe, et al. (2014). Profil Tumor Ganas Ovarium di Laboratorium Patologi Anatomi Fakultas Kedokteran Universitas Andalas Periode Januari 2011 Sampai Desember 2012. FK UNAND.

Jacques S. Abramowicz, et al. (2017). Ovarian mass differentiating benign from malignant: the value of the International Ovarian Tumor Analysis ultrasound rules. *American Journal of Obstetrics*, 2-9

Jeroen Kaijser, et al. (2013). A comparison between an ultrasound based prediction model (LR2) and the Risk of Ovarian Malignancy Algorithm (ROMA) to assess the risk of malignancy in women with an adnexal mass. *Gynecologic Oncology* 129, 377–383

Jing Bian, et al. (2013). Clinical Significance of Combined Detection of Serum Tumor Markers in Diagnosis of Patients with Ovarian Cancer . *Asian Pac J Cancer Prev*, 14 (11), 6241-6243

Jonathan S. Berek, et al. (2012). Berek & Novak's Gynecology 15th edition. Wolter Kluwer Health: Lippincott Williams & Wilkins.

José Carlos Campos Torres, et al. (2002). Risk-of-Malignancy Index in preoperative evaluation of clinically restricted ovarian cancer. *São Paulo Medical Journal* , 73-75.

Jung-Woo Park, et al. (2012). Four Risk of Malignancy Indices in Evaluation of Pelvic Masses. *Korean J Obstet Gynecol* , 636-643.

M. Farid Aziz (2009). Gynecological cancer in Indonesia. *Division of Oncology, Department of Obstetrics and Gynecology, University of Indonesia, Jakarta, Indonesia*

Mansur, S dan Sigit P. 2013. *Modified Gatot Score Has a Better Specificity in Predicting Ovarian Malignancies Compared to Risk Malignancy Index*. *Indonesia Obstetri Gynecology Journal* Volume 37 No 2 Tahun 2013. Jakarta.

Marice Sihombing and Anna Maria Sirait. (2007). Angka Ketahanan Hidup Penderita Kanker Ovarium di RS Dr. Cipto Mangunkusumo Jakarta. *Majalah Kedokteran Indonesia* , 345-352.

Milan Terzić, et al. (2011). Evaluation of the risk malignancy index diagnostic value in patients with adnexal masses. *Vojnosanit Pregl* .

- Mofrilindo. (2014). Keakuratan Uji Diagnostik Menggunakan Indeks Risiko Keganasan Dan Indeks Resistensi Neovaskuler Pada Tumor Ovarium. *Tesis*
- Monica R : Epidemiologic and genetic Factor Associated with Ovarian Cancer, *Cancer Nurs* : 2009
- Moore RG, Brown AK, Miller MC, et al. (2008). The use of multiple novel tumorbiomarkers for the detection of ovarian carcinoma in patients with a pelvic mass. *Gynecol Oncol* , 108:402-8.
- Moolthiya Watcharada and Yuenyao Pissamai. (2009). The Risk of Malignancy Index (RMI) in Diagnosis of Ovarian Malignancy. *Asian Pacific Journal of Cancer Prevention* (Department of Obstetrics and Gynecology, Faculty of Medicine, Srinagarind Hospital, Khon Kaen University, Thailand), 865-68.
- Mr Sean Duffy, et al. (2011). Ovarian cancer: the recognition and initial management of ovarian cancer. the National Collaborating Centre for Cancer at Velindre NHS Trust, Cardiff, Wales.
- Mugi Wahidin, et al. (2012). *Population-Based Cancer Registration in Indonesia*. *Asian Pacific J Cancer Prev*, 13, 1709-1710
- Nowak M, et al. (2002).[Ovarian tumors in the reproductive age group]. *Ginekologia Polska*,73(4):354-8.
- Nunes N, et al. (2014). Use of IOTA simple rules for diagnosis of ovarian cancer: meta-analysis. *Ultrasound Obstet Gynecol*. 44(5):503-14.
- Peter Boyle and Bernard Levin. (2008). *World Cancer Report 2008*. WHO. International Agency for Research on Cancer
- Pınar Yörük, et al. (2008). Comparison of the Risk of Malignancy Index and Self-Constructed Logistic Regression Models in Preoperative Evaluation of Adnexal Masses. *J Ultrasound Med* , 1469–1477.
- Prameela, et al. (2015). Role of Risk Malignancy Index in Predicting Ovarian Malignancy and its Correlation with Histopathological Report. *Scholars Journal of Applied Medical Sciences (SJAMS)* , 490-493.
- Rarung, M. (2008). Sensitivitas dan Spesifisitas Petanda Tumor CA 125 sebagai Prediksi Keganasan Ovarium. *JKM* , 9-14.

- Safia Sultana Munir, Misbah Sultana And Dawood Amin. (2010). The Evaluation Of Pelvic Mass. *Biomedica* , 70-75.
- Sayasneh A, et al. (2013). Multicentre external validation of IOTA prediction models and RMI by operators with varied training. *British Journal of Cancer* 108, 2448–2454
- Sharon E. Clarke, et al. (2009). Risk of Malignancy Index in the Evaluation of Patients With Adnexal Masses. *JOGC* (Department of Obstetrics and Gynaecology, Division of Gynaecologic Oncology, Dalhousie University, Halifax NS), 440-5.
- Ulusoy, et al. (2007). The risk of malignancy index in discrimination of adnexal masses. *International Journal of Gynecology and Obstetrics* , 186–191.
- Wouter Froyman, et al. (2017). Validation of the Performance of International Ovarian Tumor Analysis (IOTA) Methods in the Diagnosis of Early Stage Ovarian Cancer in a Non-Screening Population. Licensee MDPI, Basel, Switzerland, 1-11
- Yamamoto Y, et al. (2015). Comparison of 4 Risk-of-Malignancy Indexes in the Preoperative Evaluation of Patients With Pelvic Masses: A Prospective Study. *Clinical Ovarian and Other Gynecologic Cancer* , 8-12.
- Yasmin Jaffar, et al. (2010). Clinical Presentation of Ovarian Tumors. *Journal of Surgery Pakistan (International)* 18 (2).
- Yazbek J, et al. (2007). Accuracy of ultrasound subjective 'pattern recognition' for the diagnosis of borderline ovarian tumors. *Ultrasound Obstet Gynecol.* 29(5):489-95.
- Yogambal, M, et al. (2014). Ovarian tumours- Incidence and distribution in a tertiary referral center in south India. *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)* e-ISSN: 2279-0853, p-ISSN: 2279-0861. Volume 13, Issue 2 Ver. III. (Feb. 2014), PP 74-80
- Zoya Yurkovetsky, et al. (2010). Development of a Multimarker Assay for Early Detection of Ovarian Cancer. *Journal of Clinical Oncology*, 28 (the University of Pittsburgh), 2159-66.