

DAFTAR PUSTAKA

- Akolekar, R; de Cruz, J; Foidart, J; Munaut, C; Nicolaides, KH. (2010). Maternal plasma soluble fms-like tyrosine kinase-1 and free vascular endothelial growth factor at 11 to 13 weeks of gestation in preeclampsia. *Prenat Diagn*, 30, 191-197.
- Agarwal, I., & Karumanchi, S. A. (2011). Preeclampsia and the Anti-Angiogenic State. *PubMed Central*, 1, 17-21.
- Bahado-Singh, R. O., Syngelaki, A., Akolekar, R., Mandal, R., Bjondahl, T. C., Han, B., et al. (2015). Validation of metabolomic models for prediction of early-onset preeclampsia. *American Journal of Obstetrics & Gynecology*.
- Baumann, M. U., Bersinger, N. A., Mohaupt, M. G., Raio, L., Gerber, S., & Surbek, D. V. (2008). First-trimester serum levels of soluble endoglin and soluble fms-like tyrosine kinase-1 as first-trimester markers for late-onset preeclampsia. *American Journal of Obstetrics & Gynecology*, 266.e1-266.e6.
- Beňovská, M., Opluštilová, A., Pinkavová, J., Hodická, Z., & Čermáková, Z. (2017). The New Possibilities in Early Diagnosis of Preeclampsia by Soluble fms-Like Tyrosine Kinase-1 and Placental Growth Factor in 16–20 Weeks Gestation. *American Society for Clinical Pathology*.
- Berghella, V. (2012). *Maternal-Fetal Evidence Based Guidelines 2nd Edition*. New York : Informa Healthcare.
- Bukowski, R. (2010). Stillbirth and Fetal Growth Restriction. *Clinical Obstetrics and Gynecology*, 673-680.
- Chandra, B. (2009). *Ilmu Kedokteran Pencegahan dan Komunitas*. Jakarta: EGC.
- Coad, J., & Dunstall, M. (2007). *Anatomi Fisiologi untuk Bidan*. Jakarta: EGC.
- Cunningham, F. G., Leveno, K. J., Bloom, S. L., Spong, C. Y., Dashe, J. S., Hoffman, B. L., et al. (2014). *Williams Obstetrics (24 ed.)*. New York: Mc Graw-Hill Education.
- Dahlan, M. S. (2016). *Besar Sampel dalam Penelitian Kedokteran dan Kesehatan Seri 2 Edisi 4*. Jakarta: Epidemiologi Indonesia.
- Dahlan, M. S. (2016). *Statistik Untuk Kedokteran dan Kesehatan*. Jakarta: Salemba Medika.
- Dinas Kesehatan Kota Padang. (2015). *Profil Kesehatan kota Padang Tahun 2015*. Padang: Dinas Kesehatan Kota Padang.
- Dinas Kesehatan Provinsi Sumatera Barat. (2016). *Profil Kesehatan Provinsi Sumatera Barat Tahun 2015*. Padang: Dinas Kesehatan Provinsi Sumatera Barat.
- Erez, O; Romero, R; Espinoza, J; Fu, W; Todem, D; Kusanovic, JP; et al. (2008). The change in concentrations of angiogenic and antiangiogenic Factors in maternal plasma between the First and second trimesters in risk assessment for the subsequent development of preeclampsia and SGA. *J Matern Fetal Neonatal Med*; 21(5): 279–287.
- Erez, O., Romero, R., Maymon, E., Chaemsathong, P., Done, B., Pacora, P., et al. (2017). The prediction of late-onset preeclampsia: Results from a longitudinal proteomics study. *PLOS ONE*.

- Espinoza, J., Romero, R., Nien, J. K., Gomez, R., Kusanovic, J. P., Gonçalves, L. F., et al. (2007, April). Identification of Patients at Risk for Early Onset and/or Severe Preeclampsia With the Use of Uterine Artery Doppler Velocimetry and Placental Growth Factor. *Am J Obstet Gynecol*, 196(4), 326.e1–326.13.
- Fachinetti, F., Dekker, G. A., Baronciani, D., & Saade, G. (2010). *Stillbirth: Understanding and Management*. New York: Informa Healthcare.
- Froen, J. F., Cacciatore, J., McClure, E. M., Kuti, O., Jokhio, A. H., Islam, M., et al. (2012). Stillbirths: Why They Matter. Dalam R. Horton, *The Lancet: Global Health Series: Stillbirths*. New York: Elsevier.
- Gibbs, R. S., Karlan, B. Y., Haney, A. F., & Nygaard, I. E. (2008). *Danforth's Obstetrics and Gynecology, 10th Edition*. USA: Lippincott Williams & Wilkins.
- Gibney, M. J., Margetts, B. M., Kearney, J. M., & Arab, L. (2008). *Gizi Kesehatan Masyarakat*. Jakarta: EGC.
- Gomez-Arriaga, P., I Herraiz, E. L.-J., Escribano, D., Denk, B., & Galiando, A. (2014). Uterine artery Doppler and sFlt-1/PIGF ratio: prognostic value in early-onset pre-eclampsia. *Ultrasound Obstet Gynecol*, 43, 525-532.
- Gunardi, J. I., Mose, J., Handono, B., Satari, M. H., Anwar, A. D., Fauziah, P. N., et al. (2015). Difference of concentration of placental soluble fms-like tyrosine kinase-1(sFlt-1), placental growth factor (PIGF), and sFlt-1/ PIGF ratio in severe preeclampsia and normal pregnancy. *BMC Res Notes*, 8, 534.
- Gupte, S. (2016). *The Short Textbook of Pediatrics*. USA: Jaypee Brothers Medical Publishers.
- Guyton, A., & Hall, J. (2016). *Textbook of Medical Physiology 13th Edition*. USA: Elsevier.
- Hidayat, A. A. (2008). *Pengantar Ilmu Kesehatan Anak untuk Pendidikan Bidan*. Jakarta: Salemba Medika.
- Huppertz, B. (2008). Placental Origins of Preeclampsia: Challenging the Current Hypothesis. *Hypertension*, 970-975.
- Jacobs, M., Nassar, N., Roberts, C. L., Hadfield, R., Morris, J. M., & Ashton, A. W. (2011). Levels of soluble fms-like tyrosine kinase one in first trimester and outcomes of pregnancy: a systematic review. *Reproductive Biology and Endocrinology*, 9, 77.
- James, J. S. (2006). The regulation of trophoblast differentiation by oxygen in the first trimester of pregnancy. *Human Reproduction Update*, 137-144.
- Jeyabalan, A. (2013, Oktober). Epidemiology of preeclampsia: Impact of obesity. *NIH Public Access*, 71(1).
- Kementerian Kesehatan Republik Indonesia. (2010). *Buku Saku Pelayanan Kesehatan Neonatal Esensial*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Kementerian Kesehatan Republik Indonesia. (2013). *Riset Kesehatan Dasar Tahun 2013*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Kementerian Kesehatan Republik Indonesia. (2016). *Profil Kesehatan Indonesia Tahun 2015*. Jakarta: Kementerian Kesehatan Republik Indonesia.

- Kliegman, R. M. (2016). *Nelson Textbook of Pediatrics 20th Edition*. Philadelphia: Elsevier.
- Kosim, M. S., Yunanto, A., Dewi, R., Sarosa, G. I., & Usman, A. (2010). *Buku Ajar Neonatologi Edisi Pertama*. Jakarta: Ikatan Dokter Anak Indonesia.
- Kwiatkowski, S., Dołęgowska, B., Ewa, K., Rzepka, R., Natalia Marczuk, B. L., & Torbe, W. M.-B. (2016). A Common Profile of Disordered Angiogenic Factor Production and the Exacerbation of Inflammation in Early Preeclampsia, Late Preeclampsia, and Intrauterine Growth Restriction. *PLoS ONE*.
- Kwiatkowski, S., Dołęgowska, B., Ewa, K., Rzepka, R., Natalia Marczuk, B. L., & Torbe, W. M.-B. (2017). Do the physiological aging of the placenta and the changes in angiogenesis marker sFlt-1 and PlGF concentrations predispose patients to late-onset preeclampsia? *The Journal of Maternal-Fetal & Neonatal Medicine*.
- Luealon P. Phupong V. 2010. Risk Factor of preeclampsia in Thai Woman. Available from: <http://jmat.mat.or.th/index.php/jmat/article/viewfile/302/302>
- Lyll, F., & Belfort, M. (2007). *Pre-eclampsia: Etiology and Clinical Practice*. New York: Cambridge University Press.
- Lynch, A., Murphy, J., Gibbs, R., Levine, R., Giclas, P., Salmon, J., et al. (2010). The interrelationship of complement-activation fragments and angiogenesis-related factors in early pregnancy and their association with pre-eclampsia. *BJOG*, 456-462.
- Muy-Rivera, M; Vadachkoria, S; Woelk, GB, Qiu, C; Mahomed, K; Williams, MA. (2005). Maternal Plasma VEGF, sVEGF-R1, and PlGF Concentrations in Preeclamptic and normotensive Pregnant Zimbabwean Women. *Physiol. Res.* 54: 611-622.
- Nanjo, S., Minami, S., Mizoguchi, M., Yamamoto, M., Yahata, T., Toujima, S., et al. (2017). Levels of serum-circulating angiogenic factors within 1 week prior to delivery are closely related to conditions of pregnant women with pre-eclampsia, gestational hypertension, and/or fetal growth restriction. *The Journal of Obstetrics and Gynaecology Research*.
- Orabona, R., Vizzard, E., Sciatti, E., Prefumo, F., Bonade, I., Valcamonica, A., et al. (2016). Maternal cardiac function after HELLP syndrome: an echocardiographic study. *Ultrasound Obstet Gynecol*.
- Osungbade, K. O., & Ige, O. K. (2011). Public Health Perspectives of Preeclampsia in Developing Countries: Implication for Health System Strengthening. *Journal of Pregnancy*, 2011.
- Parra-Cordero, M; Rodrigo, R; Barja, P; Bosco, C; Rencoret, G; Sepuvela-Martinez, A; et al. (2013). Prediction of early and late pre-eclampsia from maternal characteristics, uterine artery Doppler and markers of vasculogenesis during first trimester of pregnancy. *Ultrasound Obstet Gynecol* 2013; 41: 538-544.
- Perni, U; Sison, C; Sharma, V; Helseth, G; Hawfield, A; Suthanthiran, M; et al. (2012). Angiogenic Factors in Superimposed Preeclampsia A Longitudinal Study of Women With Chronic Hypertension During Pregnancy. *Hypertension*. 2012;59:740-746.
- Pinheiro, CC; Rayol, P; Gozzani, L; dos Reis, LM; Zampieri, G; Dias, CB; et al. (2014). The relationship of angiogenic factors to maternal and neonatal manifestations of early-onset and late-onset preeclampsia. *Prenatal Diagnosis*, 34, 1084-1092.

- Putra, R. J. (2010). *Hubungan faktor biologik terhadap kadar sFlt-1 serum pada wanita hamil dengan preeklamsia berat (PEB)*. Tesis. Padang: Universitas Andalas.
- Rahmi, L., Herman, R. B., & Yusrawati. (2016). Perbedaan rerata kadar Soluble Fms-Like Tyrosine Kinase-1 (Sflt-1) Serum pada Penderita Early Onset, Late Onset Preeklamsia Berat/ Eklamsia dan Kehamilan Normal. *Jurnal Fakultas Kedokteran Universitas Andalas*, 5(1).
- Roberts, J. M., & Bell, M. J. (2013). If we know so much about preeclampsia, why haven't we cured the disease? *J Reprod Immunol*.
- Romero, R., Kusanovic, J. P., Gabor, N., Erez, O., Gotsch, F., Espinoza, J., et al. (2008). First Trimester Maternal Serum PP13 in the Risk Assessment for Preeclampsia. *Am J Obstet Gynecol*.
- Sadler, T. (2015). *Langman's Medical Embryology 13th Edition*. Philadelphia: Wolters Kluwer Health.
- Saifuddin, A. B. (2010). *Ilmu Kebidanan*. Jakarta: Yayasan Bina Pustaka Sarwono Prawirohardjo.
- Sastroasmoro, S., & Ismael, S. (2011). *Dasar-dasar Metodologi Penelitian Klinis Edisi 4*. Jakarta: Sagung Seto.
- SDKI. (2012). *Survei Demografi Dan Kesehatan Indonesia*. Jakarta: Badan Pusat Statistik.
- Singh, V. (2012). *Textbook of Clinical Embryology*. New Delhi: Elsevier.
- Singhai, S. R., Deepika, Anshu, & Nanda, S. (2009, September-Desember). Maternal and Perinatal Outcome in Severe Pre-eclampsia and Eclampsia. *Sout Asian Federation of Obstetrics and Gynecology*, 1(13), 25-28.
- Soto, E., Romeo, R., Kusanovic, J. P., Ogge, G., Hussein, Y., Yeo, L., et al. (2012). Late-Onset Preeclampsia is associated with an Imbalance of Angiogenic and Anti Angiogenic Factors in Patients with and Without Placental Lesions Consistent with Maternal Underperfusions. *Maternal Fetal Neonatal Med*, 498-507.
- Thadhani, R., Hagmann, H., Schaarschmidt, W., Roth, B., Cingoz, T., Karumanchi, S. A., et al. (2016). Removal of Soluble Fms-Like Tyrosine Kinase-1 by Dextran Sulfate Apheresis in Preeclampsia. *Journal American Society of Nephrology*, 27, 903-913.
- Triunfo, S., Crovetto, F., Crispi, F., Rodriguez-Sureda, V., Dominguez, C., Nada, A., et al. (2016). Association of first-trimester angiogenic factors with placental histological findings in late-onset preeclampsia. *Placenta*.
- Villa, PM; Hamalainen, E; Maki, A; Raikkonen, K; Pesonen, AK; Taipale, P; et al. (2013). Vasoactive agents for the prediction of early- and late-onset preeclampsia in a high-risk cohort. *BMC Pregnancy and Childbirth*, 13:110.
- Wang, A., Rana, S., & Karumanchi, S. (2009). Preeclampsia: The Role of Angiogenic Factors in Pathogenesis. *Physiology*, 147-158.
- Wang, Y., & Zhao, S. (2010). *Vascular Biology of the Placenta*. Louisiana: Morgan & Claypool Life Science.

- Wathen, KA; Tuutti, E; Stenman, UH; Alfthan, H; Halmesmaki, E; Finne, P; Ylikorkla, O; Vuorela, P. (2006). Maternal Serum-Soluble Vascular Endothelial Growth Factor Receptor-1 in Early Pregnancy Ending in Preeclampsia or Intrauterine Growth Retardation. *The Journal of Clinical Endocrinology & Metabolism* ,1, 180-184.
- Weel, I. C., Baergen, R. N., Romão-Veiga, M., Borges, V. T., Ribeiro, V. R., Witkin, S. S., et al. (2016). Association between Placental Lesions, Cytokines and Angiogenic Factors in Pregnant Women with Preeclampsia. *PLOS ONE*.
- Wikstrom, A.-K. (2007). *Biochemical and Epidemiological Studies of Early-Onset and Late-Onset Pre-Eclampsia*. Dissertation, Uppsala University, Faculty of Medicine, Swedish.

