

## DAFTAR PUSTAKA

1. Ramani G V, Uber PA, Mehra MR. Chronic heart failure: Contemporary diagnosis and management. *Mayo Clinic Proceedings*. 2010;85(2):180–95.
2. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Euro Heart Journal*. 2016;37(27):2129–2200m.
3. Fedele F, Mancone M, Adamo F, Severino P. Heart Failure With Preserved, Mid-Range, and Reduced Ejection Fraction: The Misleading Definition of the New Guidelines. *Cardiology Review*. 2017;25(1):4–5.
4. Shah S, Davies MK, Cartwright D, Nightingale P. Management of chronic heart failure in the community: role of a hospital based open access heart failure service. *Heart*. 2004;90(7):755–9.
5. Taha T, Sayed K, Saad M, Samir M. How accurate can electrocardiogram predict left ventricular diastolic dysfunction?. *Egypt Heart Journal*. 2016;68(2):117–23.
6. Bagaswoto HP, Dinarti LK, Maharani E. Scoring system based on electrocardiogram features to predict the type of heart failure in patients with chronic heart failure. *Journal Medical Science*. 2016;48(3):131–41.
7. Cincin A, Ozben B, Erdogan O. Diagnostic utility of specific electrocardiographical parameters in predicting left ventricular function. *Experimental and Clinical Cardiology*. 2012;17(4):210–4.
8. Tsunakawa H, Miyamoto N, Kawabata M, Mashima S. Electrocardiogram in heart failure. *Nippon Rinsho*. 2000;51(3):1222–32.
9. O’Neal WT, Mazur M, Bertoni AG, Bluemke DA, Al-Mallah MH, Lima JAC, et al. Electrocardiographic predictors of heart failure with reduced versus preserved ejection fraction: The multi-ethnic study of atherosclerosis. *J American Heart Association*. 2017;6(6):1–9.
10. Aiba T, Tomaselli GF. Electrical Remodeling in the Failing Heart. *Current Opinion Cardiology*. 2010;25(1):29–36.
11. Wang Y, Hill JA. Electrophysiological remodeling in heart failure. *Journal of Molecular and Cellular Cardiology*. 2010;48(4):619–32.
12. Owusu IK, Adu-boakye Y, Appiah LT. Cardiovascular Diseases & Diagnosis Electrocardiographic Abnormalities in Heart Failure Patients at a Teaching Hospital in Kumasi, Ghana. *Journal of Cardiovascular Diseases & Diagnosis*. 2014;2(2):2–4.
13. Nikolaidou T, Pellicori P, Zhang J, Kazmi S, Goode KM, Cleland JG, et al. Prevalence, predictors, and prognostic implications of PR interval prolongation in patients with heart failure. *Clinical Research Cardiology*. 2017;107(2):108–19.

14. Azevedo PS, Polegato BF, Minicucci MF, Paiva SAR, Zornoff LAM. Cardiac Remodeling: Concepts, Clinical Impact, Pathophysiological Mechanisms and Pharmacologic Treatment. *Arquivos Brasileiros de Cardiologia*. 2016;106(1):62–9.
15. Lensen JMA, Fosse VN, Bovendeerd PHM, Strijkers GJ, Kuijpers NHL, Dam PM. Electrophysiology of the heart. From cardiac electrical activity to the ECG. 2008;5-7.
16. Becker DE. Fundamentals of electrocardiography interpretation. *Anesthesia Progress*. 2006;53(2):53–63
17. Luna ABd. Ventricular Enlargement. In: A. B. d. Luna, ed. *Basic Electrocardiography Normal and Abnormal ECG Patterns*. 1st ed. Spain: Blackwell Publishing. 2007: 40-48
18. Oh C, Chang HJ, Sung JM, Kim JY, Yang W, Shim J, et al. Prognostic estimation of advanced heart failure with low left ventricular ejection fraction and wide QRS interval. *Korean Circulation Journal*. 2012;42(10):659–67.
19. Utari EL. Analisa Deteksi Gelombang QRS Untuk Menentukan Kelainan Fungsi Kerja Jantung. *Teknoin*. 2016;22:27–37.
20. Brendorp B, Elming H, Jun L, Kober L, Malik M, Jensen GB, et al. Qtc interval as a guide to select those patients with congestive heart failure and reduced left ventricular systolic function who will benefit from antiarrhythmic treatment with dofetilide. *Circulation*. 2001;103(10):1422–1427.
21. Karaye KM, Sani MU. Electrocardiographic abnormalities in patients with heart failure. *Cardiovascular Journal of Africa*. 2008;19(1):22–5.
22. Cenkerova K, Dubrava J, Pokorna V, Kaluzay J, Jurkovicova O. Clinical Study: Prognostic value of echocardiography and ECG in heart failure with preserved ejection fraction. *Bratislava Medical Journal*. 2016;117(7):407–412.
23. Lopez C, Ilie CC, Glancy DL, Quintal RE. Goldberger's electrocardiographic triad in patients with echocardiographic severe left ventricular dysfunction. *American Journal of Cardiology*. 2012;109(6):914–8.
24. Razavipour F, Nikoo MH, Abtahi F. Numerical Estimation of Ejection Fraction from 12-lead ECG. *Journal Bioengineering and Biomedical Science*. 2015;6(1):1-7
25. Xiong Y, Wang L, Liu W, Hankey GJ, Xu B, Wang S. The Prognostic Significance of Right Bundle Branch Block: A Meta-analysis of Prospective Cohort Studies. *Clinical Cardiology*. 2015;38(10):604–13.
26. Lu Z, Lense L, Sharma M. Prevalence of QT prolongation and associated LVEF changes in diabetic patients over a four-year retrospective time period. *Journal of Community Hospital Internal Medicine Perspectives*. 2017;7(2):87–94.

27. Padmanabhan S, Silvet H, Amin J, Pai RG, Linda L. Congestive Heart Failure Prognostic value of QT interval and QT dispersion in patients with left ventricular systolic dysfunction : Results from a cohort of 2265 patients with an ejection fraction of < 40 %. *American Heart Journal*. 2003;145:132-8.
28. Kashani A, Barold SS. Significance of QRS complex duration in patients with heart failure. *Journal of the American College of Cardiology*. 2005;46(12):2183–92.
29. Dhingra R, Pencina MJ, Wang TJ, Nam B-H, Benjamin EJ, Levy D, et al. Electrocardiographic QRS Duration and the Risk of Congestive Heart Failure: The Framingham Heart Study. *Hypertension*. 2006;47(5):861–7.
30. Kelly JP, Daubert JP. Prolonged QRS in Heart Failure With Preserved Ejection Fraction: Risk Marker and Therapeutic Target? *JACC Heart Failure*. 2016;4(6):487–9.
31. Dukes JW, Dewland TA, Vittinghoff E, Mandyam MC, Heckbert SR, Siscovick DS, et al. Ventricular Ectopy as a Predictor of Heart Failure and Death. *Journal of the American College of Cardiology*. 2015;66(2):101-109
32. Junior WN, West E, Santos M, Skali H, Groarke JD, Forman DE, et al. Heart Failure and Mid-Range Ejection Fraction: Implications of Recovered Ejection Fraction for Exercise Tolerance and Outcomes. *Circulation Heart Failure*. 2017;9(4):1–17
33. Rickenbacher P, Kaufmann BA, Maeder MT, Bernheim A, Goetschalckx K, Pfister O, et al. Heart failure with mid-range ejection fraction: a distinct clinical entity? Insights from the Trial of Intensified versus standard Medical therapy in Elderly patients with Congestive Heart Failure (TIME-CHF). *European Journal Heart Failure*. 2017;19(12):1586–96.
34. Steyerberg EW. *Clinical prediction models: a practical approach to development, validation and updating*. USA: Springer. 2009
35. Roger VL. Epidemiology of heart failure. *Circulation research*. 2013; 113(6):646-649
36. Deswal A, Bockert S. Comparison of morbidity in women vs men with heart failure and preserved ejection fraction. *American Journal of Cardiology*. 2006; 97: 1228-1231
37. Ghali JK, Krause-Steinraus HJ, Adams KT. Gender differences in advanced heart failure: insight from the BEST study. *Journal of the American College of Cardiology*. 2003; 42: 2128-2134
38. Azad N, Kathiravelu A, Minoosepeher S, Hebert P, Fergusson D. Gender differences in the etiology of heart failure: a systematic review. *Journal of Geriatric Cardiology*. 2011; 8: 15-23
39. Velagaleti RS, Gona P, Pencina MJ, Jayashri A, Wang TJ, Levy D, et al. Left ventricular hypertrophy patterns and incidence of heart failure with preserved versus reduced ejection fraction. *American journal of cardiology*. 2014;

113(1);1-16

40. Aro AL, Reiner K, Uy-Evanado A, Rusinaru C, Jui J, Chugh SS, et al. Prediction of moderate left ventricular systolic dysfunction from the 12-lead electrocardiogram. *European Heart Journal*. 2017;38 (Suppl 1):902
41. Olesen LL, Andersen A. ECG as a first step in the detection of left ventricular systolic dysfunction in the elderly. *ESC heart failure*. 2016;3:44-52
42. Sandhu R, Bahler RC. Prevalence of QRS prolongation in a community hospital cohort of patients with heart failure and its relation to left ventricular systolic dysfunction. *American Journal of Cardiology* 2004;93:244 –246.
43. Murkofsky RL, Dargas G, Diamond JA, Mehta D, Schaffer A, Ambrose JA. A prolonged QRS duration on surface electrocardiogram is a specific indicator of left ventricular dysfunction. *Journal of the American College of Cardiology*. 1998;32:476 –482
44. Vrtovec B, Yazdanbakhsh AP, Pintar T, Collard CD, Gregoric ID, Radovancevic B. QTc interval prolongation predicts postoperative mortality. *Texas Heart Institute Journal*. 2006; 33: 3-8
45. Fang, Wai Lee AP, Man Yu C. Review: Left atrial function in heart failure with impaired and preserved ejection fraction. *Current Opinion in Cardiology*. 2014; 29:430–436
46. Hanna EB, Glancy DL. ST-segment depression and T wave inversion: classification, differential diagnosis, and caveats. *Cleveland clinic journal of medicine*. 2011;78(6):404-414
47. Basnet BK, Manandhar K, Shrestha R, Thapa M. Electrocardiograph and chest X-Ray in prediction of left ventricular systolic dysfunction. *Journal of Nepal Medical Association*. 2009;48(176):310-3.

