

## RUJUKAN

- Anonim. (2014, Mei 10). Ethyl 4-methoxycinnamate [Web log post]. Diakses dari <http://www.lookchem.com/newseil/search.aspx?key=Ethyl%204-methoxycinnamate>.
- Arambewela, L., Perepa, A., Wijesundera, R.L.C., & Gunatileke, L. (2000). Investigation on *Kaempferia galanga*. *J. Natn. Sci. Foundation Sri Lanka*, 28(3), 225-230.
- Badan Pengawas Obat dan Makanan Republik Indonesia. (2011, Mei-Juni). Riset obat bahan alam di Pusat Riset Obat dan Makanan. *Info POM*, 12(3), 10-11.
- Badan Pusat Statistik. (2014). *Statistika Indonesia 2014*. Jakarta: BPS
- Bhuiyan, MD. N. I., Begum, J., & Anwar, M.N. (2008). Essential oils of leaves and rhizomes of *Kaempferia galangal* Linn. *The Chittagong Univ. J. B. Sci.*, 3(1&2), 65-76.
- Brown, W.H., Foote, C.S., Iverson, B.L., & Anslyn, E.V. (2012). *Organic chemistry* (6th ed.). California: Cengage Learning.
- Bruice, P.Y. (2004). *Organic chemistry* (4th ed.). New Jersey: Prentice Hall Inc.
- Carey, F.A. (2000). *Organic chemistry* (4th ed.). Boston: McGraw Hill Companies.
- Chan, E.W.C., Lim, Y.Y., Wong, L.F., Lianto, F.S., Wong, S.K., Lim, K.K., . . . Lim, T.Y. (2008). Antioxidant and tyrosinase inhibition properties of leaves and rhizomes of ginger species. *Food Chem*, 109, 477-483.
- Chang, Y.-C., Lee, F.-W., Chen, C.-S., Huang, S.-T., Tsai, S.-H., Huang, S.-H. & Lin, C.-M. (2007). Structure-activity relationship of C6-C3 phenylpropanoids on xanthine oxidase-inhibiting and free radical-scavenging activities. *Free Radical Biology & Medicine*, 43, 1541-1551.
- Choi, I. -H., Park, J. -Y., Shin, S. -C., & Park, I. -K. (2006). Nematicidal activity of medicinal plant extracts and two cinnamates isolated from *Kaempferia galanga* L. (Proh Hom) against the pine wood nematode, *Bursaphelenchus xylophilus*. *Nematology*, 8(3), 359-365.
- Choochote, W., Kanjanapothi, D., Panthong, A., Taesotikul, T., Jitpakdi, A., Chaithong, U., & Pitasawat, B. (1999). Larvicidal, adulticidal and repellent effects of *Kaempferia galanga*. *Southeast Asian J. Trop. Med. Publ. Health*, 30, 470-476.

- Departemen Kesehatan Republik Indonesia. (2008). *Farmakope herbal Indonesia*. (Edisi 1). Jakarta: Depkes RI.
- Dewoto, H.R. (2007). Pengembangan obat tradisional Indonesia menjadi fitofarmaka. *Majalah Kedokteran Indonesia*, 57(7), 205-211.
- European Medicines Agency. (2011). *Guideline on quality of herbal medicinal products/ traditional herbal medicinal products*. London.
- Gandjar, I.G. & Rohman, A. (2009). *Kimia farmasi analisis*. Yogyakarta: Pustaka Pelajar.
- Gritter, R.J., Bobbitt, J.M., & Schwarting, A.E. (1985). *Introduction to chromatography* (2nd ed.). Oakland: Holden-Day.
- Harborne, J.B. (1984). *Phytochemical methods: A guide to modern techniques of plant analysis* (2nd ed.). New York: Chapman and Hall.
- Handa, S.S., Khanuja, S.P.S., Longo, G., & Rakesh, D.D. (2008). *Extraction technologies for medicinal and aromatic plants*. Trieste: International Centre For Science and High Technology.
- Hong, T.K., Kim, S.I., Heo, J.W., Lee, J.K., Choi, D.R., & Ahn, Y.J. (2011). Toxicity of *Kaempferia galanga* rhizome constituents to *Meloidogyne incognita* juveniles and eggs. *Nematol*, 13, 235–244.
- Huang, L., Yagura, T., & Chen, S. (2008). Sedative activity of hexane extract of *Kaempferia galanga* L. and its active compounds. *Journal of Ethnopharmacology*, 120, 123-125.
- Kadota, S., Nguyen, M.T.T., Awale, S., Tezuka, Y., Tran, Q.L., & Watanabe, H. (2004). Xanthine oxidase inhibitory activity of Vietnamese medicinal plants. *Biol. Pharm. Bull.*, 27(9), 1414-1421.
- Kanjanapothi, D., Panthong, A., Lertprasertsuke, N., Taesotikul, T., Rujjanawate, C., Kaewpinit, D., . . . Pitasawat, B. (2004). Toxicity of crude rhizome extract of *Kaempferia galanga* L. (Proh Hom). *J. Ethnopharmacol*, 90, 359–365.
- Kochuthressia, K.P., Britto, S.J., Jaseentha, M.O., & Raphael, R. (2012). In vitro antimicrobial evaluation of *Kaempferia galanga* L. rhizome extract. *American Journal Biotechnology and Molecular Sciences*, 2(1), 1-5.
- Koh, H.L. (2009). *Guide to medicinal plants: An illustrated scientific and medicinal approach*. Singapore: World Scientific.

- Ko, H.-J., Kim, H. J., Kim, S. Y., Yun, H.-Y., Baek, K. J., Kwon, N. S., . . . Kim, D.-S. (2014). Hypopigmentary effects of ethyl *p*-methoxycinnamate isolated from *Kaempferia galangal*. *Phytother. Res.*, 28, 274-279.
- Kosuge, T., Yokota, M., Sugiyama, K., Saito, M., Iwata, Y., Nakura, M., & Yamamoto, T. (1985). Studies on anticancer principles in Chinese medicines.II. Cytotoxic principles in *Biota orientalis* (L.) Endl. and *Kaempferia galanga* L. *Chem. Pharm. Bull.*, 33, 5565-5567.
- Kumar, A. (2014). Chemical composition and biological activities of key constituents of essential oil from the rhizomes of *Kaempferia galanga* L. *International Journal of Pharma and Bio Sciences*, 5(1), 225-231.
- Kushwaha, S.K.S., Kushwaha, N., Maurya, N., & Rai, A.K. (2010). Role of markers in the standardization of herbal drugs: A review. *Arch. Appl. Sci. Res.*, 2(1), 225-229.
- Lakshmanan, D., Werngren, J., Jose, L., Suja, K.P, Nair, M.S., Varma, R.L., . . . Kumar, R.A. (2011). Ethyl *p*-methoxycinnamate isolated from a traditional anti-tuberculosis medicinal herb inhibits drug resistant strains of *Mycobacterium tuberculosis* in vitro. *Fitoterapia*, 82, 757-761.
- Li, S., Han, Q., Qiao, C., Song, J., Cheng, C.L., & Xu, H. (2008). Chemical markers for the quality control of herbal medicines: An overview. *Chinese Medicine*, 3, 7-23.
- Liu, B., Liu, F., Chen, C., & Gao, H. (2010). Supercritical carbon dioxide extraction of ethyl *p*-methoxycinnamate from *Kaempferia galanga* L. rhizome and its apoptotic induction in human HepG2 cells. *Nat. Prod. Res.*, 24, 1927-1932.
- Lundanes, E., Reubsaet, L. & Greibrokk, T. (2014). *Chromatography: Basic principles, sample preparations and related methods*. Weinheim, Germany: Wiley-VCH.
- Marks, D.A., Smith, C.M., & Lieberman, M.A. (1996). *Marks's basic medical biochemistry: A clinical approach* (2nd ed.). Lippincott Williams & Wilkins.
- Meyer, V.R. (2010). *Practical high-performance liquid chromatography* (5th ed.). West Sussex, UK: John Wiley and Sons, Ltd.
- Mitra, R., Orbell, J., & Muralitharan, M.S. (2007). Agriculture-medicinal plants of Malaysia. *Asia Pasific Biotech News*, 11, 105-110.

- Mukherjee, P.K. & Houghton, P.J. (2009). *Evaluation of herbal medicinal products : perspective on quality, safety and efficacy*. London: Pharmaceutical Press.
- Murray, R.K., Bender, D.A., Botham, K.M., Kennelly, P.J., Rodwell, V.W., & Weil P.A. (2009). *Harper's illustrated biochemistry* (28th ed.). New York: Mc-Graw Hill.
- Newcombe, D.S. & Robinson, D.R. (2013). *Gout : basic science and clinical practice*. London: Springer.
- Natural Health Products Directorate. (2013). *Quality of natural health products guide version 3.0*. Canada: Health Canada.
- National Center for Biotechnology Information. (2014, April 8). PubChem compound database; CID=5281783 [Web log post]. Diakses dari <https://pubchem.ncbi.nlm.nih.gov/compound/5281783>.
- Othman, R., Ibrahim, H., Mohd., M.A., Mustafa, M.R., & Awang, K. (2006). Bioassay-guided isolation of a vasorelaxant active compound from *Kaempferia galanga* L. *Phytomedicine*, 13, 61-66.
- Pavia, D.L., Lampman, G.M., Kriz, G.S., & Engel, R.G. (2011). *A small scale approach to organic laboratory techniques* (3rd ed.). Belmont, CA: Cengage Learning.
- Peter, K.V. (2006). *Handbook of herbs and spices volume 3*. Cambridge, UK: Woodhead Publishing Limited.
- Rao, V.K., Rajasekharan, P.E., Roy, T.K., & Kumar, V.T. (2009). Comparison of essential oil component in rhizomes and in-vitro regenerated whole plants of *Kaempferia galanga* L. *Journal of Medicinal and Aromatic Plant Sciences*, 31(4), 326-329.
- Riditid, W., Sae-Wong, C., Reanmongkol, W., & Wongnawa, M. (2008). Antinociceptive activity of the methanolic extract of *Kaempferia galanga* Linn. in experimental animals. *J. Ethnopharmacol*, 118, 225-230.
- Rostagno, M.A., & Prado, J.M. (2013). *Natural product extraction: Principles and applications*. Cambridge: RSC Publishing.
- Rostiana, O., Rosita, SMD., Rahardjo, M., & Taryono. (2005). *Budidaya tanaman kencur*. Bogor: Badan Penelitian dan Pengembangan Pertanian.
- Rukmana, R. (1994). *Kencur*. Yogyakarta: Penerbit Kanisius.

- Sarawek, S., (2007). *Xanthine oxidase inhibition and antioxidant activity of an artichoke leaf extract (Cynara scolymus L.) and its compounds*. (Disertasi). Florida: University of Florida.
- Sarker, S.D., Latif, Z., & Gray, A.I. (2006). *Natural product isolation* (2nd ed.). New Jersey: Humana Press Inc.
- Sarker, S.D. & Nahar, Lutfun. (2007). *Chemistry for pharmacy students: General, organic, and natural product chemistry*. West Sussex: John Wiley & Sons, Ltd.
- Seidemann, J. (2005). *World spices plants*. New York: Springer.
- Shoyama, Y. (2011). *Quality control of herbal medicines and related areas*. Rijeka, Croatia: InTech.
- Singh, B. C., Chanu, S. B., Bidyababy, Th., Devi, R. W., Singh, S.B., Nongalleima, Kh., . . . Singh, L.W. (2013). Biological and chemical properties of *Kaempferia galangal* L. – a Zingiberaceae plant. *NeBIO*, 4(4), 35-41.
- Sirisangtragul, W. & Sripanidkulchai, B. (2011). *Ethyl-p-methoxycinnamate (EPMC) and dichloromethane extract from Kaempferia galanga L. modulated pentobarbital-induced sleeping and hepatic cytochrome P450 enzyme activities in mice*. *KKU Res J (GS)*, 11(4), 9-18.
- Stabile, R.G. & Dicks, A.P. (2004). Two-step semi-microscale preparation of a cinnamate ester sunscreen analog. *Journal of Chemical Education*, 81(10), 1488-1491.
- Sutthanont, N., Choochote, W., Tuetun, B., Junkum, A., Jitpakdi, A., Chaltong, U., . . . Pitasawat, B. (2010). Chemical composition and larvicidal activity of edible plant derived essential oil against the pyrethroid susceptible and resistant strains of *Aedes aegypti* (Diptera: Culicidae). *J. Vector Ecol.*, 35, 106-115.
- Sweeney, A.P., Wyllie, S.G., Shalliker, J.L., & Markham, J.L. (2000). Xanthine oxidase inhibitory activity of selected Australian native plants. *Journal of Ethnopharmacology*, 75, 273-277.
- Tara S.V., Chandrakala, S., Sachidananda, A., Kurady, B.L., Smita, S., & Ganesh, S. (2006). Wound healing activity of alcoholic extract of *Kaempferia galanga* in Wistar rats. *Indian J. Physiol. Pharmacol*, 50, 384–390.

- Taufikurrohmah, T. (2005). Sintesis p-metoksisinamil p-metoksisinamat dari etil p-metoksisinamat hasil isolasi rimpang kencur (*Kaempferia galanga* L) sebagai kandidat tabir surya. *Indo. J. Chem.*, 5(3), 193-197.
- Tewtrakul, S. & Subhadhirasakul, S. (2007). Anti-allergic activity of some selected plants in the *Zingiberaceae* family. *J. Ethnopharmacol*, 109, 535–538.
- Tewtrakul, S., Yuenyongsawad, S., Kummee, S., & Atsawajaruwan, L. (2005). Chemical components and biological activities of volatile oil of *Kaempferia galanga* L.. *Songklanakarinn Journal of Science and Technology (Suppl.2)*, 503-507.
- Umamaheswari, M., Asokkumar K., Subhadradevi, V., & Sivashanmugam A.T. (2009). In vitro xanthine oxidase inhibitory activity of the fraction of *Erythrina stricta* Roxb. *J. Ethnopharmacol*, 124, 646-648.
- Umar, M.I., Asmawi, M.Z., Sadikun, A., Atangwho, I.J., Yam, M.F., Altaf, R., & Ahmed, A. (2012). Bioactivity-guided isolation of *Ethyl-p-methoxycinnamate*, an anti-inflammatory constituent, from *Kaempferia galanga* L. extracts. *Molecules*, 17, 8720-8734.
- Umar, M.I., Asmawi, M.Z., Sadikun, A., Majid, A.M., Al-Suede, F.S., & Hassan L.E. (2014). *Ethyl-p-methoxycinnamate* isolated from *Kaempferia galanga* inhibits inflammation by suppressing interleukin-1, tumor necrosis factor- $\alpha$ , and angiogenesis by blocking endothelial functions. *Clinics*, 69(2), 134-144.
- Voet, D. & Voet, J.G. (2011). *Biochemistry* (4th ed.). Danvers: John Wiley and Sons.
- Xue, Y. & Chen, H. (2002). Study on the anti-carcinogenic effects of three compounds in *Kaempferia galanga*. *Xei Shen Yan Jui*, 31(4), 247-251.
- Yenjai, C., Daodee, S., & Wangboonsakul, J. (2003). Antifungal activity and antimycobacterial activity of *ethyl p-methoxycinnamate* from *Kaempferia galanga* L. *Proceeding of the 3rd Symposium on the Family Zingiberaceae*, 193-195.
- Yumita, A., Suganda, A.G., & Sukandar, E.Y. (2013). Xanthine oxidase inhibitory activity of some Indonesian medicinal plants and active fraction of selected plants. *Int. J. Phar. Sci.*, 5(2), 293-296.