

DAFTAR PUSTAKA

1. Wiersinga W, Bonten M, Boersma W, Jonkers R, Aleva R, Kullberg B, et al. Management of community acquired pneumonia in Adults: 2016 Guideline update from SWAB. *Netherlands J of Med.* 2016;76: 1–13.
2. Cilloniz C, Loeches I, Vidal C, Jose A, Torres A. Microbial etiology of pneumonia: Epidemiology, diagnosis and resistance patterns. *Int J Molecul Sci.* 2016; 17:1–18
3. Jager C, Wever P, Gemen E, Kusters R, van Gageldonk-Lafeber A, Poll T, et al. The neutrophil-lymphocyte count ratio in patients with community acquired pneumonia. *Plos One.* 2012; 7: 1–8
4. Gibson G, Loddenkemper R, Lundback B, Sibille Y. Respiratory health and disease in Europe: The new European Lung White Book. *Eur Respir J.* 2013; 42:559–63
5. Rozenbaum M, Mangen M, Huijts S, van der Werf T, Postma M. Incidence, direct costs and duration of hospitalization of patients hospitalized with community acquired pneumonia: A nationwide retrospective claims database analysis vaccine. 2015; 28: 3193–9
6. Lim W, Baudouin S, George R. Pneumonia guidelines Committee of the BTS Standards of Care Committee. *BTS guidelines for management of community acquired pneumonia in adults: update 2009.* *Thorax.* 2011; 64: 1–55
7. RI KK. *Profil Kesehatan Indonesia Sitohang V, Budijanto D, Hardhana B, Soenardit, editors.* Jakarta: Kementerian Kesehatan RI; 2014.
8. Laporan Rekam Medis. Padang: RSUP.Dr. M. Djamil, Rekam Medis; 2016.
9. Moldoveanu B, Otmishi P, Jani P, Walker J, Sarmiento X, Guardiola J, et al. Inflammatory mechanisms in the lung. *J of Inflamm Research.* 2009; 2: 1–11
10. Balamayooran G, Batra S, Fessler M, Happel K, Jeyaseelan S. Mechanism of neutrophil accumulation in the lungs against bacteria. *Am J of Resp Cell and Mol Biol.* 2010; 43: 5–16
11. Mizgard J. Acute lower respiratory tract infection. *N Engl J Med.* 2008; 358: 716–27
12. Laickhalk L, Kunkel S, Strieter R. Tumor necrosis factor mediates lung antibacterial host defense in murine *Klebsiella pneumoniae*. *Infect & immun.* 1996; 64: 5211–8.

13. Tsurumi A, Que YA, Ryan CM, Tompkins RG, Rahme LG. TNF- α /IL-10 ratio correlates with burn severity and may serve as a risk predictor of increased susceptibility to infections. *Frontiers in Public Health*. 2016; 4(216): 1–6
14. Greene C, Lowe G, Taggart C, Gallagher P, McElvaney N, O'Neill S. Tumor necrosis factor- α - converting enzyme: Its role in community acquired pneumonia. *The journal of infectious disease*. 2002; 186: 1790–6
15. Puren AJ, Feldman C, Savage N, Becker PJ, Smith C. Patterns of cytokine expression in community acquired pneumonia. *CHEST*. 1995; 107(5): 1342–9
16. Akhayeva AS, Zhupenova DE, Kentzhetayeva TA, Ovsyannikova KA, Karpeeva DV. Changes in the level of TNF- α depending on severity in children with community acquired pneumonia. *Biomed Res*. 2017; 28(18): 8007–9
17. Baratawidjaja K, Rengganis I. Sel-sel sistem imun nonspesifik. Edisi 8. Jakarta: Balai penerbit Fakultas Kedokteran Universitas Indonesia; 2009: 142-54
18. Zahorec R. Ratio of neutrophil to lymphocyte counts-rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy*. 2001; 102: 5–14
19. Chalupa P, Beran O, Herwald H, Kasprkova N, Holub M. Evaluation of potential biomarker for the discrimination of bacterial and viral infections. *Infect Dis*. 2011; 7: 10
20. de Jager C, van Wijk P, Mathoera R, de Jough-Leuvenink J, van der Poll T. Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit. *Crit Care*. 2010; 14: 192
21. Lee JH , Song S, Yoon S, Lim C, Song J, Kim H. Neutrophil to lymphocyte ratio and platelet to lymphocyte ratio as diagnostic markers for pneumonia severity. *British J of Biomed Science*. 2016; 73: 140–2
22. Yang T, Wan C, Wang H, Qin J, Chen L, Shen Y, et al. The prognostic and risk-stratified value of neutrophil–lymphocyte count ratio in Chinese patients with community-acquired pneumonia. *European J of Inflamm*. 2017; 15: 22–7
23. Cataudella E, Giraffa C, di Marca S, Pulvirenti A, Alaimo S, Pisano M, et al.

Neutrophil-To-Lymphocyte Ratio: An emerging marker predicting prognosis in elderly adults with community-acquired pneumonia. *The J of American Ger soc.* 2017; 2: 2–6

24. Shah B, Ahmed W, Dhobi G, Shah N, Khurseed S, Haq I. Validity of pneumonia severity index and CURB-65 severity scoring systems in community acquired pneumonia in an Indian setting. *The J of Chest Dis and Allied Sciences.* 2010; 52: 9–17
25. Loke Y, Kwok C, Niruban A. How accurately do pneumonia severity scores predict mortality in patients with community acquired pneumonia. *Annals of Emerg Med.* 2012; 59: 51–2
26. Zhang Z, Zhang W, Liu P, Yang Y, Tan W, Ng H, et al. Prognostic value of Pneumonia Severity Index, CURB-65, CRB-65, and procalcitonin in community-acquired pneumonia in Singapore. *SAGE journals.* 2016; 25: 139–47
27. Madhu S, Augustine S, Kumar Y, Kauser K, Kumar V, Jayaraju B. Comparative study of CURB-65, Pneumonia Severity Index and IDSA/ATS scoring systems in community acquired pneumonia in an Indian tertiary care setting. *Int J of adv in med.* 2017; 4: 3
28. Wunderink R, Waterer G. Community Acquired pneumonia. *N Engl J Med.* 2014; 370: 543–51
29. Lutfiyya M, Henley E, Chang L. Diagnosis and treatment of community acquired pneumonia. *Am Fam Physician.* 2006; 73: 442–50
30. Brown J, Calverley P, Dusmet M, Elborn J, Fisher A, Fitzgerald J, et al. BTS guidelines for the management of community acquired pneumonia in adult: update 2009. *Thorax.* 2009; 64: 1–55
31. Hall M, Defrances C. National hospital discharge survey. *Adv data.* 2015; 1: 1–20
32. Castillo J, Sanchez F, Llinares P, Menendez R, Mujal A, Nava sE, et al. Guidelines for the management of community acquired pneumonia in the elderly patient. *Rev Esp Quimioter.* 2014; 27: 69–86
33. Niederman M, Mandell L, Anzueto A, Bass J, Brouhton W, Campbell G. American Thoracic Society. Guidelines for thr management of adults with community

- acquired of adults with community acquire pneumonia. Diagnosis, assestment of severity, antimicrobial therapy, and prevention. *Am J Resp Crit Care Med.* 2001; 163: 1730–54
34. Fine M, Auble T, Yealy D, Hanusa B, Weissfeld L, Singer D. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med.* 1997; 336: 243–50
35. Rajah A, Charles P, Melvani S, Burrell L, Johnson P, Grayson M. Comparing the pneumonia severity index with CURB65 in patients admitted with community acquired pneumonia. *Scan J Infect Dis.* 2008; 40(4): 293–300.
36. Greer JP, Arber DA, Glader B, List AF, Means RT, Paraskevas F, et al. *Wintrobe's Clinical Hematology.* 13th ed. Foerster J, editor. Philadelphia: Lippincott Williams & Wilkins; 2014. 2234–58
37. Gordon S. The macrofage: past, present and future. *Eur J Immunol.* 2007; 37: 9-17
38. Mukhopadhyay S, Hoidal JR, Mukherjee TK. Role of TNF- α in pulmonary pathophysiology. *Resp Research.* 2006; 7(125): 1–9
39. Williams AE, Hussell T, Lloid C. *Immunology: Mucosal and body surface defences.* Edisi 1. Oxford: Wiley-Blackwell; 2012: 152–61
40. Ryan D. *Examination of blood cells.* edisi 3. Kaushansky K, Lichtmalt M, Beutler E, Kipps T, Seligsohn U, Prchal J, editors. United State of America: MsGraw-Hill Education; 2010. 1287–96
41. Chen MM, Palmer JL, Plackett TP, Deburghraeve CR, Kovacs EJ. Age-related differences in the neutrophil response to pulmonary *Pseudomonas* infection. *Exp Gerontol.* 2015; 42–6
42. Kadioglu A, Gingles NA, Grattan K, Kerr A, Mitchell TJ, Andrew PW. Host cellular immune response to pneumococcal lung infection in mice. *Infect Immun.* 2000; 68(2): 492–501
43. Tateda K, Moore T, JC D, Newstead M, Zeng X, Matsukawa A, et al. Early recruitment of neutrophils determines subsequent T1/T2 host responses in a murine model of *Legionella Pneumophila Pneumonia.* *J Immunol.* 2001; 166: 3355–61
44. Schulte W, Bernhagen J, Bucala R. Cytokines in sepsis: potent immunoregulators

- and potential therapeutic targets- An update view. Hindawi Publishing Corp. 2013; 2013: 1–16
45. Burns A, Smith C, Walker D. Unique structural features that influence neutrophil emigration into the lung. *Physiol Rev.* 2003; 83: 309–36
 46. Filho J, Spiller F, Cunha F. Neutrophil paralysis in sepsis. *Shock.* 2010; 34: 15–21
 47. Craig A, Mai J, Cai S, Jeyaseelan S. Neutrophil recruitment to the lungs during bacterial pneumonia. *American Society for Microbiology.* 2009; 77: 568–75
 48. Yan C, Gao H. New insights for C5a and C5a receptors in sepsis. *Frontiers In Immunology.* 2012; 3: 1–11
 49. Meng F, Liu L, Chin P, D'Mello SR. Akt is a downstream target of NF-kappa B. *J Biol. Chem.* 2002; 10: 74–8
 50. Janeway C, Travers P, Walport M. *Immunobiology: The immune system in health and disease.* edisi 5. New York: Garland Science; 2001. 256–87
 51. Prussin C, Metcalfe D. IgE, mast cell, basophyl, and eosinophyl. *J Allergy Clin Immunol.* 2006; 117: 450–6
 52. Parrino J, Hotchkiss R, Bray M. Prevention of immune cell apoptosis as potential therapeutic strategy for severe infections. *Emerg Inf Dis.* 2007; 13: 191–8
 53. Gurol G, Ciftci I, Terzi H, Atasoy A, Ozbek A, Koroglu M. Are there standardized cutoff values for neutrophil-lymphocyte ratios in bacteremia or sepsis? *J Microbiol Biotechnol.* 2015; 25: 521–5
 54. Halazun K, Hardy M, Rana A, Woodland D, Luyten E. Negative impact of neutrophil-lymphocyte ratio on outcome after liver transplantation for hepatocellular carcinoma. *Ann Surg.* 2008; 250: 141–51
 55. Sarraf K, Belcher E, Raevsky E, Nicholson A, Goldstraw P. Neutrophil/lymphocyte ratio and its association with survival after complete resection in non-small cell lung cancer. *J Thorac Cardiovasc Surg.* 2009; 137: 425–8
 56. Rhee DY, Park SH, Choi HJ, Kwon MK, Cho DH. The value of neutrophil Lymphocyte Count Ratio for disease severity in nursing home acquired pneumonia patients. *J Korean Geriatr Soc.* 2013; 17(4): 213–8.
 57. Forget P, Khalifa C, Defour J, Latinne D, Van Pel M, De Kock M. What is the

- normal value of the neutrophil-to-lymphocyte ratio? *BMC Res Notes*. 2017; 10: 12–21
58. Shah BA, Singh G, Naik MA, Dhobi GN. Bacteriological and clinical profile of community acquired pneumonia in hospitalized patients. *Lung India*. 2010; 27(2): 54–7
59. Yoon NB, Son CH, Um SJ. Role of the neutrophil-lymphocyte count ratio in the differential diagnosis between pulmonary tuberculosis and bacterial community acquired pneumonia. *Ann lab Med*. 2013; 33(2): 105–10.
60. Calle LR, Seco JP, Vargas DA, Mascaros E, Redondo E, Maroto LD, et al. Incidence and risk factor prevalence of community acquired pneumonia in adults in primary care in Spain. *BMC inf Dis*. 2016; 16(645): 1–8
61. Liu X, Shen Y, Wang H, Ge Q, Fei A, Pan S. Prognostic significance of neutrophil-to-lymphocyte ratio in patients with sepsis: a prospective observational study. *Mediat of Inflamm*. 2016 Mar; 2016(1): 2–8.
62. Bermejo-martin JF, Cilloniz C, Mendez R, Almansa R, Gabarrus A, Ceccato A, et al. Lymphopenic community acquired pneumonia (LCAP), an immunological phenotype associated with higher risk of mortality. *Ebiomedicine*. 2017; 24: 231–6.
63. Kumar N, Rafiudeen R, Rashmi K. A study of clinical and etiology profile of community acquired pneumonia with special reference to atypical pneumonia. *Annals of Nigerian Medicine*. 2017; 11(1): 11–16.
64. Pantzaris ND, Platanaki C, Pierrako C, Karamouzos V, Velissaris D. Neutrophil-to-lymphocyte ratio relation to sepsis severity scores and inflammatory biomarkers in patients with community acquired pneumonia: a case series. *J Transl Int Med*. 2018; 6(1): 43–6.
65. Almirall J, Bolibar I, Serra-Prat M. New evidence of risk factors for community acquired pneumonia: a population based study. *Eur Respir J*. 2008; 31(1): 1274–84.
66. Jalil ANA. Profil pasien pneumonia komunitas di Rumah Sakit Umum Daerah Cengkareng tahun 2013-2014. Skripsi. Jakarta: UIN Syarif Hidayatullah, Fakultas Kedokteran dan Ilmu Kesehatan; 2015.

67. Luna CM, Palma I, Niederman MS, Membriani E, Giovini V, Wiemken TL, et al. The impact of age and comorbidities on the mortality of patients of different age groups admitted with community acquired pneumonia. *Ann Am Thorac Soc.* 2016; 13(9): 1519–26.
68. Lowsby R, Gomes C, Jarman I, Lisboa P, Nee PA, Vardhan M, et al. Neutrophil to lymphocyte count ratio as an early indicator of blood stream infection in the emergency department. *Emerg Med J.* 2014; 11(36): 1–5.
69. Ramirez JA, Wiemken TL, Peyrani P, Arnold FW, Kelley R, Maltingly WA, et al. Adults hospitalized with pneumonia in the United States: incidence, epidemiology, and mortality. *Clin Infect Dis.* 2017 July; 2017(65): 1806–12.
70. Nilson B. The immune system in the oldest-old clinical and immunological studies in the nona immune cohort Sweden: Linkoping; 2010.437-42
71. Dykstra B, de Haan G. Hematopoietic stem cell aging and self-renewal. *Cell Tissue Res.* 2008; 331: 91–104
72. Wesemann T, Nullmann H, Pflug MA, Heppner HJ, Pientka L, Thiem U. Pneumonia severity, comorbidity and 1-year mortality in predominantly older adults with community acquired pneumonia: a cohort study. *BMC Infect Dis.* 2015; 15(2): 1–10
73. Shehata SM, Sileem AE, Shahien NE. Prognostic values of pneumonia severity index, CURB65 and expanded CURB65 scores in community acquired pneumonia in Zagazig University Hospitals. *Egyptian J of Chest Dis and Tuberculosis.* 2017; 66(3): 549–55
74. Torres A, Peetermans WE, Viegi G, Blasi F. Risk factors for community acquired pneumonia in adults in Europa: a literature review. *Thorax.* 2013; 68(1): 1057–65
75. Taqwaningtyas E. Profil pasien pneumonia komunitas rawat inap di Rumah Sakit Paru Respira Yogyakarta periode 2014-2018. Skripsi. Yogyakarta: Universitas Islam Indonesia, Fakultas Kedokteran; 2018.
76. Polverino E, Torres MA. Community acquired pneumonia. *Minerva Anestesiol.* 2011; 77(1): 196–211.
77. Holub M, Lawrence DA, Andersen N, Davidova A, Beran O, Maresova V, et al.

Cytokines and chemokines as biomarkers of community acquired bacterial infection. *Med of Inflamm.* 2013; 2013(1): 1–8.

78. Kaya Y, Tas N, Canakci E, Cebeci Z, Ozbilen M, Keskin H, et al. Relationship of neutrophil to lymphocyte ratio with presence and severity of pneumonia. *Journal of clinical and analytical medicine.* 2018; 9(5): 452–7.
79. Ravindranath M, Raju C. Validity of pneumonia severity index/ pneumonia outcome research trial and CURB65 severity scoring systems in community acquired pneumonia in Indian setting. *International Journal of Advances Medicine.* 2016; 3(2): 338–44
80. Bacci M, Zing N, urad N, Adami F, Hinnig P, Feder D, et al. IL6 and TNF- α serum levels are associated with early death in community acquired pneumonia patients. *Braz J MedBiol Res.* 2015; 48(5): 427–32
81. Yang J, Azat M, Peng P, Chang Y, Gao R, Li W, et al. Serum levels of TNF- α , IL-1B, IL-9, and IL-15 in acute respiratory distress syndrome. *Int J Clin Exp Pathol.* 2017; 10(1): p. 781-8.

