

## DAFTAR PUSTAKA

1. Hammelgarn BR. Epidemiology of chronic kidney disease scope of the problem. In Kimmel P , Rosenberg M , editors. Chronic Renal disease.: Elsevier; 2014. 57–67.
2. Zhang L, Wang H. Chronic kidney disease in developing countries. In Turner N, editor. Oxford textbook of clinical nephrology 4th edition. Oxford: Oxford University Press; 2016. 762–66.
3. (PERNEFRI) PNI. 8th Report of Indonesia Renal Registry (IRR). 2015; 2:30–8.
4. Inker L, Astor B, Fox C, Isakova T, Lash J, Peralt J. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management. AJKD. 2014; 63(5):713–35.
5. Hill N, Fatoba ST , Oke JL , Hirst JA , O'Callaghan CA , Lasserson DS , et al. Global prevalence of chronic kidney diseases- a systematic review and meta analysis. Plos One. 2016.
6. Institute for health metrics and evaluation. The global burden of disease : generating evidence, guiding policy. IHME. 2010; 23(2):42–9
7. Chung S, Koh E, Shin S, Park C. Malnutrition in patients with chronic kidney disease. Scientific research. 2012; 2:89–99.
8. Kopple J. National kidney foundation K/DOQI clinical practice guidelines for nutrition in chronic renal failure. AJK. 2001; 37:S66–S70.
9. Lopes A, Bragg-Gresham JL , Elder SJ. Independent and joint association of nutritional status indicators with mortality risk among chronic hemodialysis patient in the Dialysis outcomes and practice patterns study(DOPPS). Journal of renal nutrition. 2010; 20:224–34.
10. Habas E, Rayani A, Khammaj A. Long term complication of hemodialysis. Sebha Med J. 2012; 11 (1) :12–8.
11. Kovesdy C. Malnutrition in dialysis patients- the need for intervention despite uncertain benefits. Semin Dial. 2015; 29: 28–34.
12. Wetmore J, Collins AJ. Global challenges posed by the growth of end-stage renal disease. Renal replacement therapy. 2016; 2:15.
13. Stenvinkel P. Inflammation in end stage renal disease- a fire that burns within. Contrib nephrol. 2005; 149:185–99.

14. Stenvinkel P , Heimburger O , Paultre F. Strong association between malnutrition, inflammation and atherosclerosis in chronic renal failure. *Kidney Int.* 1999; 55:1899–911.
15. Lindholm B, Heimburger O, Stenvinkel P. What are the cause of protein-energy malnutrition in chronic renal insufficiency? *AJKD*. 2002; 39:422–5.
16. Tsujinaka T, Fujita J, Ebisuri C. Interleukin-6 receptor antibody inhibits muscle atrophy and modulates proteolytic system in interleukin-6 transgenic mice. *JCI*. 1996; 97:244–49.
17. Pecoit-Filho R, Lindholm B, Axelsson J, Stenvinkel P. Update on interleukin-6 and its role in chronic renal failure. *Nephrol Dial Trans.* 2003; 18: 1042–45.
18. Kopple J. Pathophysiology of protein -energy wasting in chronic renal failure. *ASNS*. 1999; 129:247–51.
19. Iklizer AT. Protein and energy : Recommended intake and nutrient supplementation in chronic dialysis patients. *Sem in Dial*. 2004; 17:471-478.
20. Kalantar-Zadeh K. Proteine energy wasting. In Byham-Gray , Burrowes J, Chertow G, editors. Nutrition and kidney disease second edition. London: Humana Press; 2014. 219–30.
21. Obi Y, Kalantar-Zadeh K , Kovesdy CP , Qader H. Latest consensus and update on protein energy-wasting in chronic kidney disease. *Curr Opin Clin Nutr Metab Care*. 2015; 18(3): 254–62.
22. Wang X, Mitch W. Mechanism of muscle wasting in chronic kidney disease. *NRN*. 2014; 10(9): 504–16.
23. Fouque D, K Kalantar-Zadeh , J Kopple. A proposed nomenclature and diagnostics criteria for protein-energy wasting in acute and chronic kidney disease. *Kidney Int*. 2008; 73:391–98.
24. Syaiful H, Oenzil F, Afriant R. Hubungan Umur dan Lamanya Hemodialisis dengan Status Gizi pada Penderita Penyakit Ginjal Kronik yang menjalani Hemodialisis di RS.Dr.M.Djamil Padang. *JKA*. 2014; 3(3):381–86.
25. Bologa RA , Levine DM , Parker TS. Interleukin-6 predicts hypalbuminemia , hypcholesterolemia, and mortality in hemodialysis patients. *AJKD*. 1998; 32:107–14.
26. Carero JJ , Qureshi AR , Avesani CM , Suliman ME , Kato S , Jonsdottir SS. Comparison of nutritional and inflammatory markers in dialysis patients with

- reduce appetite. AJCN. 2007; 85:695–701.
27. Kaizu Y, Kimura M, Yoneyama T, Miyaji K, Hibi I, Kumagai H. Interleukin-6 may mediate malnutrition in chronic hemodialysis patients. AJKD. 1998; 31: 93–100.
  28. Goodman M. Interleukin-6 Induces Skletal Muscle Protein Breakdown in Rats. PSEBM. 1994; 205(2)182–5.
  29. Jones C, Akbani H, Croft D, Worth D. The relationship between serum albumin and hydration status in hemodialysis patients. JRN. 2002; 12:209–12.
  30. Yeun J, Kaysen G. Factors influencing serum albumin in dialysis patients. AJKD. 1998; 32:118–25.
  31. Liu Y, Coresh J, Eustace J, Longenecker J, Jaar B, Fink N, et al. Association between cholesterol level and mortality in dialysis patient. JAMA. 2004; 291:451–459.
  32. Honda H , Qureshi AR , Heimburger O , Barany P , Wang K , Pecoits-Filho R , et al. Serum albumin, C-reactive protein, interleukin 6, and fetuin A as predictors of malnutrition , cardiovascular disease, and mortality in patients with ESRD. AJKD. 2006; 47:139–48.
  33. Desbrow B, Bauer J , Blum J. Assesment of nutritional hemodialysis patient using patient-generated subjective global assesment. JRN. 2005; 15: 211–6.
  34. Detsky A, J McLaughkin , J Baker , N Johnston , S Whittaker , R Mandelson , et al. What is subjective global assesment of nutritional status? JPEN. 1987; 11:8–13.
  35. Kwang A, Kandiah M. Objective and subjective nutritional assesment of patients with cancer in palliative care. AJPC. 2010; 27(2):117–26.
  36. National kidney fondation. KDIGO 2012: Clinical Practice Guideline for the evaluation and management Chronic KIdney Disease. 2013; 3(1) ;1–163.
  37. NKF/KDOQI. Guidelines Nutritional for Adults. 2002.
  38. Stenvinkel P, Alverstrand A. Inflammation in end-stage renal disease: source, consequences and theray. Semin Dial. 2002; 15:329–337.
  39. Carero J, P Stenvinkel , L Cuppari , TA Ikizler , K Kalantar -Zadeh , G Keysen , et al. Etiology of the protein-energy wasting syndrom in chronic kidney disease : A consensus statement from the international society of

- renal nutrition and metabolism (ISRN). JRN. 2013; 23: 77–90.
40. Holden RM , Morton AR , Garland JS. Vitamin K dan D status in stages 3-5 chronic kidney disease. CJASN. 2010; 5 (4) : 590–7.
  41. Steiber AL. Chronic kidney disease: considerations for nutrition interventions. JPEN. 2014; 20 (10):1–9.
  42. Beharka AA , Wu D , Meydani M. Interleukin-6 production does not increase with age. J Geront. 2001; 56: B81–88.
  43. Pupim L, Martin CJ , Ikizler TA. Assesment of protein and energy nutritional status. In Kopple J, GS M, K KZ, editors. Nutritional management of renal disease 3rd edition.: Elsevier; 2013. 137–58.
  44. Baberashvili I , Azar A , Sinuani I , Kadoshi H , Shapiro G , Feldman L. Chronic kidney disease. CJASN. 2013; 8:443–451.
  45. Velasquez M, Mehrotra R , Wing M , Raj D. Causes of protein-energy wasting in chronic kidney disease. In Kopple J, Massry GS , Kalantar-Zadeh K , editors. Nutritional management of renal disease 3rd edition.: Elsevier; 2013. 159–70.
  46. Baberashvili I, Sinuani I, Azar A , Yasur H, Shapiro G, Feldman L, et al. IL-6 levels, Nutritional status, and Mortality in Prevalent Hemodialysis Patients. CJASN. 2011; 6:2253–63.
  47. Gabay C , Kushner I. Acute phase proteins and other systemic responses to inflammation. NEJM. 1999; 340:448–55.
  48. Lee B , Ahmed F , Hamm L , Teran F , Chen C , Liu Y. Association of C-reactive protein, tumor necrosis factor-alpha, and interleukin-6 with chronic kidney disease. BMC Nephrology. 2015; 16:77.
  49. Kishimoto T. IL-6: from its discovery to clinical applications. Int Immunol. 2010; 22:347–52.
  50. Barreto V, Barreto F , Liabeuf s, Temmar M, Lemke H, Triboulloy C, et al. Plasma interleukin-6 is independently associated with mortality in both hemodialysis and pre-dialysis patient with chronic kidney disease. ISN. 2010; 77;550–6.
  51. Kopf M , Baumann H , Freer G. Impaired immune and acute-phase responses in interleukin-6-deficient mice. Nature. 1994; 368: 339–49.
  52. Jones A , Scheller J , Rose-John S. Therapeutic strategies for the clinical blockade of IL-6/gp130 signaling. JCI. 2011; 121: 3375–83.

53. Jones S A , Frase D J , Fielding C A , Jones G W. Interleukin-6 in renal disease and therapy. NDT. 2014; 2:50–9.
54. Stevinkel P, Katteler M, Johnsosn R. Interleukin- 1,IL-6 and TNF alpha: important factors in altered cytokine network of end stage renal disease-the good,the bad and the ugly. Kidney Int. 2005; 67:1216–33.
55. Rao M, Wong C, Kanetsky P, Girndt M, Stenvinkel P, Reilly M, et al. Cytokine gene polymorphism and progression of renal and cardiovascular disease. Kidney Int. 2007; 72:549–56.
56. Nascimento M, Pecoits-Filho R, Lindohlm B, Riella M, Stenvinkel P. Inflammation, malnutrition and atherosclerosis in end-stage renal disease: a global perspective. Blood Purf. 2002; 20:454–58.
57. Meuwese C, Carrero J, Stenvinkel P. Recent insight in inflammation-associated wasting in patients with chronic kidney disease. Contrib Nephrol. 2011; 171:120–6.
58. Jonkowska M, Cobo G, Lindholm B, Stenvinkel P. Inflammation and protein -energy wasting in the uremic milieu. Contrib Nephrol. 2017; 191:68–71.
59. Mafra D , Farage NF , Lobo JC , Stockler-Pinto MB , Leal VO , Carvalho DP. Relationship between total ghrelin and inflammation in hemodialysis patients. peptides. 2011; 32:358–61.
60. Kalantar-Zadeh K, Block G, McAllister C. Appetite and inflammation, nutrition, anemia and clinical outcome in hemodialysis patient. AJCN. 2004; 80:299–307.
61. Kaizu Y, Kimura M, Yoneyama T, Miyaji K, Hibi I, Kumagai H. Interleukin-6 may mediated malnutrition in chronic hemodialysis patients. AJKD. 1998; 31:93–100.
62. Ma H , Sun G , Wang W , Zhou Y , Liu D , Tong Y , et al. Association between interleukin-6 -572 C>G and -174 G>C polymorphisms and hypertension. Medicine. 2016; 95:1–15.
63. Thandavan T, Abraham G, Soundrarajan P, Chandrasekaran V. Interleukin 6 strongly correlates with malnutrition inflammation score and it is strong predictor for mortality in hemodialysis patients. Urologi & Nephrology open acces journal. 2015; 2(2):20–30.
64. Oliveira CM , Kubrusly M , Mota RS , Silva CA , Oliviera VN. Malnutrition inchronic kidney failure : what is the best diagnostic method to assess? JBN. 2010; 32:55–68.

65. Friedman A, Fadem S. Reassessment of albumin as a nutritional marker in kidney disease. *JAS*. 2010; 21: 223–30.
66. Delano M, Moldawer L. The origins of cachexia in acute and chronic inflammatory disease. *NCP*. 2006; 21:68–81.
67. Kaizu Y, Ohkawa S, Odamaki M. Association between inflammatory mediators and muscle mass in long term hemodialysis patients. *AJKD*. 2003; 42:295–302.
68. Avesani C, Qureshi A, Lindholm B, Carrero J, Stevinkel P. Inflammation and wasting in chronic kidney disease: partner in crime. *Kidney International*. 2006; 70:S8–S13.
69. Mitch WE , DU J , Bailey JE. Mechanism causing muscle proteolysis in uremia. *Miner elect Lab*. 1999; 25: 216–9.
70. Seen J, Klover P, Nowak I. Interleukin -6 induce cellular insulin resistant in hepatocytes. *Diabetes*. 2002; 51:3391–9.
71. Burrgereit , Burmester G, Brand M. Bioenergetics of immune functions: fundamental and therapeutic aspect. *Immune Today*. 2000; 16:556–65.
72. Ottery FD. Rethinking nutritional support of the cancer patient: the new field of nutritional oncology. *Sem Oncology*. 1994; 6:770–8.
73. Vasantha J, Soundararajan P , Vanitha Rani N , Kannan G , Thennarasu P , Chacko RA , et al. Prediction of malnutrition using modified subjective global assesment-dialysis malnutrition score in patient on hemodialysis. *IJPS*. 2011; 73(11): 38–45.
74. Fink JS , de Mello PD , de Mello ED. Subjective Global Assessment of nutritional status – a systematic review of the literature. *Cilinical Nutrition*. 2014;(S0261-5614(14)00312-4):1–31.
75. Zabel S , Ash N , King P , Juffts J , Bauer. Relationship between appetite and quality of life in hemodialysis patients. *Apetite*. 2012; 58:194–9.
76. Mohammed F, Farhood H, Abdul-Atheemwt M. Prediction of malnutrition using modified subjective global asessment- dialysis malnutrition score in patients on chronic hemodialysis. *CMHE*. 2014; 4:291–8.
77. Santos CA , Rosa C , Ribeiro A , Ribeiro R. Patient generated subjective global assesment and classic anthropometry: comparison between the methods in detevtion of malnutrition among elderly with cancer. *Nutr Hosp*. 2015; 31(1):384–92.

78. Havinga-Kosters M, Julicher A, Duvan A, Yucesan E, van Essen, Berg AVD, et al. Diagnostic accuracy of PGSGA and MUST of patient with chronic kidney disease. CNJ. 2016; S1:91–2.
79. Kemenkes RI. In Pedoman nasional pengendalian tuberkulosis. Jakarta: Kementerian kesehatan RI; 2014. 15–7.
80. Rochmah W , Harimurti K. Demensia. In Setiati S , Sudoyo A , Alwi I , Simadibrata M , Setiyohadi B , Syam AF , editors. Buku Ajar Ilmu Penyakit Dalam.: Interna Publishing; 2014. 3801–09.
81. Mujaddid E. Pemahaman dan penanganan psikosomatik gangguan ansietas dan depresi di bidang ilmu penyakit dalam. In Setiati S , Alwi I , Sudoyo A , Simadibrata M , Setiyohadi B , Syam AF , editors. Buku ajar ilmu penyakit dalam edisi VI. Jakarta: Internal Publishing; 2014. 3581–84.
82. Dewantari EO , Taruna A , Angraini DI , Dilangga P. Hubungan adekuasi hemodialisis dengan asupan makan dan indeks massa tubuh pada pasien gagal ginjal kronik yang menjalani hemodialisis di RSUD Abdul Moeloek Bandar Lampung. MJLU. 2014; 3:24–8.
83. Sari R , Sugiarto S , Probandari A , Hanim D. Hubungan asupan energi, protein, vitamin B6, natrium dan kalium terhadap status gizi pada pasien gagal ginjal kronik dengan hemodialisis. JAB. 2017; 6:34–43.
84. Goldberg I , Krause I. The role of gender in chronic kidney disease. EMJ. 2016; 1(2):58–64.
85. Campell K, Bauer JD , Ikehiro A , Johnson WD. Role of nutition impact symptoms in predicting nutritional status and clinical outcome in hemodialysis patients: A potential screening tool. JRN. 2013; 23:302–7.
86. Alwyn T. Nutritional status of haemodialysis patients: comparison of Australian cohorts of Aboriginal and European descent. Griffith research onlline. 2013.<http://hdl.handle.net/10072/58225>.
87. Roslaini R. Hubungan kadar methylgyoxal dengan gangguan fungsi kognitif pada pasien penyakit ginjal kronik yang menjalani hemodialisa. Tesis. 2015; 47–57.
88. Kesehatan SBL. Riskesdas 2018. , Kemenkes RI; 2018.
89. Spatola L , Finazzi S , Calvetta A , Reggiani F , Morenghi E , Santostasi S , et al. Subjective global assesment-dialysis malnutrition score and cardiovascular risk in hemodialysis patients: an observational cohort study. Journal of Nephrology. 2018; 31:757–65.

90. Kaysen GA , Greene T , Mehta L , Lindsay RM , Depner TA , Hall YN , et al. The effect of frequent hemodialysis on nutrition and body composition: Frequent hemodialysis network trial. *ISN*. 2012; 82:90–9.
91. Post JB , Morin KG , Sano M. Increased presence of cognitive impairment in hemodialysis patients inthe absence of neurological events. *AJN*. 2012; 35:120–6.
92. McQuillan JR , Sarbjit V. Neuropsychiatric complications of chronic kidney disease. *NRN*. 2010; 6:471–9.
93. Borges MCC , Vogt BP , Martin LC , Caramori JCT. Malnutrition inflammation score cut-off predicting mortality in maintenance hemodialysis patients. *Clinical nutrition ESPEN*. 2016; 30:1–5.
94. Runtung Y , Kadir A , Semana A. Pengaruh hemodialisa terhadap kadar ureum kreatinin dan hemoglobin pada pasien GGK di ruang hemodialisa RSUP Dr Wahidin Sudirohusodo Makassar. *JIK*. 2013; 2(3):1–7.
95. Horl W , Vanrenterghem Y , Aljama P , Brunet P , Brunkhorst R , Gesualdo L , et al. OPTA: Optimal treatment of anemia in patients with chronic kidney disease. *NDT*. 2007; 22(3):iii20–iii26.
96. Su H , Tao Lei C , Zhang C. Interleukin -6 signaling pathway and its role in kidney disease:an update. *Frontiers in immunology*. 2017; 8:12–18.
97. Kamimura MA , Draibe SA , Dalboni MA , Cendoroglo M , Avesani CM , Manfredi SR , et al. Serum and cellular interleukin-6 in haemodialysis patients: relationship with energy expenditure. *NDT*. 2007; 22: 839–44.
98. Fishman D , Faulds G , Jeffery R. The effect of novel polymorphism in the interleukin-6 gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic -onset juvenile chronic arthritis. *JCI*. 1998; 102:1369–76.
99. Hukkanen J , Pertovaara M , Antonen J , Pasternack A , Hurme M. Elevated IL-6 plasma level are regulated by the promoter region polimorphism of the IL-6 gene in primary Sjogren's syndrome and correlate with the clinical manifestations of the disease. *Rheumatology(Oxford)*. 2001; 40: 656–61.
100. Kuijken M, Lichtart S, de Wit-van der Werf L, Donders M, van Gastel A, van den Berg G, et al. Assesment of nutritional status in haemodialysis patients using PG-SGA and handgrip strenght. *MMC*. 2015; 34:256–7.
101. Carero JJ. Identification of patients with eating disorders: clinical and biochemical signs of appetite loss in dialysis patients. *JRN*. 2009; 19(1):10–5.

102. Rambod M , Bross R , Zitterkoph J , Benner D , Colman S , Kovesdy C , et al. Association of malnutrition inflammation score with quality of life and mortality in maintenance hemodialysis patients: a -5 year prospective cohort study. AJKD. 2009; 53(2): 298–309.
103. Beberashvili I , Azar a , Sinuani i , Kadoshi H , Shapiro G , Feldman L , et al. Comparison analysis of nutritional scores for serial monitoring of nutritional status in hemodialysis patients. CJASN. 2013; 8:443–51.
104. Dai L , Mukai H , Lindholm B , Heimburger O , Barany P , Stenvinkel P , et al. Clinical global assesment of nutritional status as predictor of mortality in chronic kidney disease. pone. 2017; <https://doi.org/10.1371/journal.pone.0186659>.
105. Jones CH , Wolfenden RC , Wells LM. Is subjective global assesment a reliable measure of nutritional status in hemodialysis? JRN. 2004; 14(1) :26–30.

