

DAFTAR PUSTAKA

- Abas MH, Kader AJA, Khalil MS, Hamid AA, Isa MHM. 2013. Production of surfactin from *Bacillus subtilis* ATCC 21332 by using palm oil mill effluent (POME) as fermentation media. *IPCBE* 55 (17): 87-93.
- Abdulsalam, M., Man, H. C., Idris, A. I., Yunus, K. F., and Abidin, Z. Z. (2018b). Treatment of palm oil mill effluent using membrane bioreactor: novel processes and their major drawbacks. *Water* 10:1165. doi: 10.3390/w10091165
- Ahmad, A. L Ismail, S. and Bathia, S. (2003). Water recycling from palm oil mill effluent (pome) using membrane technology. *Desalination*. (157). 87-95
- Aisami, A., Yasid, N.A., Johari, W.L.W., Ahmad, S.A., Shukor, M.Y., 2020. Effect of temperature and pH on phenol biodegradation by a newly identified *Serratia* sp. Aq5-03. *Open Journal of Bioscience Research (OJBR)* 1 (1), 28–43.
- Akmal. 2010. Biodegradasi fenol limbah cair industri tekstil oleh *Candida tropicalis* [Skripsi]. Bogor (ID): Fakultas Matematika dan ilmu Pengetahuan Alam, Institut Pertanian Bogor.
- Atlas, RM. (1981). *Microbial Degradation of Petroleum Hydrocarbons: an Environmental Perspective*. *Microbiol Rev* 45(1):180-209.
- Ayed, L., Othman, B. and Hamdi Moktar, N. (2002). Decolorization of olive mill wastewaters by co-culture of *Geotrichum candidum* and *Lactobacillus plantarum*. In: *Proceedings of the International Symposium on Environmental Pollution Control and Waste Management (EPCOWM*2002)*. 7-10, January. Tunis, Tunisia. pp. 7-10.
- Baehaki A, Nopianti R, Saputra E, and Gofar N. 2019. Exploration of Protease Enzyme Producing Bacteria from Water in Tanjung Senai Swamp Indralaya South Sumatra. In: Herlinda S (eds) *Prosiding Seminar Nasional Lahan Suboptimal 2019*, Palembang 4-5 September 2019.
- Bala JD, Lalung J. 2015. Studies on the reduction of organic load from palm oil mill effluent (POME) by bacterial strains. *Int J Recycl Org Waste Agricult* 1 (4): 1-10.
- Bala, J. D., Lalung, J., Al-Gheethi, A. A. S., Kaizar, H., and Ismail, N. (2018). Reduction of Organic Load And Biodegradation Of Palm Oil Mill Effluent By Aerobic Indigenous Mixed Microbial Consortium Isolated From Palm Oil Mill Effluent (POME). *Water Conserv. Sci. Eng.* 3,139–156. doi: 10.1007/s41101- 018- 00439
- Bao, L.L., Li, D., Li, X.K., Huang, R.X., Zhang, J., Yang, L.V. and Xia, G.Q. (2007). Phosphorus accumulation by bacteria isolated from a continuous-flow two-sludge system. *Journal of Environmental Sciences*. 19(4), 391-395.

- Baraniya, D., Puglisi, E., Ceccherini, M. T., Pietramellara, G., Giagnoni, L., Arenella, M., et al. (2016). Protease encoding microbial communities and protease activity of the rhizosphere and bulk soils of two maize lines with different N uptake efficiency. *Soil Biol. Biochem.* 96, 176–179. doi: 10.1016/j.soilbio.2016.02.001
- Cappuccino, J. G. and C. Welsh. 2017. *Microbiology: a Laboratory Manual*. 11th Ed. Pearson Education, Inc. Edinburgh Gate Harlow, England.
- Cappuccino, J. G. and N. Sherman. 2019. *Microbiology a laboratory manual* (12th Ed). Pearson Education Inc. Publishing as Benjamin Cummings San Francisco. CA.
- Choong, Y. Y., Chou, K. W., and Norli, I. (2018). Strategies for improving biogas production of palm oil mill effluent (POME) anaerobic digestion: a critical review. *Renew. Sustain. Energy Rev.* 82, 2993–3006. doi: 10.1016/j.rser.2017.10.036
- Cooney, JJ. (1984). *The Fate of Petroleum Pollutans In Fresh Water Ecosystem - Petroleum Microbiology*.
- Dellamatrice PM, Silva-Stenico ME, de Moraes LAB, Fiore MF, Monteiro RTR. 2017. Degradation of textile dyes by cyanobacteria. *Brazilian Journal of Microbiology.* 48 : 25-31.wu
- de Souza, P. M., de Assis Bittencourt, M. L., Caprara, C. C., de Freitas, M., de Almeida, R. P. C., Silveira, D., et al. (2015). A biotechnology perspective of fungal proteases. *Braz. J. Microbiol.* 46, 337–346. doi: 10.1590/S1517-838246220140359
- Dibble, JT., Bartha R. (1979). Effect of Environmental Parameters on The Biodegradation of Oil Sludge. *Appl Environ Microbiol* 37(4):729-739.
- Direktorat Jenderal Perkebunan. 2015. *Statistik Perkebunan Indonesia: Kelapa Sawit 2014-2016*. Jakarta (ID): Direktorat Jenderal Perkebunan.
- Rajagukguk, D. P. (2018). *Isolasi dan Identifikasi Bakteri Pendegradasi Limbah Cair Pabrik Kelapa Sawit sebagai Alternatif Agen Dekolorisasi Limbah*.
- Dwidjoseputro, D. 1985. *Dasar-dasar Mikrobiologi*. Djambatan. Malang.
- Fitriana, F., Baharuddin, M., & Sappewali, S. (2016). Isolasi dan Identifikasi Bakteri Pendegradasi Fenol yang Bersumber Dari Danau Tempe Kabupaten Wajo Sulawesi Selatan. *Al-Kimia*, 4(2), 132-141.
- Fu, G., Zhao, L., Huangshen, L., and Wu, J. (2019). Isolation and identification of a salt-tolerant aerobic denitrifying bacterial strain and its application to saline wastewater treatment in constructed wetlands. *Bioresour. Technol.* 290:121725. doi: 10.1016/j.biortech.2019.121725
- Hassen-Aboushiba, A., Ramli, R., and Sofian-Azirun, M. (2013). Ecological characteristics of POME ponds with reference to study some of their invertebrate species in Peninsular Malaysia. *Journal of Animal and Plant Sciences.* 23 (5), 1305-1315.

- Hermosilla, E., Rubilar, O., Schalchli, H., da Silva, A. S. A., Ferreira-Leitao, V., and Diez, M. C. (2018). Sequential white-rot and brown-rot fungal pretreatment of wheat straw as a promising alternative for complementary mild treatments. *Waste Manag.* 79, 240–250. doi: 10.1016/j.wasman.2018.07.044
- Igwe, J. C. and Onyegbado, C.C. (2007). A review of palm oil mill effluent (POME) water treatment. *Global Journal of Environmental Research.* (2). 54-62
- Irvan, Trisakti B, Vincent M, Tandean Y. 2012. Pengolahan lanjutan limbah cair kelapa sawit secara aerobik menggunakan Effective Microorganism guna mengurangi nilai TSS. *Jurnal Teknik Kimia USU* 2 (1): 27-30.
- Juwita, rahmy 2014, Isolasi dan Karakterisasi Bakteri Pendegradasi Fenol Dari Limbah Cair Rumah Sakit Umum Daerah (RSUD) Arifin Achmad Pekanbaru, *Jom Fmipa*, 1(2): 229-237
- Louhasakul, Y., Cheirsilp, B., and Prasertsan, P. (2016). Valorization of Palm Oil Mill Effluent into Lipid and Cell-Bound Lipase by Marine Yeast *Yarrowia lipolytica* and Their Application in Biodiesel Production. *Waste Biomass Valoriz.* 7, 417–426. doi: 10.1007/s12649-015-9451-7
- Mohammed RR, Chong MF. 2014. Treatment and decolorization of biologically treated palm oil mill effluent (POME) using banana peel as novel absorbent. *Journal of Environmental Management* 132 (2014): 237-249.
- Nasution DY. 2004. Pengolahan limbah cair pabrik kelapa sawit yang berasal dari kolam akhir (final pond) dengan proses koagulasi melalui elektrolisis. *Jurnal Sains Kimia* 2 (8):38-40.
- Neoh, C. H., Yahya, A., Adnan, R., Majid, Z. A., and Ibrahim, Z. (2013). Optimization of decolorization of palm oil mill effluent (POME) by growing cultures of *Aspergillus fumigatus* using response surface methodology. *Environmental Science and Pollution Research.* 20 (5), 2912-2923.
- Nursanti I. 2013. Karakteristik limbah cair pabrik kelapa sawit pada proses pengolahan anaerob dan aerob. *Jurnal Ilmiah Universitas Jambi.* 4 (13): 67-73.
- Ohimain, E. I., and Izah, S. C. (2017). A review of biogas production from palm oil mill effluents using different configurations of bioreactors. *Renew. Sustain. Energy Rev.* 70, 242–253. doi: 10.1016/j.rser.2016.11.221
- Oswal N, Sarma PM, Zinjarde SS, Pant A (2002) Palm oil mill effluent treatment by a tropical marine yeast. *Bioresour Technol* 85(1):35–37
- Prakash, B. and rfan, M. 2011. *Pseudomonas aeruginosa* s Presentin Crude Oil Contaminated Sites of Barmer Region (India). *J Bioremed Biodegrad* 2011, 2:5
- Prasad, M. P., & Manjunath, K. (2011). Comparative study on biodegradation of lipid-rich wastewater using lipase producing bacterial species.
- Prasertsan, P., Khangkhachit, W., Duangsuwan, W., Mamimin, C., and O-Thong, S. (2017). Direct hydrolysis of palm oil mill effluent by xylanase enzyme to

enhance biogas production using two-steps thermophilic fermentation under non-sterile condition. *Int. J. Hydrogen Energy* 42, 27759–27766. doi: 10.1016/j.ijhydene.2017.05.140

- Purwanto D. 2013. Pengolahan limbah cair kelapa sawit menggunakan reaktor UAF (Upflow Anaerobic Filter). *Jurnal Riset Industri Hasil Hutan* 1 (5): 1- 7.
- Putra DS, Putra A. 2014. Analisis pencemaran limbah cair kelapa sawit berdasarkan kandungan logam, konduktivitas, TDS, dan TSS. *Jurnal Fisika Unand* 2 (3): 96-101.
- Rahmiati, R. dan Mumpuni, M. 2017. Eksplorasi bakteri asam laktat kandidat probiotik dan potensinya dalam menghambat bakteri patogen. *Elkawanie: Journal of Islamic Science and Technology*, 3(2), 141- 150
- Ratpukdi T. 2012. Declorization of anaerobically treated palm oil mill wastewater using combined coagulation and vacuum ultraviolet-hydrogen peroxide. *International Journal of Chemical Engineering Applications* 5 (3): 333-336.
- Sami M. 2012. Penyisihan COD, TSS, dan pH dalam limbah cair domestik dengan fixed-bed column up flow. *Jurnal Reaksi* 21 (10):1-11.
- Sharma, P., Singh, T. A., Bharat, B., Bhasin, S., and Modi, H. A. (2018). Approach towards different fermentative techniques for the production of bioactive actinobacterial melanin. *Beni Suf Univ. J. Basic Appl. Sci.* 7, 695–700. doi: 10.1016/j.bjbas.2018.08.002
- Sudarsono A. 2008. Isolasi dan Karakterisasi Bakteri pada Ikan Laut dalam Spesies Ikan Gindara (*Lepidocibium flavobronneum*). Skripsi. Bogor: Institut Pertanian Bogor.
- Tamrin, K. F., and Zahrim, A. Y. (2017). Determination of optimum polymeric coagulant in palm oil mill effluent coagulation using multiple-objective optimisation on the basis of ratio analysis (MOORA). *Environ. Sci. Pollut. Res.* 24, 15863–15869. doi: 10.1007/s11356-016-8235-3
- Udiharto, M. (1996). Bioremediasi minyak bumi. Di dalam: *Prosiding Pelatihan dan Lokakarya Peranan Bioremediasi dalam Pengelolaan Lingkungan; Cibinong. 24-28 Juni 1996.* hlm 24-39.
- Waluyo L. 2007. *Mikrobiologi Umum*. Edisi Revisi. Malang: Universitas Muhammadiyah Malang.
- Wiley, J. M., Sherwood, L. M. dan Woolverton., 2008. Prescott, Harley, and Klein's *Microbiology Seventh Edition*. Newyork: McGraw-Hill.
- Wong, K. M., Abdul Rahman, N., Vikineswary, S., and Hassan, M. A. (2008). Enzymatic hydrolysis of palm oil mill effluent solid using mixed cellulases from locally isolated fungi. *Research Journal of Microbiology*. 3 (6), 474-481.
- Wu TY, Mohammad AW, Jahim JM, Anuar N. 2007. Palm oil mill effluent (POME) treatment and bioresourcess recovery using ultrafiltration membrane: Effect of pressure on membrane fouling. *Biochem Eng J* 3 (35): 309-317.

Wu TY, Mohammad AW, Jahim JM, Anuar N. 2010. Pollution control technologies for the treatment of palm oil mill effluent (POME) through end-of-pipe processes. *Journal of Environmental Management*.91 : 1467- 1490.

Xu, R., Zhang, K., Liu, P., Han, H., Zhao, S., Kakade, A., et al. (2018). Lignin depolymerization and utilization by bacteria. *Bioresour. Technol.* 269, 557–566. doi: 10.1016/j.biortech.2018.08.118

Xu, Z., Qin, L., Cai, M., Hua, W., and Jin, M. (2018). Biodegradation of kraft lignin by newly isolated *Klebsiella pneumoniae*, *Pseudomonas putida*, and *Ochrobactrum tritici* strains. *Environ. Sci. Pollut. Res.* 25, 14171–14181. doi: 10.1007/s11356-018-1633-y

Yunita M, Hendrawa Y, Yulianingsih R. 2015. Analisis kuantitatif mikrobiologi pada makanan penerbangan (Aerofoof ACS) Garuda Indonesia berdasarkan TPC (Total Plate Count) dengan metode Pour Plate. *Jurnal Keteknikaan Pertanian Tropis dan Biosistem* 3 (3): 237-248.

