

DAFTAR PUSTAKA

1. Kementerian Kesehatan RI. Pedoman nasional asuhan pasca keguguran yang komprehensif. Kementerian Kesehatan Republik Indonesia. 2020. 11–15 p.
2. Sarwono Prawirohardjo. Ilmu kebidanan sarwono prawirohardjo. 3rd ed. Jakarta: PT Bina Pustaka Sarwono Prawirohardjo; 2010. 460 p.
3. Mehdi Moradinazar M x, Rostami R, Armaghan N, Shakiba M, Rai A, Abbasi Azizi S, *et al*. Epidemiological features of spontaneous abortion in the north africa and the middle east from 1990 to 2019: results from the global burden of disease study 2019. *J Fam Reprod Heal*. 2022;16(3):183–91.
4. Hikmah K. Faktor risiko umur ibu yang berisiko tinggi terhadap kejadian abortus. *Indones J Kebidanan*. 2017;1(2):113.
5. Linda Y. Gambaran faktor resiko insidensi abortus di rsud raa soewondo pati. *Medisains*. 2015;13(2):62–76.
6. Dinas Kesehatan Sumatera Barat. Profil dinas kesehatan tahun 2015. 2015;
7. Gan J, Gu T, Yang H, Ao Z, Cai G, Hong L, *et al*. Non-coding rnas regulate spontaneous abortion: a global network and system perspective. *Int J Mol Sci*. 2022;23(8):1–2.
8. Sapra KJ, Louis GMB, Sundaram R, Galea S, Ananth C V, Health C. Time-varying effects of signs and symptoms on pregnancy loss <20 weeks: Findings from a preconception prospective cohort study. *Paediatr Perinat Epidemiol*. 2018;32(1):30–9.
9. Volgsten H, Jansson C, Darj E, Stavreus-Evers A. Women's experiences of miscarriage related to diagnosis, duration, and type of treatment. *Acta Obstet Gynecol Scand*. 2018;97(12):1491–8.
10. Susiana S. Program keluarga harapan dan penurunan angka kematian ibu (studi di provinsi jambi dan provinsi kalimantan selatan). *Aspirasi*. 2019;10(1):19–31.
11. Snell RS. Anatomi klinis: berdasarkan sistem. Jakarta: EGC; 2012. 800–833 p.
12. Fatkhiyah N, Rejeki ST, Atmoko D. Kepatuhan kunjungan antenatal care berdasarkan faktor maternal. *J SMART Kebidanan*. 2020;7(1):29.
13. Moore J. First trimester bleeding. *Am Fam Physician*. 2009;79(11):985–91.
14. Bourne T, Bottomley C. When is a pregnancy nonviable and what criteria should be used to define miscarriage?. *Fertil Steril*. 2012;98(5):1091–6.
15. Kiyokawa T, Tanaka T. Change in blood fibronectin level as a prognostic factor for threatened abortion. *Japan Soc Obstet Gynecol*. 1988;(268):1207.
16. Martín GR, Prieto JP, Veiga De Cabo J, Gomez Lus L, Barberán J, Landa

- JMG, *et al.* Plasma fibronectin as a marker of sepsis. *Int J Infect Dis.* 2004;8(4):236–43.
17. Menéndez V, Fernández-Suárez A, Galán JA, Pérez M, García-López F. Diagnosis of bladder cancer by analysis of urinary fibronectin. *Urology.* 2005;65(2):284–9.
 18. Sherwood L. *Fisiologi manusia: dari sel ke sistem.* 6th ed. Yesdelita. N, editor. Jakarta: EGC; 2011. 846–865 p.
 19. WHO. Policies, programme and services for comprehensive abortion care comprehensive abortion care in south-east asia region. 2021.
 20. Kementerian Kesehatan RI. Indikator program kesehatan masyarakat dalam rpjmn dan renstra kementerian kesehatan 2020-2024. Kementeri Kesehat RI. 2020;1–99.
 21. Puspita Sari RD, Kurniati I, Utami N, Prabowo AY. Hubungan riwayat abortus dengan kejadian mola hidatidosa pada wanita usia reproduktif di rsud abdul moeloek lampung. 2018. 1–46 p.
 22. Siti Mas’udah, Tutik Ekasari HR. Hubungan usia ibu dan paritas dengan kejadian abortus di klinik utama panasea. *J Ilm Obs.* 2023;15.
 23. Lestari ID, Setiadi T, Zahrotun L. Penerapan data mining menggunakan metode naïve bayes untuk klasifikasi tindakan jenis abortus di rsud duta mulya. *J Sarj Tek Inform.* 2018;6(2):60–8.
 24. Matjino SH. Faktor risiko kejadian abortus di rsud dr. Chasan boesoirie ternate provinsi maluku utara. 2013.
 25. Hendriks E, Macnaughton H, Castillo M. First trimester bleeding: evaluation and management. *am fam physician.* 2019;99(3):166–74.
 26. Berghella V, Saccone G. Fetal fibronectin testing for reducing the risk of preterm birth. *Cochrane Database Syst Rev.* 2019;(7).
 27. Yamaguchi Y, Isemura M, Yosizawa Z, Kurosawa K, Yoshinaga K, Sato A, *et al.* Changes in the distribution of fibronectin in the placenta during normal human pregnancy. *Am J Obstet Gynecol.* 1985;152(6):715–8.
 28. Mercurio F, Votino AMC, Spizie A Di, Nappi C. Fetal fibronectin as predictor of successful induction of mid-trimester. *Acta Obs Gynecol Scand.* 2005;(84):390–4.
 29. Risza Choirunnisa, Indrayani T. Hubungan konsumsi vitamin c dengan kejadian ketuban pecah dini pada ibu hamil di puskesmas mustika jaya bekasi jawa barat tahun 2019. 2019;
 30. Kuhrt K, Hezelgrave-Elliott N, Stock SJ, Tribe R, Seed PT, Shennan AH. Quantitative fetal fibronectin for prediction of preterm birth in asymptomatic twin pregnancy. *Acta Obstet Gynecol Scand.* 2020;99(9):1191–7.
 31. J C, Lockwood, Andrew ES, M. Renate D, David C, Kumudin D, *et al.* Fetal fibronectin in cervical and vaginal secretion as a predictor of preterm

- delivery. New English J Med. 1991;323(16):1120–3.
32. Honest H, Bachmann LM, Gupta JK, Kleijnen J, Khan KS. Accuracy of cervicovaginal fetal fibronectin test in predicting risk of spontaneous preterm birth: systematic review. BMJ. 2002;325:1–2.
 33. Yeni CM, Tambunan R, Hasanuddin. Threatened preterm labor: which are become preterm labor?. Indones J Obstet Gynecol. 2020;8(4):216–22.
 34. Son M, Miller ES. Predicting preterm birth: Cervical length and fetal fibronectin. Semin Perinatol. 2017;41(8):445–51.
 35. Dahlman MS. Statistik untuk kedokteran dan kesehatan: deskriptif, bivariat, dan multivariat, dilengkapi aplikasi dengan menggunakan SPSS. 2013. 159 p.
 36. Lequin RM. Enzyme immunoassay (EIA)/enzyme-linked immunosorbent assay (ELISA). Clin Chem. 2005;51(12):2415–8.
 37. Hayrapetyan H, Tran T, Tellez-Corrales E, Madiraju C. Enzyme-Linked Immunosorbent Assay: Types and Applications. Methods Mol Biol. 2023;2612:1–17.
 38. Constantine NT, Callahan JD, Watts DM. Retroviral Testing. Essential for Quality Control and Laboratory Diagnosis. 1992. 105–107 p.
 39. Santoso K, Herowati UK, Rotinsulu DA, Murtini S, Ridwan MY, Hikman DW, et al. Comparison of Colorimetric-Based Rabies Postvaccination Antibody Titer Detection Using Elisa Reader and Mobile Phone Camera. J Vet. 2021;22(1):79–85.
 40. Noer RI, Ermawati E, Afdal A. Karakteristik ibu pada penderita abortus dan tidak abortus di RS Dr. M. Djamil Padang tahun 2011–2012. J Kesehat Andalas. 2016;5(3):575–83.
 41. Ahmad LOAI. Analisis faktor resiko usia kehamilan dan paritas terhadap kejadian abortus. J Al Maiyyah. 2016;9(1):133.
 42. Puspitasari RD, Ayu PR, Utami N. Hubungan antara polip serviks dengan ancaman abortus pada kehamilan muda. J Kedokt Unila. 2018;2(Juli 2018):157–61.
 43. Wafiyatunisa Z, Rodiani. Hubungan obesitas dengan terjadinya preeklampsia. Majority. 2016;5(5):184–90.
 44. Pipin Karlensi, Siti Aisyah, Merisa Riski. Faktor-faktor yang berhubungan dengan kejadian abortus. J Kebidanan J Ilmu Kesehat Budi Mulia. 2023;13(1):90–8.
 45. Speziale P, Arciola CR, Pietrocola G. Fibronectin and its role in human infective diseases. Cells. 2019;8(12).
 46. Min J, Watson HA, Hezelgrave NL, Seed PT, Shennan AH. Ability of a preterm surveillance clinic to triage risk of preterm birth: a prospective cohort study. Ultrasound Obstet Gynecol. 2016;48(1):38–42.

47. Zhang S, Fu S, Liu Y, Chen L, Zhu L, Tan J, *et al*. Predictive value of fetal fibronectin on the embryonic loss of patients with recurrent spontaneous abortion in early pregnancy. *Clin Exp Obstet Gynecol.* 2014;41(4):380–3.
48. Goldenberg RL, Klebanoff M, Carey JC, Macpherson C, Leveno KJ, Moawad AH, *et al*. Vaginal fetal fibronectin measurements from 8 to 22 weeks ' gestation and subsequent spontaneous preterm birth. *Am J Obs Gynecol.* 2000;183(2):469–75.
49. Lefhalm B. a fibronectin in the plasma during pregnancy and parturition. *Gynecol Obstet Invest.* 1996;41:183–8.
50. Tatra G, Graninger W, Nasr F, Hoyer E, Pirich K. Plasma fibronectin in normal pregnancy and EPH-gestosis. *Geburtsh u Frauenheilk.* 1983;43:594–7.
51. Nowacek GE, Meyer WR, McMahon MJ, Thorp JR, Wells SR. Diagnostic value of cervical fetal fibronectin in detecting extrauterine pregnancy. *Fertil Steril.* 1999;72(2):302–4.

