

## DAFTAR PUSTAKA

1. Silva VAR, Pauna HF, Lavinsky J, Hyppolito MA, Vianna MF, Leal M, et al. Task force Guideline of Brazilian Society of Otology. *Brazilian Journal Otorhinolaryngology*. 2022;89: 159–189.
2. Lieu JEC, Kenna MA, Anne S, Davidson L. Hearing loss in children. *JAMA*. 2020;324:21-95.
3. Kemenkes RI. Pedoman nasional pelayanan kedokteran tata laksana tuli sensorineural kongenital Indonesia. Keputusan Menteri Kesehatan Republik Indonesia; 2022.
4. Anastasiadou S. Hearing loss. StatPearls NCBI Bookshelf. 2023; <https://www.ncbi.nlm.nih.gov/books/NBK542323>. Diakses September 2023.
5. Ratuszniak A, Skarzynski PH, Gos E, Skarzynski H. The bonebridge implant in older children and adolescents with mixed or conductive hearing loss audiological outcomes. *International Journal of Pediatric Otorhinolaryngology*. 2019; 118: 97–102.
6. Word Health Organization (2022) . Newborn health. World Health Organization. Western Pacific Region. <https://www.who.int/westernpacific/health-topics/newborn-health>. Diakses Januari 2022.
7. Zhou X, Wang L, Feng J, Guo Y, Zhou Y, Zhang X, et al. The prevalence and risk factors for congenital hearing loss in neonates: a birth cohort study based on challenge study. *International Journal of Pediatric Otorhinolaryngology*. 2022;162:2-3.
8. Haukedal C, Lyxell B, Wie OB. Health-Related Quality of Life with Cochlear Implants: The Children's Perspective. *Ear And Hearing Journal*. 2019;41:330–343.
9. Hajare P, Mudhol RS. A study of JCIH (Joint Commission on Infant Hearing) risk factors for hearing loss in babies of NICU and well baby nursery at a tertiary care center. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2021;74:83–90.
10. Stuart A. Effect of delivery mode on neonate auditory brainstem responses to air and bone conducted stimuli. *International Journal of Pediatric Otorhinolaryngology*. 2020;139:11-23.
11. Shinta N, Novira A. Hubungan kejadian asfiksia neonatorum dengan gangguan fungsi koklea pada neonatus. *Sriwijaya Journal Of Medicine*. 2021;4:60–65.
12. Vohr B. Ear and Hearing Disorders. In: Elsevier eBooks. 2018. p.1558-1566.
13. Young A. Otoacoustic emissions. StatPearls NCBI Bookshelf. 2023; <https://www.ncbi.nlm.nih.gov/books/NBK580483/>. Diakses September 2023
14. Zizlavsky S, Supartono N, Zachreini I, Bashiruddin J, Haryuna TSH, Savitri E, et al. Factors associated with time of diagnosis and habilitation of congenital hearing loss in Indonesia: A multicenter study. *International Journal of Pediatric Otorhinolaryngology*. 2022;163:11-16.
15. Wroblewska S, Greczka G, Dąbrowski P, Szyfter W, Mazela J. The results of newborn hearing screening by means of transient otoacoustic emissions

- has anything changed over 10 years. International Journal of Pediatric Otorhinolaryngology. 2018;96:4–10.
- 16. Megantara I, Tsidkenu M, Chadir L, Anggraeni R, Sylviana N. Relation between risk factor of hearing loss and the result of otoacoustic emission in newborns at Santosa Hospital Bandung Central. Hearing, Balance and Communication. 2021;19:167–174.
  - 17. Drake R, Vogl AW, Tibbitts R, Richardson P. Gray's anatomy for students fourth edition.2019.
  - 18. Hayes S, Ding D, Salvi RJ, Allman BL. Anatomy and physiology of the external, middle and inner ear. In: Handbook of Clinical Neurophysiology. Elsevier B.V. 2013. p. 3–23.
  - 19. Hansen J, Netter FH. Netter's clinical anatomy fifth edition. Elsevier; 2021.p.435-470.
  - 20. Milenkovic I, Schiefer U, Ebenhoch R, Ungewiss J. Anatomy and physiology of the auditory pathway. Ophthalmologe Journal.2020;117:1068–1073.
  - 21. Garry W. Foundations of speech and hearing anatomy and physiology second edition. In: Plural publishing.2022.p.207-233.
  - 22. McFarland DH. Netter's atlas of anatomy for speech, swallowing, and hearing. Elsevier;2022.p.173-203.
  - 23. White H. Anatomy, head and neck, ear organ of Corti. StatPearls NCBI Bookshelf. 2023; <https://www.ncbi.nlm.nih.gov/books/NBK538335/>. Diakses September 2023
  - 24. Pickles JO. Auditory pathways. In: Handbook of Clinical Neurology. Elsevier B.V. 2015;129:3–25.
  - 25. World Health Organization(2021). World report on hearing: executive summary.<https://iris.who.int/handle/10665/339956>. Diakses Maret 2023
  - 26. Kaplama M, Yukkaldiran A. Newborn hearing screening results comparison of Syrian and Turkish newborns factors influencing the difference. International Journal of Pediatric Otorhinolaryngology. 2020;138:2-5.
  - 27. World Health Organization(2021). Ear and hearing care indicators for monitoring provision of services. World Health Organization south-east Asia Region. <https://www.who.int/multi-media/details/ear-and-hearing-care---searo>. Diakses Januari 2023
  - 28. Hardani A, Goodarzi E, Delphi M, Badfar G. Prevalence and risk factors for hearing loss in neonates admitted to the neonatal intensive care unit a hospital study. Cureus. 2020;12:1-7.
  - 29. Nieman C, Oh ES. Hearing loss. Annals of Internal Medicine. 2020;173:81–96.
  - 30. Damayanti M, Muyassaroh M. Kurang pendengaran pada anak sindrom down. Medica Hospitalia Journal of Clinical Medicine. 2022;9:236–41.
  - 31. Kreicher K, Weir FW, Nguyen SA, Meyer TA. Characteristics and Progression of Hearing Loss in Children with Down Syndrome. The Journal of Pediatrics. 2018;193:27-33.
  - 32. Faizah Z, Rochmah N, Faizi M, Hisbiyah Y, Nugroho PS. Bunga Rampai Deteksi Dini Kelainan Genetik. Airlangga University Press; 2022.
  - 33. Orland MD, Giampietro PF. Genetic etiology of syndromic congenital hearing loss. The Journal of Pediatrics. 2022;245:38.

34. Masuda S, Usui S. Comparison of the prevalence and features of inner ear malformations in congenital unilateral and bilateral hearing loss. International Journal of Pediatric Otorhinolaryngology. 2019;125:7-92.
35. Wroblewska S, Dąbrowski P, Szyfter W, Mazela J. Universal newborn hearing screening methods and results, obstacles, and benefits. Pediatric Research. 2018;81:15–22.
36. Dimopoulou D, Kourlaba G, Antoniadou A, Mariolis L, Kavatha D, Stoungioti S, et al. Low birth weight and head circumference as potential biomarkers of sensorineural hearing loss in asymptomatic congenitally CMV infected infants. Journal of Clinical Virology. 2020;129:104.
37. Yu Y, Shi K, Nielson C, Graham EM, Price MS, Haller TJ, et al. Hearing loss caused by CMV infection is correlated with reduced endocochlear potentials caused by stria damage in murine models. Hearing Research . 2022;417:108.
38. Kataoka Y, Maeda Y, Fukushima K, Sugaya A, Shigehara A, Kariya S, et al. Prevalence and risk factors for delayed-onset hearing loss in early childhood a population based observational study in Okayama Prefecture, Japan. International Journal of Pediatric Otorhinolaryngology. 2020;138:110-298.
39. Keihanidost Z. Risk factors for hearing loss and its prevalence in neonates older than 6 months with history of hospitalization in Intensive Care Unit. PubMed Central. 2018.
40. Wang Y, Liu L, Huang Y. Effect of antenatal dexamethasone therapy on hearing screening in premature infants a retrospective case-control study. American Journal of Otolaryngology. 2023;44:103.
41. Hirvonen M, Ojala R, Korhonen P, Haataja P, Eriksson K, Gissler M, et al. Visual and hearing impairments after preterm birth. Pediatrics. 2018;142:38-88.
42. Nair V, Janakiraman S, Whittaker S, Quail J, Foster T, Loganathan PK. Permanent childhood hearing impairment in infants admitted to the neonatal intensive care unit: nested case-control study. European Journal of Pediatrics. 2021;180:2083–9.
43. American Academy of Audiology. Year 2019 position statement principles and guidelines for early hearing detection and intervention programs.American Academy of Audiology. 2019; 4(2): 1–44.
44. Gultom S, Rambe AYM. Hubungan riwayat kelahiran prematur dan BBLR dengan gangguan pendengaran pada anak di RSUP haji adam malik periode 2016-2018. Scripta Score Scientific Medical Journal. 2021;2:84–91.
45. Girsang J, Safranto S, Lubis YM, Nasution SW. Gambaran hasil pemeriksaan otoakustik emisi (oae) sebagai skrining awal pendengaran pada bayi baru lahir di rsu royal prima medan. Jurnal Darma Agung. 2022;30:793-802.
46. Suchitra D, Nagaveni DP, Chary DC. Hearing assessment in asphyxiated term neonates in a tertiary, care hospital. IOSR Journal of Dental and Medical Sciences.2018;16:9-24.
47. Alenazi A, Derraugh G, Levesque M, Morris M, Shawyer AC, Min S a. L, et al. The prevalence of hearing loss in children with congenital

- diaphragmatic hernia a longitudinal population based study. *Journal of Pediatric Surgery*. 2021;56:90-226.
48. Martínez C, Poblano A, Themann P. Changes in tonal audiometry in children with progressive sensorineural hearing loss and history of Neonatal Intensive Care Unit discharge. *International Journal of Pediatric Otorhinolaryngology*. 2018;101:40-235.
49. Marliyawati D, Handayani P. Hubungan prematuritas dan berat badan lahir rendah dengan derajat gangguan pendengaran pada anak. *Jurnal Penelitian dan Pengembangan Pelayanan Kesehatan*. 2021;5:60.
50. Jeong J, Youk T, Oh J, Eo TS, Choi HS. Neonatal and maternal risk factors for hearing loss in children based on population based data of Korea. *International Journal of Pediatric Otorhinolaryngology*. 2021;147:11-80.
51. Zhu X, Lei X, Dong W. Change to hearing loss-related risks and screening in preterm infants. *American Journal of Perinatology*. 2020;39:12-50.
52. Guven SG. The effect of mode of delivery on newborn hearing screening results. *Turkish Archives of Otorhinolaryngology*. 2019;57:19–23.
53. Gupta N, Raj Rd, Modi P, Satoniya K, Patel Md. Relation between mode of delivery and findings of otoacoustic emission test results. *Journal of Clinical and Diagnostic Research*. 2021;15:1-4.
54. Kelly AF, Kelly P, Shah MD. Auditory brainstem response pass rates correlate with newborn hour of life and delivery mode. *The Journal of Pediatrics*. 2021;230:5-100.
55. Ezzeldin ZM, Sharaf EF, Hamdy M, Selim YAA. Hearing screening in neonates with hyperbilirubinemia. *International Journal of Pediatric Otorhinolaryngology*. 2021;142:110.
56. Culhaoglu B, Erbek SS, İnce DA, Ecevit AN, Erbek S. Medial olfactory complex reflex in term newborns with hyperbilirubinemia. *International Journal of Pediatric Otorhinolaryngology*. 2021;147:77-110.
57. CDC(2022). A parent's guide to genetics and hearing loss. Centers for Disease Control and Prevention. <https://www.cdc.gov/ncbddd/hearingloss/genetics.html>. Diakses September 2023
58. Chiriboga LF, Sideri KP, Figueiredo SNFR, Pinto ESM, Arteta LMC. Outcomes of a universal neonatal hearing screening program of 9941 newborns over a one-year period in Campinas, Brazil. *International Journal of Pediatric Otorhinolaryngology*. 2021;148:110.
59. Escobar-Ipuz FA, Soria-Bretones C, Garcia-Jimenez MA, Cueto E, Aranda AMT, Sotos JM. Early detection of neonatal hearing loss by otoacoustic emissions and auditory brainstem response over 10 years of experience. *International Journal of Pediatric Otorhinolaryngology*. 2019;127:46-97.
60. Wang Y, Qi Z, Yu M, Wang J, Chen R. Characteristic of stimulus frequency otoacoustic emissions: detection rate, musical training influence, and gain function. *Brain Sciences*. 2019;9:255.
61. Dhar S, Hall JW. Otoacoustic emissions principles, procedures, and protocols. Plural Publishing; 2018;2:6-7.
62. Teixeira M, Borges V, Riesgo R, Sleifer P. Hyperbilirubinemia impact on newborn hearing: a literature review. *Revista Da Associacao Medica Brasileira*. 2020;66:1002–8.

63. Martiansya F. Gambaran Faktor Risiko Gangguan Pendengaran Bayi Baru Lahir Hasil Pemeriksaan OAE Refer Di NICU RSUD Raden Mattaher Provinsi Jambi. UNJA. 2022.
64. Chavan R, Ingole S, Damodhar A, Kanchewad G. Hearing Assessment in Infants with Otoacoustic Emission and Auditory Brainstem Response: A Retrospective Study. *Otorhinolaryngology Clinics*. 2021;13(2):29–33.
65. Hajare P, Mudhol R. A Study of JCIH (Joint Commission on Infant Hearing) Risk Factors for Hearing Loss in Babies of NICU and Well Baby Nursery at a Tertiary Care Center. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2022;74:64–90.
66. Raghuwanshi S, Gargava A, Kulkarni V, Kumar A. Role of Otoacoustic Emission Test in Neonatal Screening at Tertiary Center. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2019;71:7–15.
67. Liemiyah R, Marliyati D, Irawan G, Rini AE, Wulansari N, Muyassaroh M. Otoacoustic emission (OAE) features in newborns at Dr. Kariadi Central Genital Hospital, Semarang, Indonesia. *Journal of the Medical Sciences (Berkala Ilmu Kedokteran)*. 2022;54:14.
68. Aradhana A, Sarangi G, Saboth P, Tripathy RM. Assessment of hearing impairment in NICU graduates by otoacoustic emission and brainstem evoked response audiometry tests. *Journal of Neonatology*. 2020;34:109–17.
69. Frezza S, Tiberi E, Corsello M, Priolo F, Cota F, Catenazzi P, et al. Hearing loss and risk factors in very low birth weight infants. *Journal of Clinical Medicine*. 2023;12:75-83.
70. Falerina R, Nugroho PS, Purnami N, Salsabila FA. A comparison of otoacoustic emissions examination results as early hearing screening in normal and infants at risk of hearing loss. *Bali Medical Journal*. 2023;12(2):1640–4.
71. Gulati A, Sakthivel P, Singh I, Ramji S. The hearing status of preterm infant's  $\leq 34$  weeks as revealed by Otoacoustic Emissions (OAE) Screening and Diagnostic Brainstem Evoked Response Audiometry (BERA): a tertiary center experience. *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2020;74:178–83.
72. Primadewi N, Pratiwi D, Hayustiningsih D. Hubungan asfiksia perinatal dengan gangguan fungsi sel rambut luar koklea. *Oto Rhino Laryngologica Indonesiana*. 2019;49:108.
73. Ibrahem SF, Eldin ZE, Hamdy M, Tosson AMS. Auditory assessment of neonates at high risk of hearing impairment admitted to the intensive care unit. *International Journal of Health Sciences*. 2022;11830–46.
74. Fransiska F. Ototoksitas aminoglikosida. Keluwih. 2019;1:37–47.
75. Kaveh M, Mirjalali SN, Shariat M, Zarkesh MR. Perinatal factors influencing the neonatal hearing screening results. *BMC Pediatrics*. 2021;21(1).
76. Chavan R, Ingole S, Damodhar A, Kanchewad G. Hearing Assessment in Infants with Otoacoustic Emission and Auditory Brainstem Response: A Retrospective Study. *Otorhinolaryngology Clinics*. 2021;13(2):29–33.

77. Sari R. Kehamilan dengan Infeksi TORCH. Jurnal Kedokteran Universitas Lampung. 2019;3:1.
78. Masyrukhi N, Widodo P, Muyassaroh M. Faktor risiko infeksi citomegalovirus pada anak yang dicurigai kurang pendengaran. Medica Hospitalia: Journal of Clinical Medicine. 2020;7:53–8.
79. Afshar P, Afsharmanesh J, Eslahi M, Sheikhbardsiri H, Moghadam M. Determination risk factors for severe and profound hearing loss in child candidates for cochlear implantation in southeast of Iran during 2014-2020. BMC Pediatrics. 2022;22:62.
80. Chen D, Lee I, Wang X, Wong SH. Early biomarkers and hearing impairments in patients with neonatal hypoxic ischemic encephalopathy. Diagnostics.2021;11:2056.
81. Aziza M, Bs Y. Hearing impairment in infants with asphyxia. International Tinnitus Journal.2022; 26(1):11-15.

