

DAFTAR PUSTAKA

- Abdallah MA, Yassin MA, Saber N. (2018). Dermoscopic Features of Keloid versus Hypertrophic Scar. *The Egyptian Journal of Hospital Medicine*, 70, 622-624.
- Abd El-Dayem DH, Nada HA, Hanafy NS, Elsaie ML. Laser-assisted topical steroid application versus steroid injection for treating keloids: A split side study. *J Cosmet Dermatol*. 2021 Jan;20(1):138-142
- Aditya MT, Ariyanti PR. (2016). Manfaat Gambir (Uncaria gambir Roxb) sebagai Antioksidan. *Majority*, 5(3), 129-133.
- Alavi A, Kirsner RS. Wound Healing. (2019), In (Kang S, Amagai M, Bruckner AL, Enk AH, Margolis DJ, McMichael AJ, *et al.* editors). *Fitzpatrick's Dermatology in General Medicine 9th ed*. New York: McGraw-Hill, pp 2700-2717.
- Al-hazzani AA, Alshatwi AA. Catechin hydrate inhibits proliferation and mediates apoptosis of SiHa human cervical cancer cells. *FOOD Chem Toxicol*. 2011;49(12):3281-3286.
- Amara N, Goven D, Prost F, Muloway R, Crestani B, Boczkowski J. (2010). NOX4 and NADPH oxidase expression is increased in pulmonary fibroblasts from patients with idiopathic pulmonary fibrosis and mediates TGF- β 1 induced fibroblast differentiation into myofibroblasts. *Thorax*, 65(8), 733-738.
- Amini-Nik S, Yousuf Y, Jeschke MG. (2018). Scar management in burn injuries using drug delivery and molecular signaling: Current treatments and future directions. *Advanced drug delivery reviews*, 123, 135-154.
- Ammendola M, Zuccalà V, Patruno R, Russo E, Luposella M, Amorosi A, *et al.* (2013). Tryptase-positive mast cells and angiogenesis in keloids: a new possible post-surgical target for prevention. *Updates in surgery*, 65(1), 53–57.
- Andasuryani, Purwanto YA, Budiastri IW, Syamsu K. (2014). Prediksi kandungan katekin gambir (Uncaria gambir Roxb.) dengan spektroskopi NIR. *Jurnal Teknologi Industri Pertanian*, 24(1), 43-52.
- Andisi RD, Suling PL, Kapantow MG. (2016). Profil keloid di Poliklinik Kulit dan Kelamin RSUP Prof. Dr. R. D. Kandou Manado periode Januari 2011-Desember 2015. *Jurnal e-Clinic (eCl)*, 4(2).
- Andre N. (2013). A Review of the Occurrence of Non-Alkaloid Constituents in Uncaria Species and Their Structure-Activity Relationships. *American Journal of Biomedical and Life Sciences*, 1(4), 79.

- Anggraini, Tuty T, Akihiro Y, Tomoyuki I, Tomio. (2011). Antioxidative activity and catechin content of four kinds of Uncaria gambir extracts from West Sumatra, Indonesia. *African Journal of Biochemistry Research*, 5, 33-38.
- Anthony ET, Lemonas P, Navsaria HA, Moir GC. (2010). The cost-effectiveness of intralesional steroid therapy for keloids. *Dermatologic surgery: official publication for American Society for Dermatologic Surgery*, 36(10), 1624–1626.
- Arno AI, Gauglitz GG, Barret JP, Jeschke MG. (2014). New molecular medicine-based scar management strategies. *Burns: journal of the International Society for Burn Injuries*, 40(4), 539–51.
- Arima J, Huang C, Rosner B, Akaishi S, Ogawa R. (2015) Hypertension: a systemic key to understanding local keloid severity. *Wound Rep Reg*, 23(3), 213-21.
- Arya N, Kaur D A. (2022). A Review Paper on Application of PCR-ELISA in the Molecular Diagnosis. *International Journal of Innovative Research in Engineering & Management*, (1), 341–344.
- Ashcroft KJ, Syed F, Bayat A. (2013). Site-specific keloid fibroblasts alter the behaviour of normal skin and normal scar fibroblasts through paracrine signaling. *PloS one*, 8(12), e75600.
- Sicard AA, Dao T, Suare NG, Annabi B. (2021). Diet-Derived Gallated Catechins Prevent TGF- β -Mediated Epithelial-Mesenchymal Transition, Cell Migration and Vasculogenic Mimicry in Chemosensitive ES-2 Ovarian Cancer Cells. *Nutrition and Cancer*, 73(1), 169–180.
- Bae J, Kim N, Shin Y, Kim SY, Kim YJ. (2020). Activity of catechins and their applications. *Biomedical Dermatology*, 4(8), 1-10.
- Baranowska M, Suliborska K, Chrzanowski W, Kusznierewicz B, Namieśnik J, Bartoszek A. (2018). The relationship between standard reduction potentials of catechins and biological activities involved in redox control. *Redox Biology*, 17, 355-366.
- Bartosikova L, Necas J. (2018). Epigallocatechin gallate: a review. *Veterinarni Medicina*, 63(10), 443-467.
- Bernatoniene J, Kopustinskiene DM. (2018). The Role of Catechins in Cellular Responses to Oxidative Stress. *Molecules*, 23(4), 965.
- Betarbet U, Blalock TW. (2020). Keloids: a review of etiology, prevention, and treatment. *Journal of Clinical and Aesthetic Dermatology*, 13(2), 33- 43.

- Bhatt LK, Addepalli V. (2012). Potentiation of aspirin-induced cerebroprotection by minocycline: a therapeutic approach to attenuate exacerbation of transient focal cerebral ischaemia. *Diabetes & vascular disease research*, 9(1), 25–34.
- Bian D, Zhang J, Wu X, Dou Y, Yang Y, Tan Q, et al. (2013). Asiatic acid isolated from Centella asiatica inhibits TGF- β 1- induced collagen expression in human keloid fibroblasts via PPAR- γ activation. *International journal of biological sciences*, 9(10), 1032–1042.
- Bijlard E, Kouwenberg CA, Timman R, Hovius SE, Busschbach JJ, Mureau MA. (2017). Burden of Keloid Disease: A Cross-sectional Health-related Quality of Life Assessment. *Acta dermato- venereologica*, 97(2), 225–229.
- Birawati S, Asri E. (2020). Keloids profile in the Dermatovenerology Clinic of RSUP Dr. M. Djamil Padang Indonesia from January 2014 to December 2018. *International Journal of Pharm Tech Research*, 13(04), 383-387.
- Burusapat C, Wanichjaroen N, Wongprakob N, Sapruangthong R. The Effectiveness of Immediate Triamcinolone Acetonide Injection after Auricular Keloid Surgery: A Prospective Randomized Controlled Trial. *Plast Reconstr Surg - Glob Open*. 2021;9(8):E3729.
- Boyanova L, Ilieva J, Gergova G, Vladimirov B, Nikolov R, Mitov I. (2015). Honey and green/black tea consumption may reduce the risk of Helicobacter pylori infection. *Diagnostic microbiology and infectious disease*, 82(1), 85–86.
- Carney B, Chen J, Kent R, Rummani M, Alkhalil A, Moffatt L, et al. (2019) Reactive Oxygen Species Scavenging Potential Contributes to Hypertrophic Scar Formation. *Journal of Surgical Research*, 244, 312-23
- Chen K, Chen W, Liu SL, Wu TS, Yu KF, Qi J, et al. (2018). Epigallocatechin gallate attenuates myocardial injury in a mouse model of heart failure through TGF- β 1/SMAD3 signaling pathway. *Molecular medicine reports*, 17(6), 7652–7660.
- Chen L, Schrementi ME, Ranzer M, Wilgus TA, DiPietro LA. (2014). Blockade of mast cell activation reduces cutaneous scar formation. *PloS one*, 9(1), 1-10.
- Cheresh P, Kim SJ, Tulasiram S, Kamp DW. (2013). Oxidative stress and pulmonary fibrosis. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, 1832(7), 1028-1040.
- Chu KO, Pang CCP. (2018). Pharmacokinetics and Disposition of Green Tea Catechins. In (Ed.), *Pharmacokinetics and Adverse Effects of Drugs-Mechanisms and Risks Factors*. *IntechOpen*, 17-36.

Colen, Bronden E, Geronemus, Roy, McDaniel, David H, *et al.* (2017). The Role of Elastic

Coppola M M, Salzillo R, Segreto F, Persichetti P. (2018). Triamcinolone acetonide intralesional injection for the treatment of keloid scars: Patient selection and perspectives. *Clinical, Cosmetic and Investigational Dermatology*, 11, 387–396. <https://doi.org/10.2147/CCID.S133672>

Damanik DDP, Surbakti N, Hasibuan R. (2014). Ekstraksi katekin dari daun gambir (*Uncaria gambir roxb*) dengan metode maserasi. *Jurnal Teknik Kimia USU*, 3, 10-14.

Da Silva IR, Da Tiveron LCRC, Da Silva MV, Peixoto AB, Carneiro CAX, Dos Reis MA, *et al.* (2017) Ekspresi sitokin in situ dan evaluasi morfometrik total kolagen dan kolagen Tipe I dan Tipe III pada bekas luka Keloid. Peradangan Mediator. 2017(I).

Davies P, Cuttle L, Young A. (2021). A Scoping Review of the Methodology Used in Studies of Genetic Influences on the Development of Keloid or Hypertrophic Scarring in Adults and Children After Acute Wounding. *Advances in wound care*, 10(10), 557–570.

Desdiani D, Rengganis I, Djauzi S, Setiyono A, Sadikin M, Jusman SWA, *et al.* (2020). In vitro assay and study interaction of *Uncaria gambir* (Hunter) Roxb. as anti-fibrotic activity against A549 cell line. *Pharmacognosy Journal*, 12(6), 1232-1240.

Desdiani D, Rengganis I, Djauzi S, Setiyono A, Sadikin M, Jusman S, *et al.* (2022). Fibropreventive and Antifibrotic Effects of *Uncaria gambir* on Rats with Pulmonary Fibrosis. *Evidence-based complementary and alternative medicine: eCAM*, 2022, 6721958.

Díaz-Gómez R, López-Solís RO, Obreque-Slier E, Toledo-Araya H. (2013). Comparative antibacterial effect of gallic acid and catechin against *Helicobacter pylori*. *Lwt - Food Science and Technology*, 54, 331-335.

Di Leo N, Battaglini M, Berger L, Giannaccini M, Dente L, Hampel S, *et al.* (2017). A catechin nanoformulation inhibits WM266 melanoma cell proliferation, migration, and associated neo-angiogenesis. *European Journal of Pharmaceutics and Biopharmaceutics*, 114, 1–10.

Distler JH, Györfi A, Ramanujam M, Whitfield ML, Königshoff M, Lafyatis R. (2019). Shared and distinct mechanisms of fibrosis. *Nature Reviews Rheumatology*, 15, 705-730

Ediningsih, Rahayuningsih S. (2019). Ekstraksi, Isolasi, Karakterisas dan Uji Aktivitas Antioksidan Senyawa Katekin Gambir (*Uncaria gambir* (Hunter)). *Roxb. Alkimia*, 7(2), 177–88.

- Ekstein SF, Wyles SP, Moran SL, Meves A. (2021). Keloids: a review of therapeutic management. *International journal of dermatology*, 60(6), 661–671.
- Elazhary E, Abd Al-Salam F, Abd El-Hafiz H, Maghraby H. (2022). Updates on keloid scar pathogenesis, assessment, and treatment modalities. *Journal of Recent Advances in Medicine*, 3(1), 75-86.
- Eugene EA, Robert NB, Ganiyou A, Denis YK, Ane A, Sawaliho BEH. (2022). Catechin and Epicatechin. What the more reactive? *Computational chemistry*, 10,53-70
- Fahmi K, Rahimullaily R. (2022). Strategi Pengembangan Usaha Gambir Rakyat Menggunakan Analisa SWOT di Kabupaten Pesisir Selatan. *SALAM: Jurnal Sosial Dan Budaya Syar-i*, 9(5), 1683–1698.
- Fahrudin F, Solihin DD, Kusumorini N, Ningsih S. Isolasi efektifitas ekstrakgambir (*uncaria gambir (hunter) roxburgh*) sebagai hepatoprotektor pada Tikus (*Rattus norvegicus L.*) yang diinduksi CCl₄. *Jurnal ilmu kefarmasian indonesia*.2015;13(2):115- 22
- Farah S, Zahida R, Bushra B, Faria A, Khurshid K. (2017). Comparison of efficacy of intralesional 5-fluorouracil plus triamcinolone acetonide versus intralesional triamcinolone acetonide in the treatment of keloids. *Journal of Pakistan Association of Dermatologists*, 27(2):114-119.
- Finnerty CC, Jeschke MG, Branski LK, Barret JP, Dziewulski P, Herndon DN. (2016). Hypertrophic scarring: the greatest unmet challenge after burn injury. *Lancet (London, England)*, 388(10052), 1427– 1436.
- Finnson KW, McLean S, Di Guglielmo GM, Philip A. (2013). Dynamics of Transforming Growth Factor Beta Signaling in Wound Healing and Scarring. *Advances in wound care*, 2(5),195–214.
- Frinanda D, Efrizal R, Rahayu. (2014). Efektivitas Gambir (*Uncaria gambir Roxb.*) sebagai Anti Hipercolesterolemia dan Stabilisator Nilai Darah pada Mencit Putih (*Mus musculus*) Jantan. *Jurnal Biologi Universitas Andalas*, 3(3): 231-237
- Gadkari PV, Balaraman M. (2014). Catechins: sources, extraction and encapsulation: a review. *Food and Bioproducts Processing*, 93, 122-138.
- Gauglitz GG, Korting HC, Pavicic T, Ruzicka T, Jeschke MG. (2011). Hypertrophic scarring and keloids: pathomechanisms and current and emerging treatment strategies. *Molecular medicine (Cambridge, Mass.)*, 17(1-2), 113–125.

Gauglitz GG. (2013). Management of keloids and hypertrophic scars: current and emerging options. *Clinical, cosmetic, and investigational dermatology*, 6, 103–114.

Ghitescu RE, Popa AM, Schipanski A, Hirsch C, Yazgan G, Popa VI, et al. (2018). Catechin-loaded PLGA submicron-sized fibers reduce levels of reactive oxygen species induced by MWCNT in vitro. *European journal of pharmaceutics and biopharmaceutics: official journal of Arbeitsgemeinschaft fur Pharmazeutische Verfahrenstechnik e.V.*, 122, 78–86.

Gomes FMS, da Cunha XJ, dos Santos JFS, de Matos YMLS, Tintino SR, de Freitas TS, et al. (2018). Evaluation of antibacterial and modifying action of catechin antibiotics in resistant strains. *Microbial Pathogenesis*, 115, 175–178.

Hilmi HL, Rahayu D. (2018). Artikel Tinjauan: Aktivitas Farmakologi Gambir (Uncaria gambir Roxb.). *Farmaka Suplemen*, 16(2), 134-141.

Hosen, N. (2017). Profil Sistem Usaha Pertanian Gambir di Sumatera Barat. *Jurnal Penelitian Pertanian Terapan*, 17(2), 124–131.

Huang C, Wu Z, Du Y, Ogawa R(1). (2020). The Epidemiology of Keloids.Dalam: Téot L, Mustoe TA, Middelkoop E, Gauglit GG editor. Textbook on Scar Management. Switzerland: Springer. pp. 29-34.

Hsiao CC, Chang YC, Hsiao YT, Chen PH, Hsieh MC, Wu WC, et al. Triamcinolone acetonide modulates TGF- β 2-induced angiogenic and tissue -remodeling effects in cultured human retinal pigment epithelial cells. *Mol Med Rep [Internet]*. 2021 Sep 14;24(5):802.

Husein SSS, Kamarudin MNA, Kadir HA. (+)-Catechin attenuates NF- κ B activation through regulation of Akt, MAPK, and AMPK signaling pathways in LPS-induced BV-2 microglial cells. *The American Journal of Chinese Medicine*. 2015;43(5);1-26

Isemura M. (2019). Catechin in Human Health and Disease. *Molecules (Basel, Switzerland)*, 24(3), 528.

Isnawati A, Raini M, Sampurno OD, MutiatikumD, Widowati L, Gitawati R. (2012). Karakterisasi tiga jenis ekstrak gambir (Uncaria gambir Roxb) dari Sumatera Barat. *Buletin Penelitian Kesehatan*, 40(4), 201 – 208.

Jin Q, Gui L, Niu F, Yu B, Lauda N, Liu J, et al. (2018). Macrophages in keloid are potent at promoting the differentiation and function of regulatory T cells. *Experimental cell research*, 362(2), 472– 476.

- Jusman SWA, Amalia MF, Paramita R, Ningsih S, Fadillah F. (2023). Structure-based screening of inhibitor platelet-derived growth factor from ethanol extract of Uncaria gambir (Hunter) Roxb. as an antifibrotic in keloid fibroblast cells. *Journal of Applied Pharmaceutical Science* Vol. 13(03), pp 036-043
- Komminoth P, Heitz PU, Long AA. (1994). In situ polymerase chain reaction: general methodology and recent advances. *Verhandlungen der Deutschen Gesellschaft fur Pathologie*, 78, 146–152.
- Kamsina K, Firdausni, F. (2018). Pengaruh penggunaan ekstrak gambir sebagai antimikroba terhadap mutu dan ketahanan simpan cake bengkuang (*Pachyrhizus erosus*). *Jurnal Litbang Industri*, 8(2), 111-117.
- Kang JS, Park IH, Cho JS, Hong SM, Kim TH, Lee SH, Lee H. M. (2014). Epigallocatechin-3-gallate inhibits collagen production of nasal polyp-derived fibroblasts. *Phytotherapy Research*, 28(1), 98–103. <https://doi.org/10.1002/ptr.4971>
- Kassi K, Kouame K, Kouassi A, Allou A, Kouassi I, Kourouma S, et al. (2020). Quality of life in black African patients with keloid scars. *Dermatology reports*, 12(2), 8312.
- Kenny FN, Drymoussi Z, Delaine-smith R, Kao AP, Laly AC, Knight MM, et al. (2018). Tissue stiffening promotes keratinocyte proliferation through activation of epidermal growth factor signaling, 131, 1-10.
- Khan M, Mohammad A, Patil G, Naqvi S, Chaucan L, Ahmad I. (2012). Induction of ROS, mitochondrial damage, and autophagy in lung epithelial cancer cells by iron oxide nanoparticles. 33(5), 1477-88
- Khansa I, Harrison B, Janis JE. (2016). Evidence-based scar management. *Plast Reconstr Surg*, 138, 165S–78S.
- Kim SW. (2021). Management of keloid scars: noninvasive and invasive treatments. *Arch Plast Surg*, 48(2), 149-15.
- Kimura-Ohba S, Yang Y. (2016). Oxidative DNA Damage Mediated by Intranuclear MMP Activity Is Associated with Neuronal Apoptosis in Ischemic Stroke. *Oxidative Medicine and Cellular Longevity*. Hindawi Limited.
- Kilmister EJ, Paterson C, Brasch HD, Davis PF, Tan ST. (2019). The role of the renin-angiotensin system and vitamin D in keloid disorder - a review. *Frontiers in Surgery*.6:1-12.

Klomparens K, Simman R. (2022). Treatment of Keloids: A Meta-analysis of Intralesional Triamcinolone, Verapamil, and Their Combination. *Plastic and reconstructive surgery. Global open*, 10(1), e4075.

Ko H, So Y, Jeon H, Jeong MH, Choi H, Ryu S. (2013). TGF- β 1-induced epithelial-mesenchymal transition and acetylation of Smad2 and Smad3 are negatively regulated by EGCG in Human A549 lung cancer cells, 335(1), 205-213

Ko H, So Y, Jeon H, Jeong MH, Choi HK, Ryu SH, Lee SW, Yoon HG, Choi KC.(2013). TGF- β 1-induced epithelial-mesenchymal transition and acetylation of Smad2 and Smad3 are negatively regulated by EGCG in human A549 lung cancer cells. *Cancer Lett.* 10;335(1):205-13.

Kordestani SS. (2019). *Atlas of Wound Healing: A Tissue Regeneration Approach*, Elsevier, pp 23-50.

Kumar P, Nagarajan A, Uchil PD. Analysis of cell viability by the MTT assay. *Cold Spring Harb Protoc.* 2018:469-472

Kurniawati Y, Adi S, Achadiyani, Suwarsa O, Erlangga D, Putri T. Kultur primer fibroblas: penelitian pendahuluan. *Majalah Kedokteran Andalas.* 2015; 38:33-40.

Kurniatri AA, Sulistyaningrum N, Rustanti L. (2019). Purifikasi Katekin dari Ekstrak Gambir (Uncaria gambir Roxb). *Media Penelitian dan Pengembangan Kesehatan*, 29(2), 153-160.

Koutroutsos K, et al. (2014) Effect of Smad pathway activation on podocyte cell cycle regulation: an immunohistochemical evaluation. *Ren. Fail.* 36:1310–6.

Kuwahara H, Tosa M, Egawa S, Murakami M, Mohammad G, Ogawa R. (2016). Examination of Epithelial-Mesenchymal Transition in Keloid Tissues and Possibility of Keloid Therapy Target. *Plastic and reconstructive surgery. Global open*, 4(11), 1-7.

Lee HJ, Jang YJ. (2018). Recent understandings of biology, prophylaxis, and treatment strategies for hypertrophic scars and keloids. *International journal of molecular sciences*, 19(3), 711.

Lebeko M, Khumalo NP, Bayat A. (2019). Multi-dimensional models for functional testing of keloid scars: in silico, in vitro, organoid, organotypic, ex vivo organ culture, and in vivo models. *Wound Repair and Regeneration*.27, 298-308.

- Li F, Jin H, Xiao J, Yin X, Liu X, Li D, *et al.* (2018) The simultaneous loading of catechin and quercetin on chitosan-based nanoparticles as effective antioxidant and antibacterial agent. *Food Research International*, 111, 351-360.
- Li T, Zhao N, Lu J, Zhu Q, Liu X, Hao F, Jiao X. (2019). Epigallocatechin gallate (EGCG) suppresses epithelial-mesenchymal transition (EMT) and invasion in anaplastic thyroid carcinoma cells through blocking of TGF- β 1/Smad signaling pathways. *Bioengineered*, 10(1), 282–291. <https://doi.org/10.1080/21655979.2019.1632669>
- Limandjaja GC, van den Broek LJ, Waaijman T, Breetveld M, Monstrey S, Schepers RJ, *et al.* (2018). Reconstructed human keloid models show heterogeneity within keloid scars. *Arch Dermatol Res*, 310(10), 815–26.
- Limandjaja GC, Niessen FB, Schepers RJ, Gibbs S. (2020) The Keloid Disorder: Heterogeneity, Histopathology, Mechanisms and Models. *Front. Cell Dev. Biol*, 8, 360.
- Limandjaja GC, Niessen FB, Schepers RJ, Gibbs S. (2021). Hypertrophic scars and keloids: Overview of the evidence and practical guide for differentiating between these abnormal scars. John Wiley & Sons, 30, 146–161.
- Lucarini M, Sciubba F, Capitani D, Di Cocco ME, D’Evoli L, Durazzo, A *et al* (2020). Role of catechin on collagen type I stability upon oxidation: A NMR approach. *Natural product research*, 34(1), 53-62.
- Lu Y, Wu C, Hong C, Chang K, Lee C. (2021) GLUT-1 Enhances Glycolysis, Oxidative Stress, and Fibroblast Proliferation in Keloid. Multidisciplinary Digital Publishing Institute. 11, 505, 1-18
- Lu Y, Zhang S, Wang Y, Ren X, Han J. (2019). Molecular mechanisms and clinical manifestations of rare genetic disorders associated with type I collagen. *Intractable and Rare Diseases Research*, 8(2), 98–107.
- Lysy PA, Smets F, Sibille C, Najimi M, Sokal EM. (2007). Human skin fibroblasts: From mesodermal to hepatocyte-like differentiation. *Hepatology*, 46(5):1574- 85.
- Machan S, Ruiz A RL. (2018). Dermal hypertrophy in: Bolognia J, Schaffer J, Cerroni L, editors. *Dermatology* 4th edition. New York: Elsevier. 2018:1712- 1718.
- Maiti R, Agrawal N, Dash D, Pandey B. (2007). Effect of Pentoxifylline on inflammatory burden, oxidative stress, and platelet aggregability in hypertensive type 2 diabetes mellitus patients. *Vascular Pharmacology*. 47(2),118-24

- Mangels DR, Mohler ER. (2017). Catechins as Potential Mediators of Cardiovascular Health. *Arterioscler Thromb Vasc Biol*, 37(5), 757-763.
- Mangis JW, Mansur TB, Kern KM, Schroeder JR. Selection of an optimal cytotoxicity assay for undergraduate research. Department of Biology. 2019;45(1):24-32.
- Marlinda. (2018). Identifikasi kadar katekin pada gambir (Uncaria Gambier Roxb). *Jurnal Optimalisasi*, 4(1), 47-53.
- Matsuzaki S, Darcha C. (2014). Antifibrotic properties of epigallocatechin-3-gallate in endometriosis. *Hum Reprod*, 29(8):1677-87.
- Mishra B, Arora C. (2020). Epidemiology of keloids and hypertrophic scars in a Tertiary Care Teaching Hospital of Northern India. *International Journal of Scientific Research*, 10(2), 54-55.
- Nagar H, Kim S, Lee I, Kim S, Choi SJ, Piao S, et al. (2021) Downregulation of CR6-interacting factor 1 suppresses keloid fibroblast growth via the TGF- β /SMAD signaling pathway. *Sci Rep*, 11(1), 500.
- Nangole FW, Agak GW. (2019). Keloid pathophysiology: fibroblast or inflammatory disorders? *JPRAS Open*, 22, 44-54.
- Naveeda M, BiBi J, Kamboh AA, Suherjani I, Kakar I, Fazlani SA, et al. (2018). Pharmacological values and therapeutic properties of black tea (*Camellia sinensis*): A comprehensive overview, *Biomedicine & Pharmacotherapy*, 100, 521-531.
- Ningsih, S. Fachrudin F, Rismana E, Purwaningsih E, Sumaryono W, Jusman S. (2014). Evaluation of anti-lipid peroxidation activity of gambir extract on liver homogenate in vitro. *Int.J. PharmTech Res*, 6(3), 982-989.
- Nordin A, Chowdhury SR, Saim AB, Bt R, Idrus R. (2020). Effect of kelulut honey on the cellular dynamics of TGF β -induced epithelial to mesenchymal transition in primary human keratinocytes. *International Journal of Environmental Research and Public Health*, 17, 1-15.
- Ogawa R, Okai K, Tokumura F, Mori K, Ohmori Y, Huang C, et al. (2012). The relationship between skin stretching/contraction and pathologic scarring: the important role of mechanical forces in keloid generation. *Wound Repair Regen*, 20(2), 149–57.
- Ogawa R. (2017). Keloid and hypertrophic scars are the result of chronic inflammation in the reticular dermis. *Int J Mol Sci*, 18(3), 1–10.
- Ogawa R. (2020). Total Scar Management. Ogawa R, editor. Springer.

- Ogawa R, Akaishi S. (2016). Endothelial dysfunction may play a key role in keloid and hypertrophic scar pathogenesis—keloids and hypertrophic scars may be vascular disorders. *Medical Hypotheses*, 96:51-60.
- Ojeh N, Bharatha A, Gaur U, Forde AL. (2020). Keloids: Current and emerging therapies. *Scars, Burn Heal.* 6, 1–18.
- Pan Z, Zhoua Y, Luoa X, Ruanc Y, Zhoua L, Wangc Q, et al. (2018). Against NF- κ B/thymic stromal lymphopoietin signaling pathway, catechin alleviates the inflammation in allergic rhinitis. *International Immunopharmacology*, 61, 241-248.
- Präbst K, Engelhardt H, Ringgeler S. Basic colorimetric proliferation assays: mtt, wst, and resazurin konstantin. In: Daniel F. Gilbert, Oliver Friedrich, editors. Cell viability assays: methods and protocols, methods in molecular biology. Jerman: Springer Science & Business Media. 2017:1-17
- Pellard S. (2014). Epidemiology, etiology, and management of abnormal scarring: a review of the literature. *J Wound Care*, 15(1), 44-48.
- Putra IB, Soebono H, Hadisahputra S, Bachtiar A. (2020). The effect of roselle (*Hibiscus sabdariffa* L.) flowers extract on the apoptosis of fibroblast proliferation in keloids. *Open Access Macedonian Journal of Medical Sciences*, 8, 611-614.
- Rahmawati N, Bakhtiar A, Putra DP. (2012). Isolasi katekin dari gambir (*Uncaria gambir* (Hunter) Roxb) untuk sediaan farmasi dan kosmetik. *Jurnal Penelitian Farmasi Indonesia*, 1, 6-10.
- Ramakrishnan MK, Babu M, Mathivanan T, Jayaraman V. (2014). Scar biology: fibroblast the key modulator. *Indian Journal of Burn*, 22(1), 3-9.
- Rashidinejad A, Boostani S, Babazadeh A, Rehman A, Rezaei A, Akbari-Alavijeh S, et al. (2021). Opportunities and challenges for the nano delivery of green tea catechins in functional foods. *Food Res Int*, 142, 110186.
- Ravipati AS, Reddy N, Kooyalamudi SR. (2014). Biologically active compounds from the genus *Uncaria* (Rubiaceae). 1st ed. Vol. 43, Studies in Natural Products Chemistry. Elsevier B.V, pp 381-408.
- Saad M, Goh H, Rajikan R, Roslina T, Baharum S, Bunawan H. (2020). *Uncaria gambir* (W. Hunter) Roxb: From phytochemical composition to pharmacological importance. *Tropical Journal of Pharmaceutical Research*, 19(8), 1767-1773.
- Sabarni. (2015). Teknik Pembuatan Gambir (*Uncaria gambir* Roxb) Secara Tradisional. *Journal of Islamic Science and Technology*, 1(1), 105-112.

- Santos LFS, Stolfo A, Calloni C, Salvador M. (2017). Catechin and epicatechin reduce mitochondrial dysfunction and oxidative stress induced by amiodarone in human lung fibroblasts. *Journal of Arrhythmia*, 33, 220-225.
- Saputra, R. (2017). Pengembangan Sumber Daya Lokal di Kabupaten Lima Puluh Kota Provinsi Sumatera Barat (Studi Kasus: Pengelolaan Gambir di Kecamatan Pangkalan Koto Baru). *Jom VISIP*, 4(2), 1-13.
- Sari II, Anum Q, Asri E. (2021). Pengaruh Katekin Gambir (*Uncaria Gambir Roxburgh*) Terhadap Proliferasi dan Migrasi Sel Fibroblas Keloid. Tesis. Fakultas Kedokteran Universitas Andalas.
- Satish L, Evdokiou A, Geletu E, Hahn JM, Supp DM. (2020). Pirfenidone inhibits epithelial-mesenchymal transition in keloid keratinocytes. *Burns & Trauma*, 8, 1-11.
- Sebayang, L. (2014). Inovasi Teknologi Gambir di Pakpak Barat. Balai Pengkajian Teknologi Pertanian Sumatera Utara. Medan.
- Shaheen A. (2017). Comprehensive review of keloid formation. *Clin Res Dermatol Open Access*, 4(5), 1-18.
- Shaheen AA. (2017). Risk Factors of Keloids: Mini review. *Austin J Dermatolog*, 4(2), 1074.
- Sidgwick GP, McGeorge D, Bayat A. (2016). Functional testing of topical skin formulations using an optimized ex vivo skin organ culture model. *Archives of dermatological research*, 308(5), 297–308.
- Shahid A, Ali R, Ali N, Hasan SK, Bernwal P, Afzal SM, Vafa A, Sultana S. (2016). Modulatory effects of catechin hydrate against genotoxicity, oxidative stress, inflammation, and apoptosis induced by benzo(a)pyrene in mice. *Food and Chemical Toxicology*, 92, 64 -74.
- Sinto L. (2018). Skar Hipertrofik dan Keloid: Patofisiologi dan Penatalaksanaan CDK, 45(1), 29-32.
- Sharma JR, Lebeko M, Kidzeru EB, Khumalo NP, Bayat A. (2019). In vitro and ex vivo models for functional testing of therapeutic anti-scarring drug targets in keloids. *Advances In Wound Care*, 8(12), 655-670.
- Shih B, Garside E, McGrouther DA, Bayat A. (2010). Molecular dissection of abnormal wound healing processes resulting in keloid disease. *Wound Repair and Regeneration*, 139-153.
- Song Y, Wang T, Yang L, Wu J, Chen L, Fan X, et.al (2023). EGCG inhibits hypertrophic scar formation in a rabbit ear model. *Journal of cosmetic dermatology*, 22(4), 1382-1391.

- Stewart AG, Thomas B, Koff J. (2018). TGF- β : master regulator of inflammation and fibrosis. *Respirology*, 23(12), 1096–1097.
- Sun LM, Wang KH, Lee YCG. (2014). Keloid incidence in Asian people and its comorbidity with other fibrosis-related diseases: a nationwide population-based study. *Arch Dermatol Res*, 306(9), 803–8.
- Suryavanshi SV, Kulkarni YA. (2017). NF- κ B: a potential target in the management of vascular complications of diabetes, *Front. Pharmacol*, 8, 1-12.
- Sutheno, A. (2021). Keloid after orthopedic surgery: Prevention, current therapy modalities, and emerging therapies modalities. *Bali Medical Journal*, 10(1), 225–228.
- Syed F, Bagabir RA, Paus R, Bayat A. (2013). Ex vivo evaluation of antifibrotic compounds in skin scarring: EGCG and silencing of PAI-1 independently inhibit growth and induce keloid shrinkage. *Lab Invest*, 93, 946–60.
- Thornton NJ, Garcia BA, Hoyer P, and Wilkerson MG. (2021). Keloid scars: an updated review of combination therapies. *Cureus*, 13 (1), e12999.
- Trace AP, Enos CW, Mantel A, Harvey VM. (2016). Keloids and Hypertrophic Scars: A Spectrum of Clinical Challenges. *American journal of clinical dermatology*, 17(3), 201–223.
- Treviño-Villarreal JH, Reynolds JS, Langston PK, Thompson A, Mitchell JR, Franco RA. (2021). Down-Regulation of a Profibrotic Transforming Growth Factor- β 1/Cellular Communication Network Factor 2/Matrix Metalloprotease 9 Axis by Triamcinolone Improves Idiopathic Subglottic Stenosis. *The American Journal of Pathology*, 191(8), 1412– 1430. <https://doi.org/10.1016/j.ajpath.2021.05.013>
- Tsai CH, Ogawa R. (2019). Keloid research: status and future directions. *Scars, burns & healing*, 5, 1-8.
- Udarno LRT, Setiyono. (2013). Biologi Bunga Dua Varietas Gambir (Uncaria gambir Hunter Roxb.) di Kebun Pakuwon. *Jurnal Sirinoov*, 1(2), 83–88.
- Ud-din S, Foden P, Mazhari M, Al-Habba S, Baguneid M, Paus SB, et al. (2019). A double-blind, randomized trial shows the role of zonal priming and direct topical application of epigallocatechin-3-gallate in the modulation of cutaneous scarring in human skin. *Journal of Investigative Dermatology*, 139, 1680-1690.
- Velazquez SP, Mendoza FA, Jimenez SA. (2016). Endothelial to mesenchymal transition (EndoMT) in the pathogenesis of human fibrotic diseases. *Journal of Clinical Medicine*, 1-22.

- Wahyuningsih, Mae & Yuliani, Fara & Rahmawati, Dwiki & Pratiwi, Annisa. (2018). Antifibrotic Effect of Ethanolic Extract of Nerium indicum Mill. Standardized 5 α -Oleandrin on Keloid Fibroblasts Cells. *Majalah Obat Tradisional*, 23(1), 70.
- Walliczek U, Engel S, Weiss C, Aderhold C, Lippert C, Wenzel A, et al. (2015). Clinical Outcome and Quality of Life After a Multimodal Therapy Approach to Ear Keloids. *JAMA Facial Plast Surg*, 17(5), 333-9.
- Walton KL, Johnson KE, Harrison CA. (2017). Targeting TGF- β Mediated SMAD Signaling for the Prevention of Fibrosis. *Front Pharmacol*, 8, 461.
- Wang Y, Liu N, Su X, Zhou G, Sun G, Du F, Wang, B. (2015). Epigallocatechin-3-gallate attenuates transforming growth factor- β 1-induced epithelial-mesenchymal transition via Nrf2 regulation in renal tubular epithelial cells. *Biomedicine and Pharmacotherapy*, 70(C), 260–267. <https://doi.org/10.1016/j.biopha.2015.01.032>
- Wang Z, Gao Z, Shi Y, Sun Y, Lin Z, Jiang H, et al. (2007). Inhibition of SMAD3 expression decreases collagen synthesis in keloid disease fibroblasts. *J Plast Reconstr Aesthet Surg*, 60(11):1193-9.
- Wang ZC, Zhao WY, Cao Y, Liu YQ, Sun, Q, Shi P, et al. (2020). The Roles of Inflammation in Keloid and Hypertrophic Scars. *Frontiers in immunology*, 11, 603187.
- Wardani FA, Perdanakusuma DS, Indramaya DM. (2021). Profile of Working-Age Patients with Keloid and Hypertrophic Scar at Dr. Soetomo General Hospital Surabaya in 2014-2017.
- Widaty S, Soebono H, Nilasari H, Listiawan Y, Siswati AS, Triwahyudi D, et al. (2017). Keloid. In: Panduan Praktik Klinis Bagi Dokter Spesialis Kulit dan Kelamin di Indonesia. Jakarta: PP Perdoski, p. 317.
- Wilgus TA. (2020). Inflammation as an orchestrator of cutaneous scar formation: a review of the literature. *Plastic and Aesthetic Research*, 7(54), 1-18.
- Xu X, Gu S, Huang X, Ren J, Gu Y, Wei C, et al. (2020). The role of macrophages in the formation of hypertrophic scars and keloids. *Burns&Trauma*, (8), 1-8.
- Xue M, Jackson CJ. (2018). Extracellular matrix reorganization during wound healing and its impact on abnormal scarring. *Advances in wound care*, 4(3):119-136.

Yang, T. H., Gingery, A., Thoreson, A. R., Larson, D. R., Zhao, C., & Amadio, P.

C. (2018). Triamcinolone Acetonide affects TGF- β signaling regulation of fibrosis in idiopathic carpal tunnel syndrome. *BMC Musculoskeletal Disorders*, 19(1), 1–9. <https://doi.org/10.1186/s12891-018-2260-y>

Yoo MG, Kim IH. (2014). Keloids and hypertrophic scars: characteristic vascular structures visualized by using dermoscopy. *Annals of Dermatology*, 26(5), 603-609.

You HL, Huang CC, Chen CJ, Chang CC, Liao PL, Huang ST. (2018). Anti-pandemic influenza A (H1N1) virus potential of catechin and gallic acid. *Journal of the Chinese Medical Association*, 81, 458-468.

Zhang Q, Kelly A P, Wang L, French SW, Tang X, Duong HS, et al. (2006). Green tea extract and (-)-epigallocatechin-3-gallate inhibit mast cell-stimulated type I collagen expression in keloid fibroblasts via blocking PI-3K/Akt signaling pathways. *Journal of Investigative Dermatology*, 126(12), 2607– 2613.

Zhang T, Wang XF, Wang ZC, Lou D, Fang QQ, Hu YY, et al. (2020). Current potential therapeutic strategies targeting the TGF- β /SMAD signaling pathway to attenuate keloid and hypertrophic scar formation. *Biomed Pharmacother*, 129, 1-9.

Zhu, Qiangqiang, Xiao-ying, Yang, Xiaojuan, Zhang et al. (2020). EGCG targeting Notch to attenuate renal fibrosis via inhibition of TGF β /Smad3 signaling pathway activation in streptozotocin-induced diabetic mice. *Food & Function*, 11(11), 9686-9695. Available from: 10.1039/DFO01542C

Zink A, Traidl-Hoffmann C. (2015). Green tea in dermatology--myths and facts. *Journal der Deutschen Dermatologischen Gesellschaft. Journal of the German Society of Dermatology: JDDG*, 13(8), 768–775.