

## DAFTAR PUSTAKA

- Alam, M. Z., Alhebsi, M. S. R., Ghnimi, S., and Kamal-Eldin, A. (2021). Inability of total antioxidant activity assays to accurately assess the phenolic compounds of date palm fruit (*Phoenix dactylifera* L.). *NFS Journal*, 22, 32–40. <https://doi.org/https://doi.org/10.1016/j.nfs.2021.01.001>
- Alañón, M. E., Ivanović, M., Pimentel-Mora, S., Borrás-Linares, I., Arráez-Román, D., and Segura-Carretero, A. (2020). A novel sustainable approach for the extraction of value-added compounds from *Hibiscus sabdariffa* L. calyces by natural deep eutectic solvents. *Food Research International*, 137, 109646. <https://doi.org/https://doi.org/10.1016/j.foodres.2020.109646>
- Ansar, Nazaruddin, and Azis, A. D. (2020). New frozen product development from strawberries (*Fragaria Ananassa* Duch.). *Heliyon*, 6(9), e05118. <https://doi.org/https://doi.org/10.1016/j.heliyon.2020.e05118>
- Badan Pusat Statistik. (2023). *Produksi Tanaman Buah-Buahan*. <https://www.bps.go.id/Id/Statistics-Table/2/NjIjMg==/Produksi-Tanaman-Buah-Buahan.Html>.
- Barbieri, J. B., Goltz, C., Batistão Cavalheiro, F., Theodoro Toci, A., Igarashi-Mafra, L., and Mafra, M. R. (2020). Deep eutectic solvents applied in the extraction and stabilization of rosemary (*Rosmarinus officinalis* L.) phenolic compounds. *Industrial Crops and Products*, 144, 112049. <https://doi.org/https://doi.org/10.1016/j.indcrop.2019.112049>
- Benvenuti, L., Zielinski, A. A. F., and Ferreira, S. R. S. (2022a). Pressurized aqueous solutions of deep eutectic solvent (DES): A green emergent extraction of anthocyanins from a Brazilian berry processing by-product. *Food Chemistry*, X, 13, 100236. <https://doi.org/https://doi.org/10.1016/j.fochx.2022.100236>
- Benvenuti, L., Zielinski, A. A. F., and Ferreira, S. R. S. (2022b). Pressurized aqueous solutions of deep eutectic solvent (DES): A green emergent extraction of anthocyanins from a Brazilian berry processing by-product. *Food Chemistry*, X, 13, 100236. <https://doi.org/https://doi.org/10.1016/j.fochx.2022.100236>
- Bi, Y., Chi, X., Zhang, R., Lu, Y., Wang, Z., Dong, Q., Ding, C., Yang, R., and Jiang, L. (2020). Highly efficient extraction of mulberry anthocyanins in deep eutectic solvents: Insights of degradation kinetics and stability evaluation. *Innovative Food Science and Emerging Technologies*, 66, 102512. <https://doi.org/https://doi.org/10.1016/j.ifset.2020.102512>
- Canuto, G. A. B., Oliveira, D. R., da Conceição, L. S. M., Farah, J. P. S., and Tavares, M. F. M. (2016). Development and validation of a liquid chromatography method for anthocyanins in strawberry (*Fragaria* spp.) and complementary studies on stability, kinetics and antioxidant power. *Food Chemistry*, 192, 566–574.
- Comuzzo, P., and Battistutta, F. (2019). Chapter 2 - Acidification and pH Control in Red Wines. In A. Morata (Ed.), *Red Wine Technology* (pp. 17–34). Academic Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-814399-5.00002-5>

- Cuéllar-Villarreal, M. del R., Ortega-Hernández, E., Becerra-Moreno, A., Welti-Chanes, J., Cisneros-Zevallos, L., and Jacobo-Velázquez, D. A. (2016). Effects of ultrasound treatment and storage time on the extractability and biosynthesis of nutraceuticals in carrot (*Daucus carota*). *Postharvest Biology and Technology*, 119, 18–26. <https://doi.org/https://doi.org/10.1016/j.postharvbio.2016.04.013>
- Cvjetko Bubalo, M., Ćurko, N., Tomašević, M., Kovačević Ganić, K., and Radojčić Redovniković, I. (2016). Green extraction of grape skin phenolics by using deep eutectic solvents. *Food Chemistry*, 200, 159–166. <https://doi.org/https://doi.org/10.1016/j.foodchem.2016.01.040>
- Dai, Y., Rozema, E., Verpoorte, R., and Choi, Y. H. (2016a). Application of natural deep eutectic solvents to the extraction of anthocyanins from *Catharanthus roseus* with high extractability and stability replacing conventional organic solvents. *Journal of Chromatography A*, 1434, 50–56. <https://doi.org/https://doi.org/10.1016/j.chroma.2016.01.037>
- Dai, Y., Rozema, E., Verpoorte, R., and Choi, Y. H. (2016b). Application of natural deep eutectic solvents to the extraction of anthocyanins from *Catharanthus roseus* with high extractability and stability replacing conventional organic solvents. *Journal of Chromatography A*, 1434, 50–56. <https://doi.org/https://doi.org/10.1016/j.chroma.2016.01.037>
- Della Posta, S., Gallo, V., Ascrizzi, A. M., Gentili, A., De Gara, L., Dugo, L., and Fanali, C. (2023). Development of a green ultrasound-assisted procedure for the extraction of phenolic compounds from avocado peel with deep eutectic solvents. *Green Analytical Chemistry*, 7, 100083. <https://doi.org/https://doi.org/10.1016/j.greeac.2023.100083>
- Dzhanfezova, T., Barba-Espín, G., Müller, R., Joernsgaard, B., Hegelund, J. N., Madsen, B., Larsen, D. H., Martínez Vega, M., and Toldam-Andersen, T. B. (2020). Anthocyanin profile, antioxidant activity and total phenolic content of a strawberry (*Fragaria × ananassa* Duch) genetic resource collection. *Food Bioscience*, 36, 100620. <https://doi.org/https://doi.org/10.1016/j.fbio.2020.100620>
- El Achkar, T., Fourmentin, S., and Greige-Gerges, H. (2019). Deep eutectic solvents: An overview on their interactions with water and biochemical compounds. *Journal of Molecular Liquids*, 288, 111028. <https://doi.org/https://doi.org/10.1016/j.molliq.2019.111028>
- Fajriani, L. A. (2023). *Pengaruh Penambahan Magnesium Karbonat (MgCO<sub>3</sub>) Terhadap Karakteristik Mutu Bubuk Pewarna Alami dari Labu Kabocha (Cucurbitamaxima. L.)* [Skripsi]. Universitas Andalas.
- Farias, C. A. A., Moraes, D. P., Neuenfeldt, N. H., Zabet, G. L., Emanuelli, T., Barin, J. S., Ballus, C. A., and Barcia, M. T. (2022). Microwave hydrodiffusion and gravity model with a unique hydration strategy for exhaustive extraction of anthocyanins from strawberries and raspberries. *Food Chemistry*, 383, 132446.
- Gaba, R., Kaur, N., Urvika, Pal, A., and Sharma, D. (2023). Study of intermolecular interactions present in ternary mixtures containing sugar alcohol and choline chloride at different temperatures. *Journal of Molecular Liquids*, 380, 121766. <https://doi.org/https://doi.org/10.1016/j.molliq.2023.121766>

- Gaioto, R. C., Dias, M. C. G. C., Ndiaye, P. M., Igarashi-Mafra, L., and Mafra, M. R. (2023). Choline chloride aqueous solution: A thermophysical study. *Fluid Phase Equilibria*, 574, 113903. <https://doi.org/https://doi.org/10.1016/j.fluid.2023.113903>
- García, G., Aparicio, S., Ullah, R., and Atilhan, M. (2015). Deep Eutectic Solvents: Physicochemical Properties and Gas Separation Applications. *Energy and Fuels*, 29(4), 2616–2644. <https://doi.org/10.1021/ef5028873>
- Giampieri, F., Tulipani, S., Alvarez-Suarez, J. M., Quiles, J. L., Mezzetti, B., and Battino, M. (2012). The strawberry: Composition, nutritional quality, and impact on human health. *Nutrition*, 28(1), 9–19. <https://doi.org/https://doi.org/10.1016/j.nut.2011.08.009>
- Gómez-Urios, C., Viñas-Ospino, A., Puchades-Colera, P., Blesa, J., López-Malo, D., Frígola, A., and Esteve, M. J. (2023). Choline chloride-based natural deep eutectic solvents for the extraction and stability of phenolic compounds, ascorbic acid, and antioxidant capacity from Citrus sinensis peel. *LWT*, 177, 114595. <https://doi.org/https://doi.org/10.1016/j.lwt.2023.114595>
- Hellström, J., Karhu, S., Karhu, J., Järvenpää, E., and Välimaa, A.-L. (2024). Phenolic profiles differentiate wild bilberry and cultivated blueberry fruit. *LWT*, 199, 116080. <https://doi.org/https://doi.org/10.1016/j.lwt.2024.116080>
- Hikmawanti, N. P. E., Ramadon, D., Jantan, I., and Mun'im, A. (2021). Natural deep eutectic solvents (Nades): Phytochemical extraction performance enhancer for pharmaceutical and nutraceutical product development. In *Plants* (Vol. 10, Issue 10). MDPI. <https://doi.org/10.3390/plants10102091>
- Hong, S. M., Kamaruddin, A. H., and Nadzir, M. M. (2023). Sustainable ultrasound-assisted extraction of polyphenols from Cinnamomum cassia bark using hydrophilic natural deep eutectic solvents. *Process Biochemistry*, 132, 323–336. <https://doi.org/https://doi.org/10.1016/j.procbio.2023.07.026>
- Hosseini, S. A., and Alvarez-Galvan, M. C. (2016). Study of physical–chemical properties and catalytic activities of ZnCr<sub>2</sub>O<sub>4</sub> spinel nano oxides obtained from different methods—Modeling the synthesis process by response surface methodology and optimization by genetic algorithm. *Journal of the Taiwan Institute of Chemical Engineers*, 61, 261–269. <https://doi.org/https://doi.org/10.1016/j.jtice.2015.12.024>
- Hussain, S., Sharma, M., Jarg, T., Aav, R., and Bhat, R. (2023). Natural pigments (anthocyanins and chlorophyll) and antioxidants profiling of European red and green gooseberry (*Ribes uva-crispa* L.) extracted using green techniques (UAE-citric acid-mediated extraction). *Current Research in Food Science*, 7, 100629. <https://doi.org/https://doi.org/10.1016/j.crfs.2023.100629>
- Hussain, S. Z., Naseer, B., Qadri, T., Fatima, T., and Bhat, T. A. (2021). Strawberry (*F. × ananassa*)—Morphology, Taxonomy, Composition and Health Benefits. In S. Z. Hussain, B. Naseer, T. Qadri, T. Fatima, and T. A. Bhat (Eds.), *Fruits Grown in Highland Regions of the Himalayas: Nutritional and Health Benefits* (pp. 219–228). Springer International Publishing. [https://doi.org/10.1007/978-3-030-75502-7\\_17](https://doi.org/10.1007/978-3-030-75502-7_17)
- José, A. G. M., Carrera, C., Barbero, G. F., and Palma, M. (2022). A comparison study between ultrasound–assisted and enzyme–assisted extraction of

- anthocyanins from blackcurrant (*Ribes nigrum* L.). *Food Chemistry: X*, 13, 100192. <https://doi.org/https://doi.org/10.1016/j.fochx.2021.100192>
- Jovanović, M. S., Krgović, N., Živković, J., Stević, T., Zdunić, G., Bigović, D., and Šavikin, K. (2022). Ultrasound-Assisted Natural Deep Eutectic Solvents Extraction of Bilberry Anthocyanins: Optimization, Bioactivities, and Storage Stability. *Plants*, 11(20). <https://doi.org/10.3390/plants11202680>
- Kiene, M., Zaremba, M., Januschewski, E., Juadjur, A., Jerz, G., and Winterhalter, P. (2024). Sustainable In Silico-Supported Ultrasonic-Assisted Extraction of Oligomeric Stilbenoids from Grapevine Roots Using Natural Deep Eutectic Solvents (NADES) and Stability Study of Potential Ready-to-Use Extracts. *Foods*, 13(2), 324. <https://doi.org/10.3390/foods13020324>
- Kim, H., Jung, Y. S., Song, N.-E., Yoo, M., Seo, D.-H., Kim, H.-S., and Nam, T. G. (2024). Ultrasound-assisted extraction of major anthocyanins in Korean black raspberries (*Rubus coreanus* Miquel) using natural deep eutectic solvents. *LWT*, 199, 116121. <https://doi.org/https://doi.org/10.1016/j.lwt.2024.116121>
- Koraqi, H., Trajkovska Petkoska, A., Khalid, W., Kumar, N., and Pareek, S. (2023). Optimization of experimental conditions for bioactive compounds recovery from raspberry fruits (*Rubus idaeus* L.) by using combinations of ultrasound-assisted extraction and deep eutectic solvents. *Applied Food Research*, 3(2), 100346. <https://doi.org/https://doi.org/10.1016/j.afres.2023.100346>
- Kousar, M. U., Jabeen, A., Fatima, T., Hussain, S. Z., Zargar, I. A., Amin, T., and Yaseen, M. (2023). Optimization of ultrasonic-assisted extraction of eugenol-rich fraction from basil leaves: Characterization of extract for phenols, flavonoids and antioxidant activity. *Food Chemistry Advances*, 3. <https://doi.org/10.1016/j.focha.2023.100374>
- Koutsoukos, S., Tsiaka, T., Tzani, A., Zoumpoulakis, P., and Detsi, A. (2019). Choline chloride and tartaric acid, a Natural Deep Eutectic Solvent for the efficient extraction of phenolic and carotenoid compounds. *Journal of Cleaner Production*, 241, 118384. <https://doi.org/https://doi.org/10.1016/j.jclepro.2019.118384>
- Lestario, L. N. (2018). *Antosianin: sifat kimia, perannya dalam kesehatan, dan prospeknya sebagai pewarna makanan*. Ugm Press.
- Li, D., Zhang, X., Li, L., Aghdam, M. S., Wei, X., Liu, J., Xu, Y., and Luo, Z. (2019). Elevated CO<sub>2</sub> delayed the chlorophyll degradation and anthocyanin accumulation in postharvest strawberry fruit. *Food Chemistry*, 285, 163–170.
- Lin, S., Meng, X., Tan, C., Tong, Y., Wan, M., Wang, M., Zhao, Y., Deng, H., Kong, Y., and Ma, Y. (2022). Composition and antioxidant activity of anthocyanins from *Aronia melanocarpa* extracted using an ultrasonic-microwave-assisted natural deep eutectic solvent extraction method. *Ultrasonics Sonochemistry*, 89, 106102. <https://doi.org/https://doi.org/10.1016/j.ultsonch.2022.106102>
- Liu, L., Xiao, A., Zhang, Y., and Duan, S. (2023). Efficient Extraction of Flavonoids from Lotus Leaves by Ultrasonic-Assisted Deep Eutectic Solvent Extraction and Its Evaluation on Antioxidant Activities. *Separations*, 10(2). <https://doi.org/10.3390/separations10020065>

- Mustafa, A. M., Angeloni, S., Abouelenein, D., Acquaticci, L., Xiao, J., Sagratini, G., Maggi, F., Vittori, S., and Caprioli, G. (2022). A new HPLC-MS/MS method for the simultaneous determination of 36 polyphenols in blueberry, strawberry and their commercial products and determination of antioxidant activity. *Food Chemistry*, 367, 130743. <https://doi.org/https://doi.org/10.1016/j.foodchem.2021.130743>
- National Center for Biotechnology Information. (2024a). *PubChem Compound Summary for CID 311, Citric Acid*. <https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/Citric-Acid>.
- National Center for Biotechnology Information. (2024b). *PubChem Compound Summary for CID 6209, Choline Chloride*. <https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/Choline-Chloride>.
- National Center for Biotechnology Information. (2024c). *PubChem Compound Summary for CID 61373, Oxalic acid dihydrate*. <https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/Oxalic-Acid-Dihydrate>.
- National Center for Biotechnology Information. (2024d). *PubChem Compound Summary for CID 107689, Lactic Acid, L-*. [https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/Lactic-Acid\\_-L](https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/Lactic-Acid_-L).
- National Center for Biotechnology Information. (2024e). *PubChem Compound Summary for CID 222656, (2S)-2-Hydroxybutanedioic acid*. [https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/2S\\_-2-Hydroxybutanedioic-Acid](https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/2S_-2-Hydroxybutanedioic-Acid).
- National Center for Biotechnology Information. (2024f). *PubChem Compound Summary for CID 444305, L-Tartaric acid*. <https://Pubchem.Ncbi.Nlm.Nih.Gov/Compound/L-Tartaric-Acid>.
- Nunes, A. N., Borges, A., Matias, A. A., Bronze, M. R., and Oliveira, J. (2022). Alternative Extraction and Downstream Purification Processes for Anthocyanins. *Molecules*, 27(2). <https://doi.org/10.3390/molecules27020368>
- Osamede Airouyuwa, J., Mostafa, H., Riaz, A., and Maqsood, S. (2022). Utilization of natural deep eutectic solvents and ultrasound-assisted extraction as green extraction technique for the recovery of bioactive compounds from date palm (*Phoenix dactylifera* L.) seeds: An investigation into optimization of process parameters. *Ultrasonics Sonochemistry*, 91, 106233. <https://doi.org/https://doi.org/10.1016/j.ultsonch.2022.106233>
- Pena-Pereira, F., and De La Calle, I. (2019). Solvents and eutectic solvents. In *Encyclopedia of Analytical Science* (pp. 184–190). Elsevier. <https://doi.org/10.1016/B978-0-12-409547-2.14020-X>
- Petersen, C., Wankhade, U. D., Bharat, D., Wong, K., Mueller, J. E., Chintapalli, S. V., Piccolo, B. D., Jalili, T., Jia, Z., Symons, J. D., Shankar, K., and Anandh Babu, P. V. (2019). Dietary supplementation with strawberry induces marked changes in the composition and functional potential of the gut microbiome in diabetic mice. *The Journal of Nutritional Biochemistry*, 66, 63–69. <https://doi.org/https://doi.org/10.1016/j.jnutbio.2019.01.004>
- Prakash, S., Goswami, A., Patil, R., Mitra, A., and Kutty, N. N. (2024). An eco-friendly approach to extract anthocyanins from rose flowers using natural deep

- eutectic solvents. *Industrial Crops and Products*, 210, 118059. <https://doi.org/https://doi.org/10.1016/j.indcrop.2024.118059>
- Pusty, K., Kumar Dash, K., Giri, S., Raj, G. V. S. B., Tiwari, A., Shaikh, A. M., and Béla, K. (2024). Ultrasound assisted phytochemical extraction of red cabbage by using deep eutectic solvent: Modelling using ANFIS and optimization by genetic algorithms. *Ultrasonics Sonochemistry*, 102, 106762. <https://doi.org/https://doi.org/10.1016/j.ultsonch.2024.106762>
- Radošević, K., Ćurko, N., Gaurina Srček, V., Cvjetko Bubalo, M., Tomašević, M., Kovačević Ganić, K., and Radojčić Redovniković, I. (2016). Natural deep eutectic solvents as beneficial extractants for enhancement of plant extracts bioactivity. *LWT*, 73, 45–51. <https://doi.org/https://doi.org/10.1016/j.lwt.2016.05.037>
- Raj, T., and Singh, V. (2024). Natural deep eutectic solvents (NADES) assisted deconstruction of oilcane bagasse for high lipid and sugar recovery. *Industrial Crops and Products*, 210, 118127. <https://doi.org/https://doi.org/10.1016/j.indcrop.2024.118127>
- Ratanasongtham, P., Bunmusik, W., Luangkamin, S., Mahatheeranont, S., and Suttiarporn, P. (2024). Optimizing green approach to enhanced antioxidants from Thai pigmented rice bran using deep eutectic solvent-based ultrasonic-assisted extraction. *Heliyon*, 10(1), e23525. <https://doi.org/https://doi.org/10.1016/j.heliyon.2023.e23525>
- Sainakhm, M., Jantrawut, P., Kiattisin, K., Chittasupho, C., and Singh, S. (2023). Potential of green extraction using edible deep eutectic solvents on the bioactivities from *Curcuma aromatica* rhizome extracts for food application. *Journal of Agriculture and Food Research*, 14, 100868. <https://doi.org/https://doi.org/10.1016/j.jafr.2023.100868>
- Sanjaya, Y. A., Tola, P. S., and Rahmawati, R. (2022). Ultrasound-assisted Extraction as a Potential Method to Enhanced Extraction of Bioactive Compound. *Nusantara Science and Technology Proceedings*, 191–198. <https://doi.org/10.11594/nstp.2022.2729>
- Santos-Martín, M., Cubero-Cardoso, J., González-Domínguez, R., Cortés-Triviño, E., Sayago, A., Urbano, J., and Fernández-Recamales, Á. (2023a). Ultrasound-assisted extraction of phenolic compounds from blueberry leaves using natural deep eutectic solvents (NADES) for the valorization of agrifood wastes. *Biomass and Bioenergy*, 175, 106882. <https://doi.org/https://doi.org/10.1016/j.biombioe.2023.106882>
- Santos-Martín, M., Cubero-Cardoso, J., González-Domínguez, R., Cortés-Triviño, E., Sayago, A., Urbano, J., and Fernández-Recamales, Á. (2023b). Ultrasound-assisted extraction of phenolic compounds from blueberry leaves using natural deep eutectic solvents (NADES) for the valorization of agrifood wastes. *Biomass and Bioenergy*, 175, 106882. <https://doi.org/https://doi.org/10.1016/j.biombioe.2023.106882>
- Sejati, T. M. A. (2017). *Budi Daya Stroberi*. CV. Pustaka Bengawan.
- Setiani, A. (2022). *Menanam Stroberi*. Penerbit Mutiara Aksara.

- Silva, D. T. da, Pauletto, R., Cavalheiro, S. da S., Bochi, V. C., Rodrigues, E., Weber, J., Silva, C. de B. da, Morisso, F. D. P., Barcia, M. T., and Emanuelli, T. (2020a). Natural deep eutectic solvents as a biocompatible tool for the extraction of blueberry anthocyanins. *Journal of Food Composition and Analysis*, *89*, 103470. <https://doi.org/https://doi.org/10.1016/j.jfca.2020.103470>
- Silva, D. T. da, Pauletto, R., Cavalheiro, S. da S., Bochi, V. C., Rodrigues, E., Weber, J., Silva, C. de B. da, Morisso, F. D. P., Barcia, M. T., and Emanuelli, T. (2020b). Natural deep eutectic solvents as a biocompatible tool for the extraction of blueberry anthocyanins. *Journal of Food Composition and Analysis*, *89*, 103470. <https://doi.org/https://doi.org/10.1016/j.jfca.2020.103470>
- Sodhi, G. S., and Kaur, J. (2016). Physical developer method for detection of latent fingerprints: A review. *Egyptian Journal of Forensic Sciences*, *6*(2), 44–47. <https://doi.org/https://doi.org/10.1016/j.ejfs.2015.05.001>
- Sunarya, Y. (2012). *Kimia Dasar 2 Berdasarkan Prinsip-Prinsip Kimia Terkini*. Bandung: CV. Yarma Widya.
- Syukri, D. (2021). *Bagian Alir Analisis Proksimat Bahan Pangan*. Andalas University Press.
- Taghavi, T., Patel, H., and Rafie, R. (2023). Extraction Solvents Affect Anthocyanin Yield, Color, and Profile of Strawberries. *Plants*, *12*(9). <https://doi.org/10.3390/plants12091833>
- Takla, S. S., Shawky, E., Hammada, H. M., and Darwish, F. A. (2018). Green techniques in comparison to conventional ones in the extraction of Amaryllidaceae alkaloids: Best solvents selection and parameters optimization. *Journal of Chromatography A*, *1567*, 99–110. <https://doi.org/https://doi.org/10.1016/j.chroma.2018.07.009>
- Velásquez, P., Bustos, D., Montenegro, G., and Giordano, A. (2021). Ultrasound-Assisted Extraction of Anthocyanins Using Natural Deep Eutectic Solvents and Their Incorporation in Edible Films. *Molecules*, *26*(4). <https://doi.org/10.3390/molecules26040984>
- Viñas-Ospino, A., Panić, M., Bagović, M., Radošević, K., Esteve, M. J., and Radojčić Redovniković, I. (2023). Green approach to extract bioactive compounds from orange peel employing hydrophilic and hydrophobic deep eutectic solvents. *Sustainable Chemistry and Pharmacy*, *31*, 100942. <https://doi.org/https://doi.org/10.1016/j.scp.2022.100942>
- Wang, Y., Ye, Y., Wang, L., Yin, W., and Liang, J. (2021). Antioxidant activity and subcritical water extraction of anthocyanin from raspberry process optimization by response surface methodology. *Food Bioscience*, *44*, 101394.
- Widiatmoko, H. (2016). *Sehat Tanpa Obat dengan Si Merah Stroberi*. Andi Offset.
- Xi, Y., Fan, F., and Zhang, X. (2023). Microbial L-malic acid production: History, current progress, and perspectives. *Green Carbon*, *1*(2), 118–132. <https://doi.org/https://doi.org/10.1016/j.greenca.2023.10.005>
- Yang, W., Liu, S., Marsol-Vall, A., Tähti, R., Laaksonen, O., Karhu, S., Yang, B., and Ma, X. (2021). Chemical composition, sensory profile and antioxidant capacity of low-alcohol strawberry beverages fermented with *Saccharomyces*

cerevisiae and *Torulaspota delbrueckii*. *LWT*, 149, 111910.  
<https://doi.org/https://doi.org/10.1016/j.lwt.2021.111910>

Yenrina, R. (2015). *Metode Analisis Bahan Pangan dan Komponen Bioaktif*. Andalas University Press.

Yuan, Y., Tian, Y., Gao, S., Zhang, X., Gao, X., and He, J. (2023). Effects of environmental factors and fermentation on red raspberry anthocyanins stability. *LWT*, 173, 114252. <https://doi.org/https://doi.org/10.1016/j.lwt.2022.114252>

Zannou, O., and Koca, I. (2022). Greener extraction of anthocyanins and antioxidant activity from blackberry (*Rubus spp*) using natural deep eutectic solvents. *LWT*, 158, 113184.  
<https://doi.org/https://doi.org/10.1016/j.lwt.2022.113184>

Zannou, O., Pashazadeh, H., Ibrahim, S. A., Koca, I., and Galanakis, C. M. (2022). Green and highly extraction of phenolic compounds and antioxidant capacity from kinkeliba (*Combretum micranthum* G. Don) by natural deep eutectic solvents (NADESS) using maceration, ultrasound-assisted extraction and homogenate-assisted extraction. *Arabian Journal of Chemistry*, 15(5), 103752. <https://doi.org/https://doi.org/10.1016/j.arabjc.2022.103752>

Zhang, S., Lin, S., Zhang, J., and Liu, W. (2024). Ultrasound-assisted natural deep eutectic solvent extraction of anthocyanin from *Vitis davidii* Foex. pomace: Optimization, identification, antioxidant activity and stability. *Heliyon*, 10(12), e33066. <https://doi.org/https://doi.org/10.1016/j.heliyon.2024.e33066>

