

DAFTAR PUSTAKA

- Abd Ghafar, M. F., K. N. Prasad, K. K. Weng, A. Ismail, 2010, Flavonoid, hesperidine, total phenolic contents and antioxidant activities from Citrus species. *African Journal of Biotechnology*, 9(3), 326–330
- Alam, M. N., N. J. Bristi, M. Rafiquzzaman, 2013, Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*, 21(2), 143–152. <https://doi.org/10.1016/j.jps.2012.05.002>
- Alfonso de Araujo,T., A.M.J. Barbosa, L.H. Viana, V. S. Ferreira, 2011, Electroanalytical determination of TBHQ , a synthetic antioxidant, in soybean biodiesel samples, *Fuel*, 90 : 707–712
- Andarwulan, N., R. Batari, D. A. Sandrasari, B. Bolling, H. Wijaya, 2010, Flavonoid content and antioxidant activity of vegetables from Indonesia,*Food Chemistry*. <https://doi.org/10.1016/j.foodchem.2010.01.033>
- Apak, R., K. Guclu, B. Demirata, M. Ozyurek, S.E.Celik, B. Bektasoglu, 2007, Comparative evaluation of various total antioxidant capacity assays applied to phenolic compounds with the CUPRAC assay. *Molecules*, 12:1496
- Arnao M. B., 2000, Some methodological problems in the determination of antioxidant activity using chromogen radicals: a practical case. *Trends Food Sci. Technol.*, 11:419. doi:10.1016/S0924-2244 (01)00027-9
- Benzie, I.F.F., J.J. Strain, 1999. Ferric reducing antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol*, 299, 15–27
- Berker, K. I., K. Güçlü, I. Tor, R. Iapak, 2007, Comparative evaluation of Fe(III) reducing power-based antioxidant capacity assays in the presence of phenanthroline, batho-phenanthroline, tripyridyltriazine (FRAP), and ferricyanide reagents. *Talanta*, <https://doi.org/10.1016/j.talanta.2007.01.019>
- Biparva, P., M. Ehsani, M. R. Hadjmhommadi, 2012, Dispersive liquid–liquid microextraction using extraction solvents lighter than water combined with high performance liquid chromatography for determination of synthetic antioxidants in fruit juice samples, *Journal of Food Composition and Analysis*, 27 : 87–94
- Blois, M.S., 1958,Antioxidant determinations by the use of a stable free radical. *Nature*, 26: 1199-1200

- Bolanos de la Torre, A. A. S., T. Henderson, P. S. Nigam, R. K. Owusu-Apenten, 2015, A universally calibrated microplate Ferric Reducing Antioxidant Power (FRAP) assay for foods and applications to Manuka honey, *Food Chemistry*, 174 : 119-123
- Bolton, J.L., M.A.Trush, T.M. Penning, G. Dryhurst, T.J.Monks, 2000, Role of quinones in toxicology. *Chem. Res. Toxicol.* 13, 135– 160.
- Brand-Williams, W., M. Cuvelier, C. Berset, 1995, Use of free radical method to evaluate antioxidant activity, *LWT-Food Science and Technology*, 28 (1): 25-3
- Bursal, E., E. Köksal, İ. Gülcin, G. Bilsel, A. C. Gören, 2013, Antioxidant activity and polyphenol content of cherry stem (*Cerasus avium L.*) determined by LC-MS/MS, *Food Research International*, 51 : 66–74
- Butsat, S., S.Siriamornpun, 2016, Effect of solvent types and extraction times on phenolic and flavonoid contents and antioxidant activity in leaf extracts of *Amomum chinense* C., *International Food Research Journal*, 23(1): 180-187
- Calado, J. C. P., P. A. Albertão, E. A . de Oliveira, M. H. Sisto, A. C. H. Frankland, M. C.Marcucci, 2015, Flavonoid Contents and Antioxidant Activity in Fruit , Vegetables and Other Types of Food. *Agricultural Sciences*, 6(April), 426–435. <https://doi.org/10.4236/as.2015.64042>
- Cao, G., E. Sofic, R. L. Prior, 1996, Antioxidant capacity of tea and common vegetables. *Journal Agricultural and Food Chemistry*, 44, 3426–3431
- Chang, S. Q., A. Azrina, 2017, Antioxidant content and activity in different parts of pomelo (*Citrus grandis* (L.) Osbeck) by-products., *Acta Horticulturae*, (1152), 27–34.
- Chew, K.K., S.Y. Ng, Y.Y. Thoo, M.Z. Khoo, W.M .Wan Aida, C.W.Ho, 2011, Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of *Centella asiatica* extracts. *International Food Research Journal*, 18: 571-578.
- Cortés-Rojas, D.F., C.R.F. Souza, W.P. Oliveira, 2015, Optimization of spray dry-ing conditions for production of *Bidens pilosa* L. dried extract, *Chem. Eng. Res.*, <http://dx.doi.org/10.1016/j.cherd.2014.06.010>.
- Dasgupta, A., K. Klein, 2014, Antioxidants in Food, Vitamins and Supplements: Prevention and Treatment of Disease, Elsivier, London.
- De Beer, D., E. Joubert, WCA. Gelderblom, M. Manley, 2003, Antioxidant activity of South African red and white cultivar wines: free radical scavenging. *J Agric Food Chem*, 51:902. doi:10.1021/jf026011o

Dembinska-Kiec, A., O. Mykkanen, B. Kiec-Wilk, H. Mykkanene, 2008, Antioxidant Phyto-chemicals against Type 2 Diabetes. *British J. Nutri.*, 99: 109-117.

Deng, G. F., X. R. Xu, Y.J. Guo, E. Q. Xiaa, S. Lia, S. Wu, F. Chen, W. H. Linga, H. B. Li, 2013, Determination of antioxidant property and their lipophilic and hydrophilic phenolic contents in cereal grains, *Journal of functional foods*, 4 : 906 –914

Flis, S., Z. Jastrzebski, J. Namiesnik, P. Arancibia-Avila, F. Toledo, H. Leontowicz, M. Suhaj, S. Trakhtenber, S. Gorinsteing, 2012, Evaluation of inhibition of cancer cell proliferation in vitro with different berries and correlation with their antioxidant levels by advanced analytical methods, *Journal of Pharmaceutical and Biomedical Analysis*, 62 : 68– 78

Frei, B., 1991, Ascorbic acid protects lipids in human plasma and lowdensity lipoprotein against oxidative damage. *Am. J. Clin. Nutr.*, 54 (6 SS), S1113-S1118.

Garcia, A., M. G. Alriols, G. Spigno, J. Labidi, 2012, Lignin as natural radical scavenger, , Effect of the obtaining and purification processes on the antioxidant behaviour of lignin, *Biochemical Engineering Journal*, 67 : 173– 185

Ghasemi, K., Y. Ghasemi,M. A. Ebrahimzadeh, 2009. Antioxidant activity, phenol and flavonoid contents of 13 citrus species peels and tissues. *Pakistan Journal of Pharmaceutical Sciences*, 22(3): 277-281.

Gheldorf, N., N.J. Engeseth, 2002,Antioxidant capacity of honeys from various floral sources based on the determination of oxygen radical absorbance capacity and inhibition of in vitro lipoprotein oxidation in human serum samples. *J Agric Food Chem*, 50:3050. doi:10.1021/jf0114637

Gomes, S. M., M. E.Ghica, I. A.Rodrigues, E.de Souza Gil, A. M. Oliveira-Brett, 2016, Flavonoids electrochemical detection in fruit extracts and total antioxidant capacity evaluation. *Talanta*, 154, 284–291.

Hardiana, R., R. Zaharah, T . Anita, 2012, Antivitas antioksidan senyawa golongan fenol dari beberapa jenis tumbuhan famili Malveceae, *JKK*, 1(1), 8-13, ISSN 2303 - 1077

Haris, D.C.,2003, Determination of Iron with 1,10-Phenanthroline, in: Quantitative Chemical Analysis, 6th ed., W. H.Freeman & Company, New York, pp. 258-261, 407-422, 453, 461-476, 707-709.

Han, X., T. Shen, H. Lou, 2007, Dietary Polyphenols and Their Biological Significance. *International Journal Molecular Science*, 8, 950988.<http://dx.doi.org/10.3390/i8090950>

Ho ,HY.,ML. Cheng, DT. Chiu, 2007, Glucose 6-phosphate dehydrogenase: from oxidative stress to cellular functions and degenerative diseases,*Redox Rep J*;12:109_18.

Houston, M. C., 2010, The role cellular micronutrient analysis, nutraceuticals, vitamins and antioxidants and minerals in the prevention and treatment of hypertension and cardiovascular disease,*Therapeutic Advances in Cardiovascular Disease*, 4(3), 165–183

Huang, H., Y. Sun, S. Lou, H. Li, X. Ye, 2014, In vitro digestion combined with cellular assay to determine the antioxidant activity in Chinese bayberry (*Myrica rubra Sieb. et Zucc.*) fruits: A comparison with traditional methods, *Food Chemistry*, 146 : 363–370

Huang,D., B. Ou, R.L. Prior,2005, The chemistry behind antioxidant capacity assays, *J Agric Food Chem*,53:1841. doi:10.1021/jf030723c

Ismail, A.,T. Hong, 2002,Antioxidant activity of selected commercial seaweeds. *Malaysian Journal of Nutrition*, 8 (2): 167-177.

Ismail, A., Z. M. Marjan, C. W. Foong, 2004, Total antioxidant activity and phenolic content in selected vegetables. *Food Chemistry*,<https://doi.org/10.1016/j.foodchem.2004.01.010>

Jargar, J. G. , S. H. Hattiwale , S. Das, S. A. Dhundasi, K. K. Das, 2012, A modified simple method for determination of serum α -tocopherol (vitamin E), *J Basic Clin Physiol.Pharmacol.*, 23(1) : 45–48

Juániz, I., I. A. Ludwig, E. Huarte, G. Pereira-Caro, J. M. Moreno-Rojas, C.Cid, M.P. De Peña, 2016, Influence of heat treatment on antioxidant capacity and (poly)phenolic compounds of selected vegetables. *Food Chemistry*. <https://doi.org/10.1016/j.foodchem.2015.10.139>

Karadag, A., B. Ozcelik, S. Saner, 2009, Review of Methods to Determine Antioxidant Capacities, *Food Anal. Methods*, 2:41–60, DOI 10.1007/s12161-008-9067-7

Katsube ,N., K , Iwashita, T. Tsushida, K . Yamaki, M . Kobori , 2003, Induction of apoptosis in cancer cells by bilberry (*Vaccinium myrtillus*) and the anthocyanins. *J Agric Food Chem*, 51:68.doi:10.1021/jf025781x

Kinnula, V.L., J.D . Crapo, 2004, Superoxide dismutases in malignant cells and human tumors. *Free Radic. Biol. Med.*, 36, 718–744.

Kraujalyte, V., P. R. Venskutonis, A. Pukalskas, L. C. esoniene, R. Daubaras, 2015, Antioxidant properties, phenolic composition and potentiometric sensor array

evaluation of commercial and new blueberry (*Vaccinium corymbosum*) and bog blueberry (*Vaccinium uliginosum*) genotypes, *Food Chemistry*, 188 : 583–590

Kumazawa, S., T. Nakayama, 2001, Polyphenols in propolis and their antioxidant activity. *Honeybee Science*, 22 (1): 1-8.

Kunwar ,A., KI. Priyadarsini, 2011, Free radicals, oxidative stress and importance of antioxidants in human health. *J Med Allied Sci*;1:53-60.

Kwon, S. W., S. S. Hong, J. I. Kim, I. Ahn, 2003,Antioxidant properties of heat treated Hibiscus syriacus. *Biology Bulletin*, 30 (1), 15-16.

Langrish, J.P., X. Li, S. Wang, M.M . Lee, al. Et, 2012, Reducing personal exposure to particulate air pollution improves cardiovascular health in patients with coronary heart disease. *Environ Health Perspect*;120:367_72.

Li ,X.Y., PS , Gilmour, K. Donaldson, W . MacNee, 1996, Free radical activity and pro-inflammatory effects of particulate air pollution (PM10) in vivo and in vitro. *Thorax*,51:1216_22.

Lim, B. P., A. Nagao, J. Terao, K. Tanaka, T . Suzuki,K. Takama, 1992,Antioxidant activity of xanthophylls on peroxy radical mediated phospholipid peroxidation. *Biochim. Biophys. Acta*, 1126, 178-184.

Lim, S. M.,S. P. Loh,2016, In vitro antioxidant capacities and antidiabetic properties of phenolic extracts from selected citrus peels. *International Food Research Journal*, 23(1), 211–219.

Lin, Z., H. Wang, Y. Xu, J. Dong, Y. Hashi, S. Chen, 2012, Identification of antioxidants in *Fructus aurantii* and its quality evaluationusing a new on-line combination of analytical techniques, *Food Chemistry*, 134 : 1181–1191

Litescu ,S.C., A.V. Sandra, S.A.V. Eremia, M. Diaconu, A . Tache, al. Et, 2011, Biosensors Applications on Assessment of Reactive Oxygen Species and Antioxidants. Environmental Biosensors. *In Tech Rijeka Croatia*.

Loizzo, M. R., A. Pugliese, M. Bonesi, F. Menichini, R. Tundis, 2015, Evaluation of chemical profile and antioxidant activity of twenty cultivars from *Capsicum annuum*, *Capsicum baccatum*, *Capsicum chacoense* and *Capsicum chinense*: A comparison between fresh and processed peppers, *LWT - Food Science and Technology*, 64 : 623-631

Lopez, M. D. M. C., J. M. L. Villarino, M. V. G. Rodriguez, L. F. B. Losada, 2011, Development, Validation and Application of Micellar Electrokinetic Capillary Chromatography Method for Routine Analysis of Catechins, Quercetin and Thymol in Natural Samples, *Microchemical Journal*, 99, 461–469

MacDonald-Wicks , L.K., L.G . Wood, M.L. Garg , 2006, Methodology for the determination of biological antioxidant capacity in vitro: a review. *J Sci Food Agric*, 86(13):2046. doi:10.1002/jsfa.2603

Madhujith ,T., M. Izydorczyk, F. Shahidi, 2006,Antioxidant properties of pearled barley fractions. *J Agric Food Chem*, 54:3283

Magalhaes , L.M., M.A . Segundo, S. Reis, J.L.F.C . Lima , 2008, Methodological aspects about in vitro evaluation of antioxidant properties. *Anal Chim Acta*, 613:1. doi:10.1016/j.aca.2008.02.047

Maranz ,S.,Z .Wiesman, N. Garti, 2003, Phenolic constituents of shea (*Vitellaria paradoxa*) kernels. *J Agric Food Chem*, 51:6268. doi:10.1021/jf034687t

Martínez, R., P. Torres , M.A. Meneses, J.G. Figueroa, J.A. Pérez-Álvarez , M. Viuda-Martos, 2012, Chemical, technological and in vitro antioxidant properties of cocoa (*Theobroma cacao* L.) co-products, *Food Research International*, 49 : 39-45

Mehta, P. S., V. B. Patel, 2012, Spectrophotometric method for determination of Fe(II) and Zn(II) in multivitamin soft gel capsule, *International Journal of Pharmaceutical Research & Analysis*, 2(2), 87-91

Meyer, A. S., M. Heinonen, E. N. Frankel, 1998, Antioxidant interactions of catechin, cyanidin, cafeic acid, quercetin, and ellagic acid on human LDL oxidation. *Food Chemistry*, 61, 71–75.

Miller, H., F. Rigelhof, L. Marquart, A. Prakash, M. Kanter, 2000,Antioxidant Content of Whole Grain Breakfast Cereals, Fruits and Vegetables. *Journal of the American College of Nutrition*, 19 (3): 312S-319S.

Molyneux, P., 2004, The Use of The Stable Free Radical Diphenylpicryl-hydrazone (DPPH) for Estimating Antioxidant Activity, *Songklanakarin J. Sci. Technol.*, 26(2), 211-21

Moreira, F. T. C., J. R. L. Guerreiro, R. Barros, M. G. F. Sales, 2012, The Effect of Method, Standard and Sample Components on The Total Antioxidant Capacity of Commercial Waters Assessed by Optical Conventional Assays, *Food Chemistry*, 134 : 564–571

Motilva, M-J., A. Macia, M-P. Romero, A. Labrador, A. Dominguez, L. Peiro, 2014, Optimisation and validation of analytical methods for the simultaneous extraction of antioxidants: Application to the analysis of tomato sauces, *Food Chemistry*, 163 : 234–243

Naczk, M., F. Shahidi, 2004, Extraction and analysis of phenolics in food. *Journal of Chromatography A.*, 1054(1-2): 95-111.

Nakamura, Y., S. Tsuji, Y. Tonogai, 2003, Method for analysis of tannic acid and its metabolites in biological samples: application to tannic acid metabolism in the rat. *J Agric Food Chem*, 51:331. doi:10.1021/jf020847+

Novita, M., I. Sulaiman, S. Ura, 2016, Pengaruh Jenis Pelarut terhadap Aktivitas Antioksidan dan Kandungan Fenol Beberapa Jenis Bayam dan Sayuran Lain, *JIM Pertanian Unsyiah – THP*, Vol. 1, No. 1, November 2016: 935-940

Ohlsson, T., N. Bengtsson, 2002, Minimal Processing technologies in food industry. Retrieved from (<http://books.google.com>).

Ojezele, O.J., M.O. Ojezele, A.M. Adeosun, 2016, Comparative Phytochemistry and Antioxidant Activities of Water and Ethanol Extract of *Annona muricata Linn* Leaf, Seed and Fruit, *Advan. Biol. Res.*, 10 (4): 230-235,

Ozcelik ,B., J.H. Lee, D.B. Min, 2003, Effects of light, oxygen, and pH on the absorbance of 2,2-diphenyl-1-picrylhydrazyl (DPPH). *J Food Sci* 68(2):487. doi:10.1111/j.1365-2621.2003.tb05699.x

Pichaiyongvongdee, S., B. Rattanapun, R. Haruenkit, 2014, Total polyphenol content and antioxidant properties in different tissues of seven pomelo (*Citrus grandis* (L .) Osbeck, *Cultivars*, 996, 989–996.

Pisoschi, A.M., G.P. Negulescu, 2011, 1:1 Methods for Total Antioxidant Activity Determination: A Review, *Biochem & Anal Biochem*, 1 (1), 1 -10.

Porgal, E., E. Buyuktuncel, 2012, Determination of Phenolic Composition and Antioxidant Capacity of Native Red Wines by High Performance Liquid Chromatography and Spectrophotometric Methods, *Food Research International*, 45, 145–154

Powere, S. K., M.J. Jackson, 2008, Exercise induced oxidative stress: cellular mechanism and impact on muscle force production. *Physiol Rev*, 88:1243_76.

Prakash, A., F. Rigelhof, E. Miller, 2001, *Antioxidant Activity, Medalliaon Laboratories Analytical Progress*, vol 10, No.2

Prior, RL., X. Wu, K. Schaich, 2005, Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J Agric Food Chem*, 53(8):3101–3113. doi:10.1021/jf0478861

Rahman, N. F. A., R. Shamsudin, A. Ismail, N. N. A. K. Shah, J. Varith, 2018, Effects of drying methods on total phenolic contents and antioxidant capacity of the

pomelo (*Citrus grandis* (L.) Osbeck) peels. *Innovative Food Science and Emerging Technologies*. <https://doi.org/10.1016/j.ifset.2018.01.009>

Roberts, W. G., M. H. Gordon, 2003, Determination of the total antioxidant activity of fruits and vegetables by a liposome assay. *Journal of Agricultural and Food Chemistry*. <https://doi.org/10.1021/jf025983t>

Rufino, M., M. do S. R. E . Alves, E. S. de Brito, J. Pérez-Jiménez, F. Saura-Calixto,J. Mancini-Filho, 2010, Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. *Food Chemistry*, 121(4), 996–1002. <https://doi.org/10.1016/j.foodchem.2010.01.037>

Sanchez-Moreno, C., J. A. Larrauri, F. Saura-Calixto, 1999, Free radical scavenging capacity of selected red, rosé and white wines. *Journal of the Science of Food and Agriculture*, 79 (10): 1301-1304

Sanchez-Moreno, C., 2002, Review: Methods used to evaluate the free radical scavenging activity in foods and biological systems, *Food Sci.Technol.Int.*, 8, 121 - 137

Sas, K., Robotka, H., J. Toldi, L. Vecsei, 2007, Mitochondrial, metabolic disturbances, oxidative stress and kynurenone system, with focus on neurodegenerative disorders. *J. Neurol. Sci.*, 257, 221–239.

Sayuti, K., R. Yenrina, 2015, Antioksidan, Alami dan Sintetik, Andalas University Press, Padang

Schlesier, K., M. Harwat, V.Bohm, R. Bitsch, 2002, Assessment of antioxidant activity by using different in vitro methods. *Free Radical Research*, 36(2), 177–187. <https://doi.org/10.1080/10715760290006411>

Segev, A., H. Badani, K.Kapulnik, I.Shomer, M. Oren-Shamir, S. Galili, 2010, Determination of Polyphenols, Flavonoids and Antioxidant Capacity in Colored Chick-Pea (*Cicer arietinum* L.). *Journal Food Science*, 75, S115- S119. <http://dx.doi.org/10.1111/j.1750-3841.2009.01477.x>

Shahidi, F., C.M . Liyana-Pathirana, D. S.Wall, 2006, Antioxidant activity of white and black sesame seeds and their hull fractions. *Food Chem.*, 99(3):478. doi:10.1016/j.foodchem.2005.08.009

Singh, U., I. Jialal, 2006, Oxidative stress and atherosclerosis. *Pathophysiology* 13, 129–142.

Smith, H., S. Doyle, R. Murphy, 2015, Filamentous fungi as a source of natural antioxidants, *Food Chemistry*, 185:389–397

- Smith, M.A., C.A. Rottkamp, A. Nunomura, A.K . Raina, G. Perry, 2000, Oxidative stress in Alzheimer's disease. *Biochim. Biophys. Acta*, 1502, 139–144.
- Srivastava, J.K., E. Shankar,S. Gupta, 2010, Chamomile: a tanaman obat medicine of thepast with a bright future (Review). *Mol. Med. Rep.* 3, 895–901.
- Stratil, P., B. Klejdus, V. Kuban, 2006, Determination of total content of phenolic compounds and their antioxidant activity in vegetablesevaluation of spectrophotometric methods. *J Agric Food Chem*, 54:607. doi:10.1021/jf052334j
- Stratil, P., B. Klejdus, V. Kubáň, 2007, Determination of phenolic compounds and their antioxidant activity in fruits and cereals. *Talanta*, 71(4), 1741–1751. <https://doi.org/10.1016/j.talanta.2006.08.012>
- Sudhakar, S., R. A. Nazeer, 2015, Preparation of potent antioxidant peptide from edible part of shortclub cuttlefish against radical mediated lipid and DNA damage, *LWT - Food Science and Technology*, 64 : 593-601
- Sultana, B., F.Anwar, M. Ashraf, 2009, Effect of Extraction Solvent/Technique on the Antioxidant Activity of Selected Medicinal Plant Extracts, *Molecules*, 14, 2167-2180; doi:10.3390/molecules14062167
- Szydłowska-Czerniak, A., A. Łaszewska, 2015, Effect of refining process on antioxidant capacity, total phenolics and prooxidants contents in rapeseed oils *LWT - Food Science and Technology*, 64 :853-859
- Szydłowska-Czerniak, A., A. Tułodziecka, 2013, Comparison of a silver nanoparticle-based method and the modified spectrophotometric methods for assessing antioxidant capacity of rapeseed varieties, *Food Chemistry*, 141, 1865–1871
- Szydłowska-Czerniak, A., C. Dianoczki, K. Recseg, G. Karlovits, E. Szłyk, 2008, Determination of antioxidant capacities of vegetable oils by ferric-ion spectrophotometric methods. *Talanta*. <https://doi.org/10.1016/j.talanta.2008.04.055>
- Tepe, B., S. Degerli, S. Arslan, E. Malatyali, C. Sarikurkcu, 2011, Determination of chemical profile, antioxidant, DNA damage protection and antiamoebic activities of Teucrium polium and Stachys iberica, *Fitoterapia*, 82 : 237–246
- Thaipong, K., U. Boonprakob, CK. rosby, CL. isneros-Zevallos, D. Hawkins Byrne, 2006, Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19(6–7), 669–675.
- Tokusoglu, O., Y.Z. Ildrim, 2012, Effects of Cooking Methods on The Anthocyanin Levels and Antioxidant Activity of Local Turkish Sweetpotato (*Ipomoea batatas*

L) Cultivar Hatay Kirmizi : Boiling, Steaming, and Frying Effects, *J. Field Crops*, 17(1), 87 – 90.

Tsuda, T., M. Watanabe, K. Ohshima, A. Yamamoto, S. Kawakishi, T. Osawa, 1994, Antioxidative components isolated from the seed of tamarind (*Tamarindus indica L.*). *Journal of Agricultural and Food Chemistry*, 42, 2671–2674.

Tsumbu, C.N., G.D. Dupont,M. Tits, A.L. ngenot, T. Franck, D . Serteyn,A.M. Mickalad, 2011, Antioksidant dan Antiradical Activities of Manihot esculenta Crantz (Euphorbiaceae) Leaves and Other Selected Tropical Green Vegetables Investigated on Lipoperoxidation and Phorbol-12-myristate-13-acetate (PMA) Activated Monocytes.*Nutrient*, 3: 818-838.

Velioglu, Y.S., G. Mazza, L. Gao, B.D. Oomah, 1998, Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *J Agric Food Chem*, 46(10):4113. doi:10.1021/jf9801973

Vinson, J.A., X.H. Su, L. Zubik, P. Bose, 2001, Phenol antioxidant quantity and quality in foods: fruits. *J Agric Food Chem*, 49:5315.doi:10.1021/jf0009293

Viuda-Martos, M. Y., J. Ruiz-Navajas, E. Fernández-López, E. Sendra, J. A.Sayas-Barberá, Pérez-Álvarez, 2011, Antioxidant properties of pomegranate (*Punica granatum L.*) bagasses obtained as co-product in the juice extraction, *Food Research International*, 44 : 1217–1223

Vyas, N., N. Tailang, GN.P. avatia, B.K. Gupta, 2010, Antioxidant Potential of *Psidiumguajava* Linn, *International Journal of pharmtech Research*, Vol.2, No.1, pp 417-419

Wang, M., J.E. Simon, I.F. Aviles, K. He, Q. Zheng, Y. Tadmor, 2003, Analysis of antioxidative phenolic compounds in artichoke (*Cynara scolymus L.*). *J Agric Food Chem*, 51(3):601. doi:10.1021/jf020792b

Wright, J. S. E. R. Johnson, G. A. DiLabio,2001, Predicting the activity of phenolic antioxidants: Theoretical method, analysis of substituent effects, and application to major families of antioxidants. *J. Am. Chem. Soc.*, 123, 1173-1183.

Xu ,Y.,Y., Gu, S.Y. Qian, 2012, An advanced electron spin resonance (ESR) spin trapping and LC (ESR) MS technique for the study of lipid peroxidation. *Int J Mol Sci*, 13:14648_66.

Xu, Y., Y. Gu, S.Y. Qian, 2012, An advanced electron spin resonance (ESR) spin trapping and LC (ESR) MS technique for the study of lipid peroxidation. *Int J MolSci*, 13:14648 - 66.

Yadav, A., R., Kumari, A., Yadav, J.P. Mishra, S. Srivatva, S. Prabha, 2016, Antioxidants and its functions in human body - A Review, *Res. Environ. Life Sci.* 9(11) 1328-1331

Yefrida, M., Ulfaningsih, Refilda, 2014, Validasi metode penentuan antioksidan total (dihitung sebagai asam sitrat) dalam sampel jeruk secara spektrofotometri dengan menggunakan oksidator FeCl₃ dan peng kompleks orto-fenantrolin, *J. Riset Kimia*, 7(2), 186-193

Yefrida, N., Ashikin, Refilda, 2015, Validasi metode FRAP modifikasi pada penentuan kandungan antioksidan total dalam sampel mangga dan rambutan, *J. Riset Kimia*, 8(2), 170-175

Yefrida, Y., H. Suyani, H. Aziz, M. Efdi, 2018, Comparison of Iron Reduction Methods on the Determination of Antioxidants Content in Vegetables Sample. *Oriental Journal of Chemistry*, 34 (5), 2418-2424, <https://doi.org/10.13005/ojc/340525>

Yefrida, Y., H. Suyani, A. Alif, H. Aziz, M. Efdi, 2018, Modification of Phenanthroline Method to Determine Antioxidant Content in Tropical Fruits Methanolic Extract, *Res. J. Chem. Environ.*, 22 (4), 28-35

Zhang, X., Z. Lin, J. Fang, M. Liu, Y. Niu, S. Chen, H. Wang, 2015, An on-line high-performance liquid chromatography-diode-arraydetector-electrospray ionization-ion-trap-time-of-flight-massspectrometry-total antioxidant capacity detection