

## DAFTAR PUSTAKA

- Aliasgharpour, M., Zabolypour, S., Asadinoghabi, A., Haghani, H., & Lesanpezeshki, M. (2018). The Effect of Increasing Blood Flow Rate on Severity of Uremic Pruritus in Hemodialysis Patients: A Single Clinical Trial. *Journal of the National Medical Association*, 110(3), 270–275. <https://doi.org/10.1016/j.jnma.2017.04.008>
- Anding, K., Bär, T., Trojniak-Hennig, J., Kuchinke, S., Krause, R., Rost, J. M., & Halle, M. (2015). A structured exercise programme during haemodialysis for patients with chronic kidney disease: Clinical benefit and long-term adherence. *BMJ Open*, 5(8), 1–9. <https://doi.org/10.1136/bmjopen-2015-008709>
- Arici, M. (2014). *Management of Chronic Kidney Disease*. London: Springer. <https://doi.org/10.1007/978-3-642-54637-2>
- Artom, M., Moss-Morris, R., Caskey, F., & Chilcot, J. (2014). Fatigue in advanced kidney disease. *Kidney International*, 86(3), 497–505. <https://doi.org/10.1038/ki.2014.86>
- Ashley, C., & Morlidge, C. (2008). *Introduction to Renal Therapeutics* (First). London: Pharmaceutical Press.
- Avesani, C. M., Trolonge, S., Deleaval, P., Baria, F., Mafra, D., Faxen-Irving, G., ... Fouque, D. (2012). Physical activity and energy expenditure in haemodialysis patients: an international survey. *Nephrology, Dialysis, Transplantation: Official Publication of the European Dialysis and Transplant Association - European Renal Association*, 27(6), 2430–2434. <https://doi.org/10.1093/ndt/gfr692>
- Barcellos, F. C., Santos, I. S., Umpierre, D., Bohlke, M., & Hallal, P. C. (2015). Effects of exercise in the whole spectrum of chronic kidney disease: a systematic review. *Clinical Kidney Journal*, 8(6), 753–765. Retrieved from <http://dx.doi.org/10.1093/ckj/sfv099>
- Barzegar, H., Moosazadeh, M., Jafari, H., & Esmaeili, R. (2016). Evaluation of dialysis adequacy in hemodialysis patients: A systematic review. *Urology Journal*, 13(4), 2744–2749.

- Black, J M, & Hawks, J. H. (2009). *Medical Surgical Nursing Clinical Management for Positive Outcome* (8th ed.). St.Louis: Elsevier.
- Black, Joyce M, & Hawks, J. H. (2014). *Keperawatan Medikal Bedah: Manajemen Klinis untuk Hasil yang Diharapkan*. (A. Susila, F. Ganiajri, P. P. Lestari, & R. W. Sari, Eds.) (8th ed.). Jakarta: Salemba Medika.
- Bonner, A., Wellard, S., & Caltabiano, M. (2010). The impact of fatigue on daily activity in people with chronic kidney disease. *Journal of Clinical Nursing*, 19(21–22), 3006–3015. <https://doi.org/10.1111/j.1365-2702.2010.03381.x>
- Borg, G. (1990). Psychophysical scaling with applications in physical work and the perception of exertion. *Scandinavian Journal of Work, Environment & Health*, 16 Suppl 1, 55–58.
- Borzou, S. R., Gholyaf, M., Zandiha, M., Amini, R., Goodarzi, M. T., & Torkaman, B. (2009). The effect of increasing blood flow rate on dialysis adequacy in hemodialysis patients. *Saudi Journal of Kidney Diseases and Transplantation : An Official Publication of the Saudi Center for Organ Transplantation, Saudi Arabia*, 20(4), 639–642.
- Bossola, M., Di Stasio, E., Antocicco, M., Panico, L., Pepe, G., & Tazza, L. (2015). Fatigue Is Associated with Increased Risk of Mortality in Patients on Chronic Hemodialysis. *Nephron*, 130(2), 113–118. <https://doi.org/10.1159/000430827>
- Bossola, M., Vulpio, C., & Tazza, L. (2011). Fatigue in Chronic Dialysis Patients. *Seminars in Dialysis*, 24(5), 550–555. <https://doi.org/10.1111/j.1525-139X.2011.00956.x>
- Broers, N. J. H., Martens, R. J. H., Ornelis, T., Van Der Sande, F. M., Diederen, N. M. P., Hermans, M. M. H., ... Kooman, J. P. (2017). Physical Activity in End-Stage Renal Disease Patients: The Effects of Starting Dialysis in the First 6 Months after the Transition Period. *Nephron*, 137(1), 47–56. <https://doi.org/10.1159/000476072>
- Brown, P. D. S., Rowed, K., Shearer, J., MacRae, J. M., & Parker, K. (2017). Impact of intradialytic exercise intensity on urea clearance in hemodialysis patients. *Applied Physiology, Nutrition, and Metabolism*, 43(1), 101–104. <https://doi.org/10.1139/apnm-2017-0460>
- Canadian Institute for Health Information. (2015). *Canadian Organ Replacement Register Annual Report: Treatment of End-Stage Organ Failure in Canada*,

2004 to 2013. Ottawa.  
[https://doi.org/https://secure.cihi.ca/free\\_products/2011\\_CORR\\_Annua\\_Report\\_EN.pdf](https://doi.org/https://secure.cihi.ca/free_products/2011_CORR_Annua_Report_EN.pdf)

Caplin, B., Kumar, S., & Davenport, A. (2011). Patients' perspective of haemodialysis-associated symptoms. *Nephrology Dialysis Transplantation*, 26(8), 2656–2663. <https://doi.org/10.1093/ndt/gfq763>

CDC: Centers for Disease Control and Prevention. (2015). Target Heart Rate and Estimated Maximum Heart Rate. Retrieved from <https://www.cdc.gov/physicalactivity/basics/measuring/heartrate.htm>

Chang, Y., Cheng, S. Y., Lin, M., Gau, F. Y., & Chao, Y. F. C. (2010). The effectiveness of intradialytic leg ergometry exercise for improving sedentary life style and fatigue among patients with chronic kidney disease: A randomized clinical trial. *International Journal of Nursing Studies*, 47(11), 1383–1388. <https://doi.org/10.1016/j.ijnurstu.2010.05.002>

Chen, C.-K., Tsai, Y.-C., Hsu, H.-J., Wu, I.-W., Sun, C.-Y., Chou, C.-C., ... Wang, L.-J. (2010). Depression and suicide risk in hemodialysis patients with chronic renal failure. *Psychosomatics*, 51(6), 528-528.e6. <https://doi.org/10.1176/appi.psy.51.6.528>

Clarke, A. L., Young, H. M. L., Hull, K. L., Hudson, N., Burton, J. O., & Smith, A. C. (2015). Motivations and barriers to exercise in chronic kidney disease: A qualitative study. *Nephrology Dialysis Transplantation*, 30(11), 1885–1892. <https://doi.org/10.1093/ndt/gfv208>

Collins, A. J., Foley, R. N., Gilbertson, D. T., & Chen, S.-C. (2015). United States Renal Data System public health surveillance of chronic kidney disease and end-stage renal disease. *Kidney International Supplements*, 5(1), 2–7. <https://doi.org/10.1038/kisup.2015.2>

Costanzo, L. S. (2012). *Essential Fisiologi Kedokteran*. (M. Wiyanto, Ed.) (5th ed.). Jakarta: Binarupa Aksara.

Couser, W. G., Remuzzi, G., Mendis, S., & Tonelli, M. (2011). The contribution of chronic kidney disease to the global burden of major noncommunicable diseases. *Kidney International*, 80(12), 1258–1270. <https://doi.org/10.1038/ki.2011.368>

Dahlan, M. S. (2014). *Statistik untuk Kedokteran dan Kesehatan: deskriptif, bivariat, dan multivariat, dilengkapi aplikasi menggunakan SPSS* (6 Seri 1). Jakarta978-: Epidemiologi Indonesia.

Daugirdas, J. T. (1995). Simplified equations for monitoring Kt/V, PCRn, eKt/V, and ePCRn. *Advances in Renal Replacement Therapy*, 2(4), 295–304.

Daugirdas, J. T., Blake, P. G., & Ing, T. S. (2015). *Handbook of Dialysis*. (J. Goolsby, Ed.) (Fifth). Philadelphia: Wolters Kluwer Health.

Davis, M. P., & Walsh, D. (2010). Mechanisms of Fatigue. *The Journal of Supportive Oncology J Support Oncol*, 8(November), 164–174. <https://doi.org/10.1016/B978-0-443-07427-1.50014-5>

Dharma, K. K. (2011). *Metodologi Penelitian Keperawatan: Panduan Melaksanakan dan Menerapkan Hasil Penelitian*. Jakarta: Trans Info Media.

Dobsak, P., Homolka, P., Svojanovsky, J., Reichertova, A., Soucek, M., Novakova, M., ... Siegelova, J. (2012). Intra-dialytic electrostimulation of leg extensors may improve exercise tolerance and quality of life in hemodialyzed patients. *Artificial Organs*, 36(1), 71–78. <https://doi.org/10.1111/j.1525-1594.2011.01302.x>

EBMcalc system. (2017). Kt/V Dialysis Dose Formulae MultiCalc. Retrieved from [https://www.merckmanuals.com/medical-calculators/Kt\\_V\\_Formulae\\_Multi.htm](https://www.merckmanuals.com/medical-calculators/Kt_V_Formulae_Multi.htm)

Eckardt, K. U., Coresh, J., Devuyst, O., Johnson, R. J., Köttgen, A., Levey, A. S., & Levin, A. (2013). Evolving importance of kidney disease: From subspecialty to global health burden. *The Lancet*, 382(9887), 158–169. [https://doi.org/10.1016/S0140-6736\(13\)60439-0](https://doi.org/10.1016/S0140-6736(13)60439-0)

El-Sheikh, M., & El-Ghazaly, G. (2016). Assessment of hemodialysis adequacy in patients with chronic kidney disease in the hemodialysis unit at Tanta University Hospital in Egypt. *Indian Journal of Nephrology*, 26(6), 398–404. <https://doi.org/10.4103/0971-4065.168141>

El Shemy, M. B. A., ElghiteElhossinyElkazeh, E. A., & RadwaELkually, R. (2016). Efficacy Of Exercise Program In Intra Hemodialysis On patients' Quality Of Life. *IOSR Journal of Nursing and Health Science (IOSR-JNHS)*, 5(1), 17–30. <https://doi.org/10.9790/1959-05151730>

Foley, R. N., & Collins, A. J. (2007). End-Stage Renal Disease in the United States: An Update from the United States Renal Data System. *Journal of the American Society of Nephrology*, 18(10), 2644–2648. <https://doi.org/10.1681/ASN.2007020220>

Fouque, D., Kalantar-Zadeh, K., Kopple, J., Cano, N., Chauveau, P., & Cuppari, L. (2008). A proposed nomenclature and diagnostic criteria for protein-energy wasting in acute and chronic kidney disease. *Kidney Int*, 73. <https://doi.org/10.1038/sj.ki.5002585>

Garfinkel, M. S., Singhal, A., Katz, W. A., Allan, D. A., Reshetar, R., & Schumacher, H. R. J. (1998). Yoga-based intervention for carpal tunnel syndrome: a randomized trial. *JAMA*, 280(18), 1601–1603.

Gellish, R. L., Goslin, B. R., Olson, R. E., McDonald, A., Russi, G. D., & Moudgil, V. K. (2007). Longitudinal modeling of the relationship between age and maximal heart rate. *Medicine and Science in Sports and Exercise*, 39(5), 822–829. <https://doi.org/10.1097/mss.0b013e31803349c6>

Giannaki, C. D., Hadjigeorgiou, G. M., Karatzafiri, C., Maridaki, M. D., Koutedakis, Y., Founta, P., ... Sakkas, G. K. (2013). A single-blind randomized controlled trial to evaluate the effect of 6 months of progressive aerobic exercise training in patients with uraemic restless legs syndrome. *Nephrology Dialysis Transplantation*, 28(11), 2834–2840. Retrieved from <http://dx.doi.org/10.1093/ndt/gft288>

Giannaki, C. D., Stefanidis, I., Karatzafiri, C., Liakos, N., Roka, V., Ntente, I., & Sakkas, G. K. (2011). The effect of prolonged intradialytic exercise in hemodialysis efficiency indices. *American Society of Artificial Internal Organs Journal*, 57, 213–218. <https://doi.org/10.1097/MAT.0b013e318215dc9e>

Girija, K., & Radha, R. (2013). Beneficial Effect of Physical Activity in Hemodialysis Patients. *Universal Journal of Engineering Science*, 1(2), 40–44. <https://doi.org/10.13189/ujes.2013.010204>

Gordon, P. L., Doyle, J. W., & Johansen, K. L. (2011). Postdialysis fatigue is associated with sedentary behavior. *Clinical Nephrology*, 75(5), 426–433.

Grassmann, A., Gioberge, S., Moeller, S., & Brown, G. (2005). ESRD patients in 2004: Global overview of patient numbers, treatment modalities and associated trends. *Nephrology Dialysis Transplantation*, 20(12), 2587–2593.

<https://doi.org/10.1093/ndt/gf159>

Greenwood, S. A., Naish, P., Clark, R., O'Connor, E., Pursey, V. A., Macdougall, I. C., ... Koufaki, P. (2014). INTRA-DIALYTIC EXERCISE TRAINING: A PRAGMATIC APPROACH. *Journal of Renal Care*, 40(3), 219–226. <https://doi.org/10.1111/jorc.12080>

Gulati, M., Shaw, L. J., Thisted, R. A., Black, H. R., Bairey Merz, C. N., & Arnsdorf, M. F. (2010). Heart rate response to exercise stress testing in asymptomatic women: the st. James women take heart project. *Circulation*, 122(2), 130–137. <https://doi.org/10.1161/CIRCULATIONAHA.110.939249>

Hartanti, R. D. (2016). Exercise Introdialisis Meningkatkan Nilai URR Pasien Gagal Ginjal Kronik Dengan Hemodialisis. *The 3 Rd University Research Colloquium 2016*, 533-541. ISSN 2407-9189.

Hassanpour-Dehkordi, A., & Jalali, A. (2016). Effect of Progressive Muscle Relaxation on the Fatigue and Quality of Life Among Iranian Aging Persons. *Acta Medica Iranica*, 54(7), 430–436. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/27424013>

Hastono, S. P., & Sabri, L. (2011). *Statistik Kesehatan* (ke-6). Jakarta: PT. Raja Grafindo Persada, Jakarta.

Heidenheim, A. P., Muirhead, N., Moist, L., & Lindsay, R. M. (2003). Patient quality of life on quotidian hemodialysis. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 42(1 Suppl), 36–41.

Hemmelgarn, B. R., Manns, B. J., Lloyd, A., James, M. T., Klarenbach, S., Quinn, R. R., ... Tonelli, M. (2010). Relation between kidney function, proteinuria, and adverse outcomes. *JAMA - Journal of the American Medical Association*, 303(5), 423–429. <https://doi.org/10.1001/jama.2010.39>

Hendriks, C., Drent, M., Elfferich, M., & De Vries, J. (2018). The Fatigue Assessment Scale: quality and availability in sarcoidosis and other diseases. *Current Opinion in Pulmonary Medicine*, 24(5), 495–503. <https://doi.org/10.1097/MCP.0000000000000496>


Hill, N. R., Fatoba, S. T., Oke, J. L., Hirst, J. A., Callaghan, A. O., Lasserson, D. S., & Hobbs, F. D. R. (2016). Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis. *PLOS ONE*, 1–18.

<https://doi.org/10.5061/dryad.3s7rd.Funding>

Horigan, A. E. (2012). Fatigue in hemodialysis patients: A review of current knowledge. *Journal of Pain and Symptom Management*, 44(5), 715–724. <https://doi.org/10.1016/j.jpainsympman.2011.10.015>

Hyun, Y. Y., Lee, K. B., Han, S. H., Kim, Y. H., Kim, Y. S., Lee, S. W., ... Ahn, C. (2017). Nutritional status in adults with predialysis chronic kidney disease: KNOW-CKD study. *Journal of Korean Medical Science*, 32(2), 257–263. <https://doi.org/10.3346/jkms.2017.32.2.257>

Indonesian Renal Registry. (2015). *8th Report Of Indonesian Renal Registry. Program Indonesia Renal Registry*. <https://doi.org/10.2215/CJN.02370316>



Jadeja, Y. P., & Kher, V. (2012). Protein energy wasting in chronic kidney disease: An update with focus on nutritional interventions to improve outcomes. *Indian Journal of Endocrinology and Metabolism*, 16(2), 246–251. <https://doi.org/10.4103/2230-8210.93743>

Jhamb, M., Pike, F., Ramer, S., Argyropoulos, C., Steel, J., Dew, M. A., ... Unruh, M. (2011). Impact of fatigue on outcomes in the hemodialysis (HEMO) study. *American Journal of Nephrology*, 33(6), 515–523. <https://doi.org/10.1159/000328004>



Jhamb, M., Weisbord, S. D., Steel, J. L., & Unruh, M. (2008). Fatigue in Patients Receiving Maintenance Dialysis: A Review of Definitions, Measures, and Contributing Factors. *American Journal of Kidney Diseases*, 52(2), 353–365. <https://doi.org/10.1053/j.ajkd.2008.05.005>



Johansen, K. L., Finkelstein, F. O., Revicki, D. A., Evans, C., Wan, S., Gitlin, M., & Agodoa, I. L. (2012). Systematic review of the impact of erythropoiesis-stimulating agents on fatigue in dialysis patients. *Nephrology Dialysis Transplantation*, 27(6), 2418–2425. <https://doi.org/10.1093/ndt/gfr697>

Kalantar-Zadeh, K., & Unruh, M. (2005). Health related quality of life in patients with chronic kidney disease. *International Urology and Nephrology*, 37(2), 367–378. <https://doi.org/10.1007/s11255-004-0012-4>

Kallenbach, Gutch, J. Z., Stoner, C., & Corea, M. (2005). *Review of Hemodialysis for Nurses And Dialysis Personnel* (7th ed.). Elsevier.

- Karakan, S., Sezer, S., & Ozdemir, F. N. (2011). Factors related to fatigue and subgroups of fatigue in patients with end-stage renal disease. *Clinical Nephrology*, 76(5), 358–364.
- KDIGO. (2013). Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Journal of The International Society of Nephrology*, 3(1), Supplement. <https://doi.org/10.1038/kisup.2012.76>
- Kemenkes RI. (2013). Riset Kesehatan Dasar 2013. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. Retrieved from [www.depkes.go.id/resources/download/general/Hasil\\_Riskesdas\\_2013.pdf](http://www.depkes.go.id/resources/download/general/Hasil_Riskesdas_2013.pdf)
- Kemenkes RI. (2017). Situasi Penyakit Ginjal Kronis. InfoDatin Pusat Data Dan Informasi Kemenkes RI. ISSN 2442-7659. Retrieved from <http://www.pusdatin.kemkes.go.id/folder/view/01/structure-publikasi-pusdatin-info-datin.html>
- Khalaj, A., Sanavi, S., Afshar, R., & Rajabi, M. (2010). Effect of Intradialytic Change in Plasma Volume on Blood Pressure in Patients Undergoing Maintenance Hemodialysis. *Journal of Laboratory Physicians*, 2(2), 66–69. <https://doi.org/10.4103/0974-2727.72151>
- Kirkman, D. L., Roberts, L. D., Kelm, M., Wagner, J., Jibani, M. M., & Macdonald, J. H. (2014). Interaction between intradialytic exercise and hemodialysis adequacy. *American Journal of Nephrology*, 38(6), 475–482. <https://doi.org/10.1159/000356340>
- Koh, K. P., Fassett, R. G., Sharman, J. E., Coombes, J. S., & Williams, A. D. (2009). Intradialytic versus home based exercise training in hemodialysis patients: A randomised controlled trial. *BMC Nephrology*, 10(1), 1–6. <https://doi.org/10.1186/1471-2369-10-2>
- Kong, C. H., Tattersall, J. E., Greenwood, R. N., & Farrington, K. (1999). The effect of exercise during haemodialysis on solute removal. *Nephrology Dialysis Transplantation*, 14, 2927–2931.
- Kosmadakis, G. C., Bevington, A., Smith, A. C., Clapp, E. L., Viana, J. L., Bishop, N. C., & Feehally, J. (2010). Physical exercise in patients with severe kidney disease. *Nephron - Clinical Practice*, 115(1), 7–16. <https://doi.org/10.1159/000286344>

- Larun, L., Brurberg, K. G., Odgaard-Jensen, J., & Price, J. R. (2015). Exercise therapy for chronic fatigue syndrome (Review). *The Cochrane Library*, (2). <https://doi.org/10.1002/14651858.CD003200.pub6.www.cochranelibrary.com>
- Laville, M., & Fouque, D. (2000). Nutritional aspects in hemodialysis. *Kidney International. Supplement*, 76, S133-9.
- Levy, J., Morgan, J., & Brown, E. a. (2016). *Oxford Handbook of Dialysis. Oxford Medicine Online* (Fourth). United Kingdom: Oxford University Press. <https://doi.org/10.1093/med/9780199235285.001.0001>
- Liao, M.-T., Liu, W.-C., Lin, F.-H., Huang, C.-F., Chen, S.-Y., Liu, C.-C., ... Wu, C.-C. (2016). Intradialytic aerobic cycling exercise alleviates inflammation and improves endothelial progenitor cell count and bone density in hemodialysis patients. *Medicine*, 95(27), e4134. <https://doi.org/10.1097/MD.0000000000004134>
- Lindsay, R. M., Heidenheim, P. A., Nesrallah, G., Garg, A. X., & Suri, R. (2006). Minutes to recovery after a hemodialysis session: a simple health-related quality of life question that is reliable, valid, and sensitive to change. *Clinical Journal of the American Society of Nephrology : CJASN*, 1(5), 952–959. <https://doi.org/10.2215/CJN.00040106>
- Lindsay, R. M., & Kortas, C. (2001). Hemeral (Daily) Hemodialysis. *Advances in Renal Replacement Therapy*, 8(4), 236–249. <https://doi.org/10.1053/jarr.2001.27593>
- Liu, H. . (2006). Fatigue and Associated Factors in Hemodialysis Patients in Taiwan. *Research in Nursing & Health*, 40–50. <https://doi.org/10.1002/nur.20109>
- Locatelli, F., Buoncristiani, U., Canaud, B., Köhler, H., Petitclerc, T., & Zucchelli, P. (2005). Dialysis dose and frequency. *Nephrology Dialysis Transplantation*, 20(2), 285–296. Retrieved from <http://dx.doi.org/10.1093/ndt/gfh550>
- MacDonald, J. H., Fearn, L., Jibani, M., & Marcra, S. M. (2012). Exertional fatigue in patients with CKD. *American Journal of Kidney Diseases*, 60(6), 930–939. <https://doi.org/10.1053/j.ajkd.2012.06.021>
- Maharani, D. (2017). Penyakit Ginjal Sedot Dana BPJS Lebih Dari Rp 2 Triliun. *Kompas*. Retrieved from

<https://sains.kompas.com/read/2017/03/09/120300123/penyakit.ginjal.sedot.dana.bpjs.lebih.dari.rp.2.triliun>

Mahon, A., Jenkins, K., & Burnapp, L. (2013). *Oxford Handbook of Renal Nursing* (First). United Kingdom: Oxford University Press.  
<https://doi.org/10.1093/med/9780199600533.001.0001>

Mahrova, A., & Svagrova, K. (2013). Exercise Therapy - Additional Tool for Managing Physical and Psychological Problems on Hemodialysis. In *Hemodialysis* (pp. 753–821). INTECH.  
<https://doi.org/http://dx.doi.org/10.5772/53058>

Makhlough, A., Ilali, E., Mohseni, R., & Shahmohammadi, S. (2012). Effect of Intradialytic Aerobic Exercise on Serum Electrolytes Levels in Hemodialysis Patients. *Iranian Journal of Kidney Diseases*, 6(2), 119–123.

Mathew, A., & Lathas S, L. S. (2014). Effectiveness Of Intradialytic Exercise On Fatigue And Quality Of Life Among Chronic Renal Failure Patients Undergoing Hemodialysis In A Selected Hospital At Mangalore. *Journal Of International Academic Research For Multidisciplinary*, 2(7), 418–424.

MedCalc.com. (2018). MedCalc: Hemodialysis - Kt/V and URR. Retrieved from <http://www.medcalc.com/hd.html>

Mohseni, R., Emami Zeydi, A., Ilali, E., Adib-Hajbaghery, M., Makhlough, A., & Student, G. (2013). The Effect of Intradialytic Aerobic Exercise on Dialysis Efficacy in Hemodialysis Patients: A Randomized Controlled Trial. *Oman Medical Specialty Board Oman Medical Journal*, 28(5), 345–349.  
<https://doi.org/10.5001/omj.2013.99>

Moledina, D. G., & Wilson, F. P. (2015). Pharmacologic Treatment of Common Symptoms in Dialysis Patients: A Narrative Review. *Seminars in Dialysis*, 28(4), 377–383. <https://doi.org/10.1111/sdi.12378>


Morishita, S., Tsubaki, A., & Shirai, N. (2017). Physical function was related to mortality in patients with chronic kidney disease and dialysis. *Hemodialysis International*, 21(4), 483–489. <https://doi.org/10.1111/hdi.12564>

Motedayen, Z., Nehrir, B., Tayebi, A., Ebadi, A., & Einollahi, B. (2014). The Effect of the Physical and Mental Exercises During Hemodialysis on fatigue: A Controlled Clinical Trial. *Nephro-Urology Monthly*, 6(4), 1–6.  
<https://doi.org/https://doi.org/DOI: 10.5812/numonthly.14686>

Mula-Abed, W.-A. S., Rasadi, K. Al, & Riyami, D. Al. (2012). Estimated Glomerular Filtration rate (eGFR): A Serum Creatinine-based Test for the Detection of Chronic Kidney Disease and its Impact on Clinical Practice. *Oman Medical Journal*, 27(4), 339–340. <https://doi.org/10.5001/omj.2012.87>


Murtagh, F. E. M., Addington-Hall, J., & Higginson, I. J. (2007). The Prevalence of Symptoms in End-Stage Renal Disease: A Systematic Review. *Advances in Chronic Kidney Disease*, 14(1), 82–99. <https://doi.org/10.1053/j.ackd.2006.10.001>

National Kidney Foundation. (2006). Hemodialysis Adequacy 2006. *American Journal of Kidney Diseases*, 48, S2–S90. <https://doi.org/10.1053/j.ajkd.2006.03.051>



National Kidney Foundation. (2015). Clinical Practice Guideline for Hemodialysis Adequacy: 2015 Update. *American Journal of Kidney Diseases*, 66(5). <https://doi.org/10.1053/j.ajkd.2015.07.015>

Nelson, V., Gopalakrishnan, S., Simon, S., Babu, V., Vikraman, V., Abraham, S., ... Medical, T. (2016). Depression among dialysis patients, 40(2), 34–37.



Nissenson, A. R., & Fine, R. N. (2017). *Handbook of Dialysis Therapy*. (A. R. Nissenson & R. N. Fine, Eds.) (Fifth). Philadelphia: Elsevier.



Painter, P. (2000). *Exercise: A Guide for the People on Dialysis* (Vol. 1). The Life Options Rehabilitation Advisory Council. <https://doi.org/10.1017/CBO9781107415324.004>



Paluchamy, T., & Vaidyanathan, R. (2018). Effectiveness of intradialytic exercise on dialysis adequacy, physiological parameters, biochemical markers and quality of life - A pilot study. *Saudi Journal of Kidney Diseases and Transplantation*, 29(4), 902–910. <https://doi.org/10.4103/1319-2442.239661>



Parsons, T. L., Toffelmire, E. B., & King-VanVlack, C. E. (2006). Exercise Training During Hemodialysis Improves Dialysis Efficacy and Physical Performance. *Archives of Physical Medicine and Rehabilitation*, 87(5), 680–687. <https://doi.org/10.1016/j.apmr.2005.12.044>



Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia (PAPDI). (2012). *Konsensus Nutrisi Enteral & Parenteral Pada Bidang Ilmu Penyakit Dalam*. (A. F. Syam, E. J. Nelwan, H. Rasyid, & L. Sukrisman, Eds.). Jakarta: Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia.

Persatuan Nefrologi Indonesia. (2003). *Konsensus Dialisis Pernefri*. (Persatuan Nefrologi Indonesia, Ed.) (1st ed.). Jakarta: PERNEFRI.

Persatuan Nefrologi Indonesia. (2011). *Konsensus Manajemen Anemia Pada Penyakit Ginjal Kronik* (1st ed.). Jakarta: PERNEFRI.

Pescatello, L. S., Arena, R., Riebe, D., & Thompson, P. D. (2013). *ACSM's Guidelines for Exercise Testing and Prescription* (ninth). Wolters Kluwer Health.

Price, S. A., & Wilson, L. M. (2015). *Patofisiologi Konsep Klinis Proses-Proses Penyakit* (Edisi 6 Vo). Jakarta: EGC.

Quinn, E. (2017). How to Use the Perceived Exertion Scale During Your Workout: Estimating Heart Rate and Exercise Intensity Zones. Retrieved August 10, 2018, from <https://www.verywellfit.com/rating-of-perceived-exertion-scale-3119445>

Raghavendra, V., Nagendra, N., Patil, S., Diwakar, R. B., Shashidhara, H. P., Gopinath, K. S., & Ajaikumar, B. (2017). Effects of Yoga in Managing Fatigue in Breast Cancer Patients: A Randomized Controlled TRial. *Indian Journal of Palliative Care*, 23(3), 247–252. <https://doi.org/10.4103/IJPC.IJPC9517>

Raiesifar, A., Torabpour, M., Mohsenizad, P., Shabani, H., Tayebi, A., & Masoumi, M. (2009). Dialysis adequacy in patients of Abadan hemodialysis center. *Iranian Journal of Critical Care Nursing Fall Issue*, 2.

Ream, E., & Richardson, A. (1996). Fatigue: a concept analysis. *International Journal of Nursing Studies*, 33(5), 519–529.

Resic, H., Vavra-Hadziahmetovic, N., Celik, D., Kablar, A., Kukavica, N., Masnic, F., ... Coric, A. (2014). The effect of intradialytic exercise program on the quality of life and physical performance in hemodialysis patients. *Acta medica Croatica : casopis Hrvatske akademije medicinskih znanosti*, 68(2), 79–84.

Riebe, D., Ehrman, J. K., Liguori, G., & Magal, M. (2016). *Guidelines for Exercise Testing and Prescription*. (D. Riebe, J. K. Ehrman, G. Liguori, & M. Magal, Eds.) (Tenth). Wolters Kluwer.

Ronny, R., Setiawan, S., & Fatimah, S. (2010). *Fisiologi Kardiovaskuler: Berbasis Masalah Keperawatan*. (E. Wahyuningsih, Ed.). Jakarta: EGC.

Sakitri, G., Makiyah, N., & Khoiriayati, A. (2017). Pengaruh Intradialytic Exercise Terhadap Fatigue Pasien Hemodialisis Di RSUP dr . Soeradji Tirtonegoro Klaten. *Media Publikasi Penelitian*, 15(1), 58–64. Retrieved from <https://ejournal.stikespu.ac.id/index.php/mpp/article/viewFile/58/172>

Sakkas, G. K., & Karatzasferi, C. (2012). Hemodialysis fatigue: Just “simple” fatigue or a syndrome on its own right? *Frontiers in Physiology*, 3(July), 1–5. <https://doi.org/10.3389/fphys.2012.00306>

Sanathan, S. R., Bharathan, V., Alla, P., Madhuri, S., Shetty, M. S., & Ram, D. (2014). Depressive symptoms in chronic kidney disease patients on maintenance hemodialysis. *World Journal of Pharmacy and Pharmaceutical Sciences*, 3(8), 535–548. <https://doi.org/ISSN 2278 - 4357>

Sanavi, S., Afshar, R., Shegarfy, L., & Shavandi, N. (2010). Effects of aerobic exercise and resistance training on lipid profiles and inflammation status in patients on maintenance hemodialysis. *Indian Journal of Nephrology*, 20(4), 185. <https://doi.org/10.4103/0971-4065.73442>

Santoso, S. (2012). *Panduan Lengkap SPSS Versi 20* (1st ed.). Jakarta: PT Elex Media Komputindo Kompas Gramedia.

Sawant, A., Garland, S. J., House, A. A., & Overend, T. J. (2011). Morphological, electrophysiological, and metabolic characteristics of skeletal muscle in people with end-stage renal disease: a critical review. *Physiotherapy Canada*, 63(3), 355–376. <https://doi.org/10.3138/ptc.2010-18>

Shahid, A., Wilkinson, K., Marcu, S., & Shapiro, C. M. (2012). *STOP, THAT and ONE Hundred Other Sleep Scales*. Springer Science + Business Media, LLC. [https://doi.org/10.1007/978-1-4419-9893-4\\_33](https://doi.org/10.1007/978-1-4419-9893-4_33)

Shariati AR, Asayesh H, Nasiri H, Tajbakhsh R, Hesam M, Mollaee E, ... Nik Sier M. (2012). Comparison of dialysis adequacy in patient's that referred to Golestan province hemodialysis centers. *Journal of Health Promotion Management*, 1(3), 55–63. Retrieved from <http://jhpm.ir/article-1-82-en.html>

Silverman, M. N., & Deuster, P. A. (2014). Biological mechanisms underlying the role of physical fitness in health and resilience. *Interface Focus*, 4(5), 20140040. <https://doi.org/10.1098/rsfs.2014.0040>

Singarimbun, M., & Effendi, S. (2016). *Metode Penelitian Survai*. Jakarta.: PT Midas Surya Grafindo.

Sirajuddin, S., Surmita, & Astuti, T. (2018). *Bahan Ajar Gizi: Survey Konsumsi Pangan.* (A. Suhardianto, Ed.) (1st ed.). Jakarta: Kementerian Kesehatan Republik Indonesia: Pusat Pendidikan Sumber Daya Manusia Kesehatan.

Sklar, A. H., Riesenbergs, L. A., Silber, A. K., Ahmed, W., & Ali, A. (1996). Postdialysis fatigue. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 28(5), 732–736.

Sklar, A., Newman, N., Scott, R., Semenyuk, L., Schultz, J., & Fiacco, V. (1999). Identification of factors responsible for postdialysis fatigue. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 34(3), 464–470. <https://doi.org/10.1053/AJKD03400464>

Smeltzer, S. ., & Bare, B. . (2014). *Textbook of Medical-Surgical Nursing*. Jakarta: EGC.

Soliman, H. M. M. (2015). Effect of intradialytic exercise on fatigue, electrolytes level and blood pressure in hemodialysis patients: A randomized controlled trial. *Journal of Nursing Education and Practice*, 5(11), 16–28. <https://doi.org/10.5430/jnep.v5n11p16>

Taylor, R. R., Jason, L. A., & Torres, A. (2000). Fatigue rating scales: an empirical comparison. *Psychological Medicine*, 30(4), 849–856.

Tentori, F., Elder, S. J., Thumma, J., Pisoni, R. L., Bommer, J., Fissell, R. B., ... Robinson, B. M. (2010). Physical exercise among participants in the Dialysis Outcomes and Practice Patterns Study (DOPPS): Correlates and associated outcomes. *Nephrology Dialysis Transplantation*, 25(9), 3050–3062. <https://doi.org/10.1093/ndt/gfq138>

United States Renal Data System. (2014). *2014 USRDS annual data report: Epidemiology of kidney disease in the United States*. Bethesda. Retrieved from <https://www.usrds.org/2014/view/Default.aspx>

Unruh, M., Miskulin, D., Yan, G., Hays, R. D., Benz, R., Kusek, J. W., & Meyer, K. B. (2004). Racial differences in health-related quality of life among hemodialysis patients. *Kidney International*, 65, 1482–1491. <https://doi.org/10.1111/j.1523-1755.2004.00529.x>

Valliant, A., & Hofmann, R. M. (2013). Managing dialysis patients who develop anemia caused by chronic kidney disease: focus on peginesatide. *International Journal of Nanomedicine*, 8, 3297–3307. <https://doi.org/10.2147/IJN.S44944>

- van Vilsteren, M. C. B. A., de Greef, M. H. G., & Huisman, R. M. (2005). The effects of a low-to-moderate intensity pre-conditioning exercise programme linked with exercise counselling for sedentary haemodialysis patients in The Netherlands: results of a randomized clinical trial. *Nephrology, Dialysis, Transplantation*, 20(1), 141–146. <https://doi.org/10.1093/ndt/gfh560>
- Waehner, P. (2018). Karvonen Formula for Calculating Your Target Heart Rate. Retrieved September 25, 2018, from <https://www.verywellfit.com/karvonen-formula-1229753>
- Weisbord, S. D., Fried, L. F., Arnold, R. M., Fine, M. J., Levenson, D. J., Peterson, R. A., & Switzer, G. E. (2005). Prevalence, Severity, and Importance of Physical and Emotional Symptoms in Chronic Hemodialysis Patients. *Journal of the American Society of Nephrology*, 16(8), 2487–2494. <https://doi.org/10.1681/ASN.2005020157>
- Williams, A. G., Crane, P. B., & Kring, D. (2007). Fatigue in African American women on hemodialysis. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 34(6), 610–617, 644; quiz 618.
- World Kidney Day. (2018). *Chronic Kidney Disease*. Retrieved from <http://www.worldkidneyday.org/faqs/chronic-kidney-disease/>
- Yong, D. S. P., Kwok, A. O. L., Wong, D. M. L., & Centre, C. M. (2009). Symptom burden and quality of life in end-stage renal disease : a study of 179 patients on dialysis and palliative care. *Palliative Medicine*, 111–119. <https://doi.org/10.1177/0269216308101099>
- Zuraida, R., & Chie, H. H. (2014). Pengujian Skala Pengukuran Kelelahan (SPK) Pada Responden Di Indonesia. *ComTech*, 5(9), 1012–1020.
- Zyga, S., Alikari, V., Sachlas, A., Fradelos, E. C., Stathoulis, J., Panoutsopoulos, G., ... Lavdaniti, M. (2015). Assessment of Fatigue in End Stage Renal Disease Patients Undergoing Hemodialysis: Prevalence and Associated Factors. *Medical Archives (Sarajevo, Bosnia and Herzegovina)*, 69(6), 376–380. <https://doi.org/10.5455/medarh.2015.69.376-380>