

## REFERENCES

- Addaha H., D. H. Tjong, W. Novarino. 2014. Variasi Morfologi Katak Pohon Bergaris *Polypedates leucomystax* Gravenhorst, 1829 (Anura: Rhacophoridae) di Sumatera Barat. *Online Journal of Natural Science*, 4(3):348-354.
- Addis, Brett & Lowe, Winsor & Hossack, Blake & Allendorf, Fred. 2015. Population genetic structure and disease in montane boreal toads: more heterozygous individuals are more likely to be infected with amphibian chytrid. *Conservation Genetics*. 16. 10.1007/s10592-015-0704-6.
- Ahluwalia K. B. 2009. *Genetics: Second Edition*. New Delhi: New Age International Publishers.
- Allendorf F.W. and L. Gordon. 2007. *Conservation and the Genetics of Population*. Victoria: Blackswell Publishing.
- Anwar, J, J. Damanik, N Hisyam dan A.J. Whitten. 1984. *Ekologi Ekosistem Sumatera*. Yogyakarta : UG Press.
- Ardila M.Z., Titrawani, E. Cahyadi. 2017. Variasi morfometrik *Fejervarya cancrivora* Gravenhorst (1829) di Kawasan Universitas Riau. *Jurnal Riau Biologia*. retrieved online from Variasi Morfometrik *Fejervarya cancrivora* Gravenhorst (1829) di Kawasan Universitas Riau (unri.ac.id)
- Arifin, Umilaela, S. Utpal., S. Hertwig., E. Smith, D.T. Iskandar, and A. Haas. 2018. Molecular phylogenetic analysis of a taxonomically unstable ranid from Sumatra, Indonesia, reveals a new genus with gastromyzophorous tadpoles and two new species. *Zoosystematics and Evolution*. 94. 10.3897/zse.94.22120.
- Arruda M. P and E. M. Versute 2008. Cytogenetic and Random Amplified Polymorphic DNA Analysis of *Leptodactylus* Species from Rural And Urban Environments (Anura, Amphibia). *Genet. Mol. Res.* 7 (1): 161-176.
- Atlas, J. E., and J. Fu. 2019. Isolation by resistance analysis reveals major barrier effect imposed by the Tsinling Mountains on the Chinese wood frog. *Journal of Zoology*. doi:10.1111/jzo.12702
- Bagley, M. J., E. Susan, S. E. Franson, S. A. Christ, E. R., Waits G. P. Toth. 2002. *Genetic Diversity as an Indicator of Ecosystem Condition and Sustainability: Utility for Regional Assessments of Stream Condition in the Eastern United States*. U.S. Environmental Protection Agency, Cincinnati, OH.

- Bardacki, F. 2001. Random Amplified Polymorphic DNA (RAPD) Markers. *Turk. J. Biol.* 25: 185-196.
- Bird, M.I., D. Taylor. C. Hunt. 2005. Palaeoenvironments of insular Southeast Asia during the Last Glacial Period: a savanna corridor in Sundaland? *Quatern. Sci. Rev.*, 24, 2228–2242.
- Borzée A., S. Y.Park, A. B.Kim, H. T.Kim, Y. KJang. 2013. Morphometrics of two sympatric species of tree frogs in Korea: a morphological key for the critically endangered *Hyla suweonensis* in relation to *H. japonica*. *Anim Cell Syst*, 17(5): 348–356.
- Cao, D. and J.H. Oard, 1997. Pedigree and RAPD-based DNA analysis of commercial U.S. rice cultivars. *Crop Sci.*, 37: 1630-1635. DOI: 10.2135/cropsci1997.0011183X003700050037x
- Che J, Zhou WW, Hu JS, et al. 2010. Spiny frogs (Paini) illuminate the history of the Himalayan region and Southeast Asia. *Proceedings of the National Academy of Sciences of the United States of America*. Aug;107(31):13765-13770. DOI: 10.1073/pnas.1008415107. PMID: 20643945; PMCID: PMC2922240.
- De Bruyn, M., B. Stelbrink, R.J.Morley, R. Hall, G.R. Carvalho, C.H. Cannon, G. van den Bergh, E. Meijaard, I. Metcalfe, L. Boitani, L. Maiorano, R Shoup, and T. von Rintelen. 2014. Borneo and Indochina are Major Evolutionary Hotspots for Southeast Asian Biodiversity. *Systematic Biology*, 63(6), 879–901. doi: 10.1093/sysbio/syu047.
- Diouf, J. 2007. *The State of the World's Animal Genetic Resources for Food and Agriculture*. Rome, Italy. Pp.
- Djong H.T, Matsui M., D. M. Belabut, Y. H. Sen, M. Nishioka and M. Sumida. 2007. Morphological Divergence, Reproductive Isolating Mechanism, and Molecular Phylogenetic Relationships among Indonesia, Malaysia and Japan Populations of The *Fejervarya limnocharis* Complex (Anura: Ranidae). *Zoological Science*, 24(12), 1197-1212.
- Dubois, A., and A. Ohler. 2000. Systematics of *Fejervarya limnocharis* (Gravenhorst, 1829) (Amphibia, Anura, Ranidae) and related species. 1. Nomenclatural status and type-specimens of the nominal species *Rana limnocharis* Gravenhorst, 1829. *Alytes*. Paris 18: 15–50.
- Fatchiyah, Arumingtiyas, Widyawati dan Rahayu. 2011. *Biologi Molekuler*. Erlangga. Jakarta.



- Fauzan. 2009. *Differensiasi Morfometrik Fejervarya limnocharis (Gravenhorst, 1829) Di Sumatera Barat*. [Skripsi]. Jurusan Biologi FMIPA Universitas Andalas, Padang.
- Finkeldey, R. 2005. *An Introduction to Tropical Forest Genetics*. Gottingen: Institute of Forest Genetics and Forest Tree Breeding Georg-August-University Gottingen.
- Frost, D. R. 2007. Amphibian Species of the World: An Online Reference. Version 5.0.
- Fukuoka, S., K. Hosaka and O. Kamijima, 1992. Use of Random Amplified Polymorphic DNAs (RAPDs) for identification of rice accessions. *Japanese J.Genet.*, 67: 243-252.
- Funk, W.C., M. S. Blouin, P. S. Corn, B. A. Maxell, D. S. Pilliod, S. Amish and F.W. Allendorf. 2005. Population Structure of Columbia Spotted Frogs (*Rana luteiventris*) is Strongly Affected by The Landscape. *MolecularEcology*, 14: 483-496.
- Gravenhorst, J. L. C. 1829. *Deliciae Musei Zoologici Vratislaviensis. Fasciculus primus. Chelonios et Batrachia*. Leipzig: Leopold Voss. Leipzig 1, 41 pp.
- Gurdebeke, S., J. P., Maelfait and T. Backeljau. 2003. Contrasting Allozyme and RAPD Variation in Spider Populations from Patchy Forest Habitats. *Genetica*, 119: 27-34.
- Hall, R., 1996. Reconstructing Cenozoic SE Asia. In: Hall, R. and Blundell, D. (Eds.), *Tectonic Evolution of Southeast Asia*, vol. 106. Geological Society Special Publications, 153–184.
- Hall, R. 2009. Southeast Asia's changing palaeogeography. *Blumea*, 54, 148–161
- Harrison I., M. Laverty, E. Sterling. 2004. Genetic Diversity. *Connexions module*: m12158.
- Hendri, W., D. H. Tjong, Dahelmi, D. I. Roesma. 2019. Genetic Variation of *Limnonectes blythii* (Anura : Dicroglossidae) Using RAPD (Random Amplified Polymorphic DNA) Analysis in West Sumatra. *International Journal of Zoological Research*, 15 (1) : 1-5
- Huang, M.C., Y.M. Horng, H.L. Huang, Y.L. Sin, M.J. Chen. (2003). RAPD Fingerprinting for the Species Identification of Animals. *Asian-Australasian Journal of Animal Sciences*. 16. 10.5713/ajas.2003.1406.

- Ilić, M. S., Gorana, N., V. Marković, N. Paunović., M. C, Jelka. 2016. Identification of syntopic anuran species in early tadpole stages: Correspondence between morphometric and genetic data. *Applied Ecology and Environmental Research*. 14. 381-397. 10.15666/aeer/1402\_381397.
- Iskandar D.T. 1998. *Amfibi Jawa dan Bali*. Bogor: Puslitbang Biologi LIPI.
- Islam M.M., M.M.R. Khan, D.H.Tjong, M.S.Alam, and M.Sumida. 2008. Genetic Differentiation of the *Fejervarya limnocharis* Complex from Bangladesh and Other Asian Countries Elucidated by Allozyme Analyses. *Zoological science*, 25: 261-272.
- IUCN. 2004. *Fejervarya cancrivora*, retrieved online from [Fejervarya cancrivora \(iucnredlist.org\)](http://www.iucnredlist.org)
- Jaeger, R.G. 1994. Patch sampling. In: Heyer, W.R., M.A. Donnelly, R.W. MacDiarmid, L.C. Hayek & M.S. Foster (Eds). *Measuring and Monitoring Biological Diversity: Standard Methods for Amphibians*. Smithsonian Institution Press, Washington DC. Pp. 107-109.
- Joshi, M and J. D. Deshpande. 2010. Polymerase Chain Reaction: Methods, principles And Application. *IJBR* 1 (5): 81-97
- Kimberling D.N., A.R.Ferreira, S.M.Shuster, P.Keim. 1996. RAPD marker estimation of genetic structure among isolated northern leopard frog populations in the south-western USA. *Molecular ecology*, 5: 521-529.
- Kumari, N and S.K. Thakur. 2014. Randomly Amplified Polymorphic DNA – A Brief Review. *American Journal of Animal and Veterinary Sciences*, 9(1) : 6-13.
- Kurniati H., W. Crampton, A. Goodwin, A. Locket and A. Sinkins. 2000. Herpetofauna diversity of Ujung kulon National Park: An inventory results in 1990. *J. Biol. Res.* 6 (2): 113-128.
- Kurniati H. 2016. *Metode Cepat Penentuan Keragaman, Kepadatan dan Kelimpahan Jenis Kodok*. Cibinong: Pusat Penelitian Biologi LIPI.
- Kurniati H. and E.Sulistiyadi. 2017. Kepadatan Populasi Kodok *Fejervarya cancrivora* di Persawahan Kabupaten Karawang, Jawa Barat. *Jurnal Biologi Indonesia* 13(1): 71-83.
- Kurniawan N. M., M. Islam, D. H. Tjong, M.Nishioka. 2010. “Genetic Divergence and Evolutionary Relationship in *Fejervarya cancrivora* from Indonesia and Other Asian Countries Inferred from Allozyme and MtDNA Sequence Analyses.” *Zoological Science*, 27 (3):222–233.



- Kurniawan, N., D.H. Tjong, M.M. Islam, T. Nishizawa, D.M. Belabut, Y.H. Sen, R. Wanichanon, I. Yasir, dan M. Sumida. 2011. Taxonomic Status of Three Types of *Fejervarya cancrivora* from Indonesia and Other Asian Countries Based on Morphological Observations and Crossing Experiments. *Zoological Science* 28 (1) : 12-24.
- Kurniawan, N. H. T. Djong. T. Meideliza. A. Hamidy. M. Hasan, T. Igawa and M. Sumida. 2014. Genetic Divergence and Geographic Distribution of Frogs in Genus *Fejervarya* from Indonesia Inferred from Mitochondrial 16s rRNA Gene Analysis. *Treubia* 41: 1-16
- Kusrini M. D. and R. Alford. 2006. Indonesia's Exports Of Frogs Legs. *Traffic Bulletin*, 21:13-24.
- Larson, A., Chippindale, P. T. 1993. Molecular approaches to the evolutionary biology of plethodontid salamanders. – *Herpetologica* 49: 204-215.
- Laumonier, Y. 1997. *The Vegetation and Physiography of Sumatra*. Kluwer Academic Publishers. Netherlands.
- Liu, Z. J and J. F. Cores. 2004. Review: DNA Marker Technology and Their Applications in Aquaculture Genetics. *Aquaculture*. 238:1-37.
- Matsui M.,D. M.Belabut, N.Ahmad. 2014. “Two new species of fanged frogs from Peninsular Malaysia (Anura: Dicroglossidae).” *Zootaxa*, (1):75–93.
- McKay, J.L. 2006. *A Field Guide to the Amphibians and Reptiles of Bali*. Krieger Publishing Company, Malabar, Florida.
- Méndez M.A., C. Correa, A.Veloso, E.Vergara, Sallaberry, P.Iturra, E. R.Soto. 2004. “Morphological and genetic differentiation among Chilean populations of *Bufo spinulosus* (Anura: Bufonidae). *Revista Chilena de Historia Natural*, 77: 559-567.
- Mikulek, P and J. Pialek. 2003. Molecular Identification of Three Crested Newt Species (*Triturus cristatus* superspecies) by RAPD Markers. *Amphibia-Reptilia*. 24:201-207
- Moslehi, T., M. Mahdieh, A. Shayestehfar and S. M. Talebi. 2015 Genetic Diversity of *Rana (Pelophylax) ridibunda* and *Bufo (Pseudepidelea) viridis* in different populations. *Biodiversitas* Vol.16 No.2 Pages:128-131.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. dan Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.

- Nesty, R., D.H. Tjong, dan H. Herwina. 2013. Variasi Morfometrik Kodok *Duttaphrynus melanostictus* (Schneider, 1799) (Anura : Bufonidae) di Sumatera Barat yang Dipisahkan oleh Bukit Barisan. *Jurnal Biologi Universitas Andalas (J. Bio. UA.)* 2 (1) : 37-42.
- Peakall, R. and Smouse, P.E. 2006 GenAlEx 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol. Ecol. Notes*, 6, 288–295.
- Pemerintah Provinsi Sumatera Barat. 2021. Peraturan Daerah no. 6 tahun 2021 tentang Rencana Pembangunan Jangka Menengah Daerah tahun 2021-2026. Padang, II.8.
- Rasolonjatovo S.M., M.D. Scherz, R. Schmidt, J. Glos, A. Rakotoarison, A.P. Raselimanana and M. Vences. 2022. Population Diversification in the Frog *Mantidactylus bellyi* on an Isolated Massif in Northern Madagascar based on Genetic, Morphological, Bioacoustic and Ecological Evidence. *PLoS ONE* 17(3): e0263764. <https://doi.org/10.1371/journal.pone.0263764>
- Ritu, D and B. Mohapatra. 2018. Genetic Variation. J. Vonk, T. K. Shackelford (eds.), *Encyclopedia of Animal Cognition and Behavior*, [https://doi.org/10.1007/978-3-319-47829-6\\_20-1](https://doi.org/10.1007/978-3-319-47829-6_20-1)
- Sambrook, J., E.F. Fritsch and T. Maniatis, 1989. *Molecular Cloning: A Laboratory Manual*. 2nd Edn., Cold Spring Harbor Laboratory, ISBN-10: 0879693096, pp: 3.
- Shao, Y.W., L.Z. Li., M.C. Hongman., R. Wu, D. Zhang, Y.C. Jing. 2015. Transcriptomes reveal the genetic mechanisms underlying ionic regulatory adaptations to salt in the crab-eating frog. *Scientific Reports*. 5. 17551. 10.1038/srep17551.
- Shimada T., M.Matsui, P.Yambun, A.Sudin. 2011. A survey of morphological variation in adult *Meristogenys amoropalamus* (Amphibia, Anura, Ranidae), with a description of a new cryptic species. *Zootaxa*, 33-56
- Silva D. D. M. E., A. D.D. Cruz, R. P. Bastos, R. L. Reis, M. P. D. C. Telles, J. A. F. F. D. Filho. 2007. Population structure of *Eupemphix nattereri* (Amphibia, Anura, Leiuperidae) from Central Brazil. *Genetics and Molecular Biology*, 30 (4): 1161-1168.
- Sneath, P.H. and Sokal, R.R. 1973. *Numerical Taxonomy: The Principles and Practice of Numerical Classification*. 1st Edition, W. H. Freeman, San Francisco.



- Stuart, S. N., J. S. Chanson, N. A. Cox, B. E. Young, A. S. L. Rodrigues, D. L. Fischman and R. W. Waller. 2005. Status and trends of amphibian declines and extinctions worldwide. *Science*, 306: 1783-1786.
- Tamura, K., J. Dudley, M. Nei, S. Kumar. 2007. *MEGA4 Molecular Evolutionary Genetics Analysis* (MEGA) software version 4.0.
- Telles, M. P. D. C., R. R. P. Bastos, T. N. Soares, L. V. Resende and J.A.F. Dinizfilho, 2006. RAPD Variation and Population Genetic Structure of *Physalaemus cuvieri* (Anura: Leptodactylidae) in Central Brazil. *Genetica* 128:323-332.
- Vignal, A., D. Milan, M. S. Cristobal and A. Eggen. 2002. Review on SNP and other types Of Molecular Markers and Their Use in Animal Genetics. *Genet. Sel. Evol.* 34 (275-305)
- Vitt L.J. and J.P. Caldwell. 2009. *Herpetology: An Introductory Biology of Amphibians and Reptiles Third Edition*. USA: Elsevier Inc.
- Wakita, K. & Metcalfe, I., 2005. Ocean Plate Stratigraphy in East and Southeast Asia. *Journal of Asian Earth Sciences*, 24, 679–702.
- Welsh, W and M. McClelland, 1990. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Research*, Vol. 18, No. 24 7213
- Wien, J. J. 2000. *Phylogenetic, Analysis of Morphological Data*. Smithsonian Institution. USA.
- Williams, J. G. K., A. R. Kubelik, K. J. Livak, J. A. Rafalski and S. V. Tingey. 1990. DNA Polymorphisms Amplified by Arbitrary Primers are Useful as Genetic Markers. *Nucleic Acids Res*, 18(4): 6531-6535.
- Yadav, A. and B.R. Yadav, 2007. Genetic diversity among six breeds of Indian goat using RAPD markers. *Biotechnology*, 6: 57-60. DOI: 10.3923/biotech.2007.57.60
- Yeh, F.C., R.C. Yang, T. Boyle. 1999. POPGENE, The User-Friendly Shareware For Population Genetic Analysis. *Molecular Biology and Biotechnology* Centre, University of Alberta, Canada.
- Yodthong S.B. L. Stuart, A. Aowphol. 2019. Species delimitation of crab-eating frogs (*Fejervarya cancrivora* complex) clarifies taxonomy and geographic distributions in mainland Southeast Asia. *ZooKeys*, 883: 119-153.
- Zhang, D.R, M.Y. Chen, R.W. Murphy, J. Che, J.F. Pang, J.S. Hu, Y.P. Zhang. 2010. Genealogy and palaeodrainage basins in Yunnan Province: phylogeography of the Yunnan spiny frog, *Nanorana yunnanensis*

(Dicroglossidae). *Molecular Ecology*, 19(16), 3406–3420. doi:10.1111/j.1365-294x.2010.0474(4):

