

DAFTAR PUSTAKA

- Afiyanti, & Rachmawati. (2014). *Metodologi Penelitian Kualitatif dalam Riset Keperawatan* (1st ed.). Jakarta: Rajawali Pers.
- Agency for Healthcare, & Research and Quality. (2008). 2007 National healthcare disparities report. *U.S. Department of Health and Human Services, Agency for Healthcare Research and Quality*, (Book, Whole), 244. <https://doi.org/10.1037/e593252009-001>
- Aliasgharpour, M., Abbasi, Z., Pedram Razi, S., & Kazemnezhad, A. (2016). The effect of stretching exercises on severity of restless legs syndrome in patients on hemodialysis. *Asian Journal of Sports Medicine*, 7(2). <https://doi.org/10.5812/asjsm.31001>
- Allen, R. P. (2007). Controversies and Challenges in Defining the Etiology and Pathophysiology of Restless Legs Syndrome. *American Journal of Medicine*, 120(1 SUPPL. 1), 13–21. <https://doi.org/10.1016/j.amjmed.2006.11.003>
- Allen, R. P., Picchietti, D., Hening, W. A., Trenkwalder, C., Walters, A. S., & Montplaisi, J. (2003). Restless legs syndrome: diagnostic criteria, special considerations, and epidemiology. A report from the restless legs syndrome diagnosis and epidemiology workshop at the National Institutes of Health. *Sleep Med*, 4(2), 101–119. [https://doi.org/S1389945703000108 \[pii\]](https://doi.org/S1389945703000108)
- American College of Sports Medicine. (1998). Position stand on exercise and physical activity for older adults. *Medicine and Science in Sports and Exercise*, 30(6), 992–1008. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9624662
- American College of Sports Medicine. (2014). ACSM's Guidelines for Exercise Testing and Prescription. In *Igarss 2014*. <https://doi.org/10.1007/s13398-014-0173-7.2>
- Anand, S., Johansen, K. L., Grimes, B., Kaysen, G. A., Dalrymple, L. S., Kutner, N. G., & Chertow, G. M. (2013). Physical activity and self-reported symptoms of insomnia, restless legs syndrome, and depression: The comprehensive dialysis study. *Hemodialysis International*, 17(1), 50–58. <https://doi.org/10.1111/j.1542-4758.2012.00726.x>
- Araujo, S. M. H. A., Bruin, V. M. S. de, Nepomuceno, L. A., Maximo, M. L., Daher, E. de F., Correia Ferrer, D. P., & Bruin, P. F. C. de. (2010). Restless legs syndrome in end-stage renal disease: Clinical characteristics and associated comorbidities. *Sleep Medicine*, 11(8), 785–790. <https://doi.org/10.1016/j.sleep.2010.02.011>

- Atimiati. (2012). Tingkat kualitas hidup pasien gagal ginjal kronik terapi hemodialisis. *Kesehatan Masyarakat*.
- Batool-Anwar, S., Malhotra, A., Forman, J., Winkelman, J., Li, Y., & Gao, X. (2011). Restless legs syndrome and hypertension in middle-aged women. *Hypertension*, 58(1), 791–796. <https://doi.org/10.1161/HYPERTENSIONAHA.111.174037>
- Baumgaertel, M. W., Kraemer, M., & Berlit, P. (2014). Neurologic complications of acute and chronic renal disease. *Handbook of Clinical Neurology*, 119, 383–393. <https://doi.org/10.1016/B978-0-7020-4086-3.00024-2>
- Bennett, P. (2015). Resource packs to facilitate exercise on dialysis. *Australian Nursing & Midwifery Journal*, 23(6), 39. Retrieved from <http://ezproxy2.library.drexel.edu/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=c8h&AN=111145541&site=ehost-live>
- Bennett, P., Corradini, A., Ockerby, C., & Cossich, T. (2012). Exercise during hemodialysis the intradialytic Zumba Gold. *Nephrol News Issues*, 26(9), 31–32.
- Bennett, P. N., Breugelmans, L., Barnard, R., Agius, M., Chan, D., Fraser, D., ... Potter, L. (2010). Sustaining a hemodialysis exercise program: A review. *Seminars in Dialysis*, Vol. 23, pp. 62–73. <https://doi.org/10.1111/j.1525-139X.2009.00652.x>
- Birdee, G. S., Rothman, R. L., Sohl, S. J., Wertebaker, D., Wheeler, A., Bossart, C., ... Ikizler, T. A. (2015). Feasibility and Safety of Intradialysis Yoga and Education in Maintenance Hemodialysis Patients. *Journal of Renal Nutrition*, 25(5), 445–453. <https://doi.org/10.1053/j.jrn.2015.02.004>
- Black, J. M., & Hawks, J. H. (2014). *Medical surgical nursing: Clinical management for positive outcomes*. St. Louis: Elsevier.
- Brouns, R., & De Deyn, P. P. (2004). Neurological complications in renal failure: a review. *Clin Neurol Neurosurg*, 107(1), 1–16. [https://doi.org/S0303-8467\(04\)00113-1 \[pii\]\r10.1016/j.clineuro.2004.07.012](https://doi.org/S0303-8467(04)00113-1 [pii]\r10.1016/j.clineuro.2004.07.012)
- Centers for Disease Control and Prevention. (2014). National Chronic Kidney Disease Fact Sheet. *CKD Surveillance Project*, 1–4.
- Centers for Disease Control and Prevention. (2017). National Chronic Kidney Disease Fact Sheet 2017. *US Department of Health and Human Services, Center for Disease Control and Prevention*, 1–4.
- Cheema, B. S. B., & Fiatarone Singh, M. A. (2005). Exercise training in patients receiving maintenance hemodialysis: A systematic review of clinical trials.

- American Journal of Nephrology*, Vol. 25, pp. 352–364.
<https://doi.org/10.1159/000087184>
- Cheema, B. S. B., O'Sullivan, A. J., Chan, M., Patwardhan, A., Kelly, J., Gillin, A., & Fiatarone Singh, M. A. (2006). Progressive resistance training during hemodialysis: Rationale and method of a randomized-controlled trial. *Hemodialysis International*, 10(3), 303–310. <https://doi.org/10.1111/j.1542-4758.2006.00112.x>
- Chung, Y.-C., Yeh, M.-L., & Liu, Y.-M. (2017). Effects of intradialytic exercise on the physical function, depression and quality of life for haemodialysis patients: a systematic review and meta-analysis of randomised controlled trials. *Journal of Clinical Nursing*, 26(13–14), 1801–1813. <https://doi.org/10.1111/jocn.13514>
- Comella, C. (2014). Treatment of Restless Legs Syndrome. *Neurotherapeutics*, 11(1), 177–187. <https://doi.org/10.1007/s13311-013-0247-9>
- Dahlan, S. (2014). *Besar Sampel dalam Penelitian Kedokteran dan Kesehatan* (6th ed.). Jakarta: Salemba Medika.
- Dharma, K. K. (2011). *Metodologi Penelitian Keperawatan : Panduan Melaksanakan dan Menerapkan Hasil Penelitian*. Jakarta: Trans Info Media.
- Earley, C. J., Allen, R. P., & Hening, W. (2011). Restless legs syndrome and periodic leg movements in sleep. Retrieved from Handb Clin Neurol website: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed10&NEWS=N&AN=21056236>
- Fuhrmann, I., & Krause, R. (2004). Principles of exercising in patients with chronic kidney disease, on dialysis and for kidney transplant recipients. *Clin Nephrol*, 61 Suppl 1, S14-25. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15233243>
- Fuhs, A., Bentama, D., Antkowiak, R., Mathis, J., Trenkwalder, C., & Berger, K. (2014). Effects of short- and long-term variations in RLS severity on perceived health status - the COR-Study. *PLoS ONE*, 9(4). <https://doi.org/10.1371/journal.pone.0094821>
- Garcia-Borreguero, D., Silber, M. H., Winkelman, J. W., Högl, B., Bainbridge, J., Buchfuhrer, M., ... Allen, R. P. (2016). Guidelines for the first-line treatment of restless legs syndrome/Willis-Ekbom disease, prevention and treatment of dopaminergic augmentation: A combined task force of the IRLSSG, EURLSSG, and the RLS-foundation. *Sleep Medicine*, 21, 1–11. <https://doi.org/10.1016/j.sleep.2016.01.017>
- Giannaki, C. D., Hadjigeorgiou, G. M., Karatzafiri, C., Maridaki, M. D., Koutedakis, Y., Founta, P., ... Sakkas, G. K. (2013). A single-blind

randomized controlled trial to evaluate the effect of 6 months of progressive aerobic exercise training in patients with uraemic restless legs syndrome. *Nephrology, Dialysis, Transplantation : Official Publication of the European Dialysis and Transplant Association - European Renal Association*, 28(11), 2834–2840. <https://doi.org/10.1093/ndt/gft288>

Giannaki, C. D., Sakkas, G. K., Hadjigeorgiou, G. M., Karatzafiri, C., Patramani, G., Lavdas, E., ... Stefanidis, I. (2010). Non-pharmacological management of periodic limb movements during hemodialysis session in patients with uremic restless legs syndrome. *ASAIO Journal*, 56(6), 538–542. <https://doi.org/10.1097/MAT.0b013e3181f1cc04>

Giannaki, C. D., Sakkas, G. K., Karatzafiri, C., Hadjigeorgiou, G. M., Lavdas, E., Kyriakides, T., ... Stefanidis, I. (2013). Effect of exercise training and dopamine agonists in patients with uremic restless legs syndrome: A six-month randomized, partially double-blind, placebo-controlled comparative study. *BMC Nephrology*, 14(1). <https://doi.org/10.1186/1471-2369-14-194>

Giannaki, C. D., Sakkas, G. K., Karatzafiri, C., Hadjigeorgiou, G. M., Lavdas, E., Liakopoulos, V., ... Stefanidis, I. (2011). Evidence of increased muscle atrophy and impaired quality of life parameters in patients with Uremic restless legs syndrome. *PLoS ONE*, 6(10). <https://doi.org/10.1371/journal.pone.0025180>

Graef, P., Michaelsen, S. M., & Dadalt, M. L. R. (2016). *Effects of functional and analytical strength training on upper-extremity activity after stroke: a randomized controlled trial*. 20(6), 543–552.

Greenwood, S. A., Naish, P., Clark, R., O'Connor, E., Pursey, V. A., Macdougall, I. C., ... Koufaki, P. (2014). Intra-dialytic exercise training: A pragmatic approach. *Journal of Renal Care*, 40(3), 219–226. <https://doi.org/10.1111/jorc.12080>

Hidayat, A. A. (2007). *Metode Penelitian Keperawatan dan Analisa Data*. Jakarta: Salemba Medika.

Ikizler, T. A., Pupim, L. B., Brouillette, J. R., Levenhagen, D. K., Farmer, K., Hakim, R. M., & Flakoll, P. J. (2002). Hemodialysis stimulates muscle and whole body protein loss and alters substrate oxidation. *American Journal of Physiology. Endocrinology and Metabolism*, 282(1), E107-16. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11739090>

IRLSSG. (2003). Validation of the International Restless Legs Syndrome Study Group rating scale for restless legs syndrome. *Sleep Medicine*, 4(2), 121–132. [https://doi.org/10.1016/S1389-9457\(02\)00258-7](https://doi.org/10.1016/S1389-9457(02)00258-7)

IRR (Indonesian Renal Registry). (2015). *Program Indonesian Renal Registry (IRR)*. 1–45.

IRR (Indonesian Renal Registry). (2017). *9 th Report Of Indonesian Renal Registry 2016*.

IRR (Indonesian Renal Registry). (2018). *10 th Report Of Indonesian Renal Registry 2017 10 th Report Of Indonesian Renal Registry 2017*.

Jaber, B. L., Schiller, B., Burkart, J. M., Daoui, R., Kraus, M. A., Lee, Y., ... Finkelstein, F. O. (2011). Impact of short daily hemodialysis on restless legs symptoms and sleep disturbances. *Clinical Journal of the American Society of Nephrology*, 6(5), 1049–1056. <https://doi.org/10.2215/CJN.10451110>

Johns, M. W. (1991). *A New Method for Measuring Daytime Sleepiness : The Epworth Sleepiness Scale*. 14(6), 540–545.

Johnson, R., Feehally, J., & Floege, J. (2015). Comprehensive Clinical Nephrology, 5th Edition. In *Saunders, an imprint of Elsevier Inc*.

Kallenbach, J.Z., Gutch, C.F., Martha, S.H., & Corca, A. L. (2005). *Review of Hemodialysis For nurses and dialysis personel* (7th ed.). St. Louis: Elsevier Mosby.

KDIGO. (2013). KDIGO 2012: Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Journal of the International Society of Nephrology*, 3(1), Supplement. <https://doi.org/10.1038/kisup.2012.76>

Kemenkes, R. (2017). *InfoDATIN*.

Kidney Disease Improving Global Outcomes. (2012). KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney International Supplements*, 2(4), 279–335. <https://doi.org/10.1038/kisup.2012.40>

Knap, B., Buturovic-Ponikvar, J., Ponikvar, R., & Bren, A. F. (2005). Regular Exercise as a Part of Treatment for Patients With End-stage Renal Disease. *Therapeutic Apheresis and Dialysis*, 9(3), 211–213. <https://doi.org/10.1111/j.1774-9987.2005.00256.x>

Kohnen, R., Martinez-martin, P., Benes, H., & Trenkwalder, C. (2016). Validation of the Kohnen Restless Legs Syndrome – Quality of Life instrument. *Sleep Medicine*, 24, 10–17. <https://doi.org/10.1016/j.sleep.2016.04.019>

Kontos, P. C., Miller, K.-L., Brooks, D., Jassal, S. V., Spanjevic, L., Devins, G. M., ... Naglie, G. (2007). Factors influencing exercise participation by older adults requiring chronic hemodialysis: a qualitative study. *International Urology and Nephrology*, 39(4), 1303–1311. <https://doi.org/10.1007/s11255-007-9265-z>

- La Manna, G., Pizza, F., Persici, E., Baraldi, O., Comai, G., Cappuccilli, M. L., ... Stefoni, S. (2011). Restless legs syndrome enhances cardiovascular risk and mortality in patients with end-stage kidney disease undergoing long-term haemodialysis treatment. *Nephrology Dialysis Transplantation*, 26(6), 1976–1983. <https://doi.org/10.1093/ndt/gfq681>
- Lekha. (2016). *Effectiveness of Intradialytic Stretching Exercise on Prevention and Reduction of Muscle Cramps among Patients Undergoing Hemodialysis at PSG Hospitals* (The Tamil Nadu Dr. M G R Medical University). Retrieved from <https://scholar.google.co.id>
- LeMone, & Burke. (2008). *Medical surgical nursing : Critical thinking in client care* (4th ed.). Pearson Prentice Hall : New Jersey: Lumongga L. N.
- Levey, A. S., Coresh, J., Balk, E., Kausz, A. T., Levin, A., Steffes, M. W., ... Willis, K. (2003). National Kidney Foundation Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification, and Stratification. *Annals of Internal Medicine*, Vol. 139, p. 137–147+I36. <https://doi.org/10.7326/0003-4819-139-2-200307150-00013>
- Li, Y., Walters, A. S., Chiuve, S. E., Rimm, E. B., Winkelman, J. W., & Gao, X. (2012). Prospective study of restless legs syndrome and coronary heart disease among women. *Circulation*, 126(14), 1689–1694. <https://doi.org/10.1161/CIRCULATIONAHA.112.112698>
- Lima, M. C. De, Cicotoste, C. D. L., Cardoso, S., Alberto, L., Junior, F., Monteiro, M. B., & Dias, A. S. (2013). *Effect of Exercise Performed during Hemodialysis : Strength versus Aerobic*. (December 2012), 1–8. <https://doi.org/10.3109/0886022X.2013.780977>
- Mahrova, A., & Svagrova, K. (2013). *Exercise Therapy – Additional Tool for Managing Physical and Psychological Problems on Hemodialysis*. 753–821. <https://doi.org/10.5772/53058>
- Martins, J. V., Santos, T. M., Coutinho, E. S., & Laks, J. (2015). *Comparison of strength training , aerobic training , and additional physical therapy as supplementary treatments for Parkinson ' s disease : pilot study*. 183–191.
- McMurray, A.; Blazey, L.; Fetherston, C. (2008). The effect of intradialytic foot pedal exercise on blood pressure phosphate removal efficiency and health related quality of life in haemodialysis patients. *Ren Soc Aust J*, 4(July), 38–44.
- Miranda, M., Kagi, M., Fabres, L., Aguilera, L., Alvo, M., Elgueta, L., ... Venegas, P. (2004). Pramipexole for the treatment of uremic restless legs in patients undergoing hemodialysis. *Neurology*, 62(5), 831–832. <https://doi.org/10.1212/01.WNL.0000113752.14744.15>
- Molnar, M. Z., Novak, M., Ambrus, C., Szeifert, L., Kovacs, A., Pap, J., ...

- Mucsi, I. (2005). Restless Legs Syndrome in patients after renal transplantation. *Am.J.Kidney Dis.*, 45(2), 388–396.
- Mortazavi, M., Vahdatpour, B., Ghasempour, A., Taheri, D., Shahidi, S., Moeinzadeh, F., ... Dolatkhah, S. (2013). Aerobic exercise improves signs of restless leg syndrome in end stage renal disease patients suffering chronic hemodialysis. *Thescientificworldjournal*, 2013, 628142. <https://doi.org/https://dx.doi.org/10.1155/2013/628142>
- Motedayen, Z., Nehrir, B., Tayebi, A., Ebadi, A., & Einollahi, B. (2014). The effect of the physical and mental exercises during hemodialysis on fatigue: A controlled clinical trial. *Nephro-Urology Monthly*, 6(4). <https://doi.org/10.5812/numonthly.14686>
- Mubarak, W. I., & Chayatin, N. (2008). *Buku ajar kebutuhan dasar manusia teori & aplikasi dalam praktik*. Jakarta: EGC.
- Mucsi, I., Molnar, M. Z., Ambrus, C., Szeifert, L., Kovacz, A. Z., Zoller, R., ... Novak, M. (2005). Restless legs syndrome, insomnia and quality of life in patients on maintenance dialysis. *Nephrology Dialysis Transplantation*, 20(3), 571–577. <https://doi.org/10.1093/ndt/gfh654>
- National Kidney Foundation. (2015). Clinical Practice Guideline for Hemodialysis Adequacy: 2015 Update. *American Journal of Kidney Diseases*, 66(5). <https://doi.org/10.1053/j.ajkd.2015.07.015>
- Nonoyama, M. L., Brooks, D., Ponikvar, A., Jassal, S. V., Kontos, P., Devins, G. M., ... Naglie, G. (2010). Exercise program to enhance physical performance and quality of life of older hemodialysis patients: A feasibility study. *International Urology and Nephrology*, 42(4), 1125–1130. <https://doi.org/10.1007/s11255-010-9718-7>
- Notoatmodjo, S. (2010). *Metodologi Penelitian Kesehatan*. Jakarta: Rineka Cipta.
- Notoatmodjo, S. (2012). *Promosi Kesehatan dan Perilaku Kesehatan*. Jakarta: Rineka Cipta.
- Novak, M., Winkelman, J. W., & Unruh, M. (2015). Restless Legs Syndrome in Patients With Chronic Kidney Disease. *Seminars in Nephrology*, Vol. 35, pp. 347–358. <https://doi.org/10.1016/j.semephrol.2015.06.006>
- Nur, A., Erika, K. A., & Sinrang, A. W. (2018). *The Effect of Intradialysis Streching Exercise on the Scale of Restless Legs Syndrome*. 3, 16–24.
- Parker, K. (2016). *Exercise is Medicine Intradialytic Exercise is Medicine for Hemodialysis Patients*. 1(10), 269–275.
- Parsons, T. L., Toffelmire, E. B., & King-VanVlack, C. E. (2006). Exercise

Training During Hemodialysis Improves Dialysis Efficacy and Physical Performance. *Archives of Physical Medicine and Rehabilitation*, 87(5), 680–687. <https://doi.org/10.1016/j.apmr.2005.12.044>

Pellecchia, M. T., Vitale, C., Sabatini, M., Longo, K., Amboni, M., Bonavita, V., & Barone, P. (2004). Ropinirole as a treatment of restless legs syndrome in patients on chronic hemodialysis: An open randomized crossover trial versus levodopa sustained release. *Clinical Neuropharmacology*, 27(4), 178–181. <https://doi.org/10.1097/01.wnf.0000135480.78529.06>

Potter, P. A., & Perry, A. G. (2006). *Buku ajar fundamental keperawatan* (4 volume 2). Jakarta: EGC.

Rafiudin. (2004). *Insomnia dan Gangguan Tidur*. Jakarta: PT.Elex Media Komputindo.

Raj, D. S. C., Sun, Y., & Tzamaloukas, A. H. (2008). Hypercatabolism in dialysis patients. *Current Opinion in Nephrology and Hypertension*, 17(6), 589–594. <https://doi.org/10.1097/MNH.0b013e32830d5bfa>

Rapley. (2004). Quality of Life Research: A Critical Introduction. *JSTOR Springer*, 13, 1021–2024.

Ratnawati. (2014). *Efektivitas Dialiser Proses Ulang (DPU) pada Penderita Gagal Ginjal Kronik (Hemodialisa)*. 2(April), 2–6.

Sacher, R., & Pherson. (2004). *Tinjauan Klinis Hasil Pemeriksaan Laboratorium* (11th ed.; Pendit & Wulandari, Eds.). Jakarta: EGC.

Sakkas, G. K., Hadjigeorgiou, G. M., Karatzafiri, C., Maridaki, M. D., Giannaki, C. D., Mertens, P. R., ... Stefanidis, I. (2008). Intradialytic Aerobic Exercise Training Ameliorates Symptoms of Restless Legs Syndrome and Improves Functional Capacity in Patients on Hemodialysis. *ASAIO Journal*, 54(2), 185–190. <https://doi.org/10.1097/MAT.0b013e3181641b07>

Saxena, S., & Orley, J. (1997). *Quality of life assessment : the World Health Organization perspective*. 12(Suppl 3).

Segura, E. (2010). *evidence based nephrology Exercise in haemodialysis patients : a systematic review evidence based*.

Seidel, U. K., Gronewold, J., Volsek, M., Todica, O., Kribben, A., Bruck, H., & Hermann, D. M. (2014). *Physical , Cognitive and Emotional Factors Contributing to Quality of Life , Functional Health and Participation in Community Dwelling in Chronic Kidney Disease.* 9(3). <https://doi.org/10.1371/journal.pone.0091176>

Seifert, T., Brassard, P., Wissenberg, M., Rasmussen, P., Nordby, P., Stallknecht,

- B., ... Hb, N. (2010). *Endurance training enhances BDNF release from the human brain*. 372–377. <https://doi.org/10.1152/ajpregu.00525.2009>.
- Shahgholian, N., Jazi, S. K., Karimian, J., & Valiani, M. (2016). *The effects of two methods of reflexology and stretching exercises on the severity of restless leg syndrome among hemodialysis patients*. 219–224. <https://doi.org/10.4103/1735-9066.180381>
- Sherwood, L. (2014). *Fisiologi manusia dari sel ke sistem* (8th ed.). Jakarta: EGC.
- Silva, S. F. Da, Pereira, A. A., Silva, W. A. H. Da, Simões, R., & Barros Neto, J. D. R. (2013). Physical therapy during hemodialyse in patients with chronic kidney disease. *Jornal Brasileiro de Nefrologia: 'orgão Oficial de Sociedades Brasileira E Latino-Americana de Nefrologia*, 35(3), 170–176. <https://doi.org/10.5935/0101-2800.20130028>
- Smelzer, S. C., & Bare, B. G. (2002). *Buku Ajar Keperawatan Medikal Bedah Brunner and Suddarth* (8th ed.). Jakarta: Penerbit Buku Kedokteran EGC.
- Smyth, C. (2012). The Epworth Sleepiness Scale (ESS). *Medsurg Nursing: Official Journal of the Academy of Medical-Surgical Nurses*, 18(2), 134, inside back cover. <https://doi.org/10.1093/occmed/kqm009>
- Stenvinkel, P., Carrero, J. J., von Walden, F., Ikizler, T. A., & Nader, G. A. (2015). Muscle wasting in end-stage renal disease promulgates premature death: established, emerging and potential novel treatment strategies. *Nephrology, Dialysis, Transplantation: Official Publication of the European Dialysis and Transplant Association - European Renal Association*. [https://doi.org/gfv122 \[pii\]](https://doi.org/gfv122 [pii])
- Stenvinkel, P., Carrero, J. J., Von Walden, F., Ikizler, T. A., & Nader, G. A. (2016). Muscle wasting in end-stage renal disease promulgates premature death: Established, emerging and potential novel treatment strategies. *Nephrology Dialysis Transplantation*, Vol. 31, pp. 1070–1077. <https://doi.org/10.1093/ndt/gfv122>
- Sugiyono. (2017). *Metode Penelitian Kuantitatif, Kualitatif*. Bnadung: Alfabeta.
- Suyono, S. (2001). *Buku Ajar Ilmu Penyakit Dalam Jilid II* (3rd ed.). Jakarta: Balai Penerbit FKUI.
- Takhreem, M. (2008). The effectiveness of intradialytic exercise prescription on quality of life in patients with chronic kidney disease. *Medscape Journal of Medicine*, 10(10), 226.
- Thomas, N. (2003). *Renal Nursing* (2nd ed.). London: Bailiere Tindall.
- Tsekoura, D., & Manolis, A. J. (2014). *The association of Restless Legs Syndrome*

with hypertension and cardiovascular disease. 654–659.
<https://doi.org/10.12659/MSM.890252>

Unruh, M. L., Levey, A. S., D'Ambrosio, C., Fink, N. E., Powe, N. R., & Meyer, K. B. (2004). Restless Legs Symptoms among Incident Dialysis Patients: Association with Lower Quality of Life and Shorter Survival. *American Journal of Kidney Diseases*, 43(5), 900–909.
<https://doi.org/10.1053/j.ajkd.2004.01.013>

Wawan, & Dewi. (2010). *Teori dan Pengukuran Pengetahuan, Sikap dan Perilaku Manusia*. Jakarta: Nuha Medika.

Widianti, A. T., Hermayanti, Y., & Kurniawan, T. (2017). *Pengaruh Latihan Kekuatan terhadap Restless Legs Syndrome Pasien Hemodialisis Effect of Strength Training on Restless Legs Syndrome: Hemodialysis patients*. 5(April), 47–56.

Winkelmann, J. W., Chertow, G. M., & Lazarus, J. M. (1996). Restless legs syndrome in end-stage renal disease. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 28(3), 372–378.
<https://doi.org/10.1016/j.sleep.2004.01.014>

Winter, A. C., Berger, K., Glynn, R. J., Buring, J. E., Gaziano, J. M., Schürks, M., & Kurth, T. (2013). Vascular Risk Factors , Cardiovascular Disease , and Restless Legs Syndrome in Men. *AJM*, 126(3), 228–235.e2.
<https://doi.org/10.1016/j.amjmed.2012.06.039>