

DAFTAR PUSTAKA

1. Irianti T, Mada UG, Ugm S, et al. Antioksidant. Yogyakarta: 2017;
2. Nurkhasanah MA, Si A, Mochammad S, et al. Antioksidan dan Stres Oksidatif. Yogyakarta: UAD PRESS, 2023;
3. Dungir SG, Katja DG, Kamu VS. Aktivitas Antioksidan Ekstrak Fenolik dari Kulit Buah Manggis (*Garcinia mangostana* L.). J MIPA 2012;1(1):11.
4. Jahanbakhshi A, Abbaspour-Gilandeh Y, Heidarbeigi K, Momeny M. Detection of fraud in ginger powder using an automatic sorting system based on image processing technique and deep learning. Comput Biol Med 2021;136(August):104764. Available from: <https://doi.org/10.1016/j.combiomed.2021.104764>
5. Hu W, Yu A, Wang S, et al. Extraction, Purification, Structural Characteristics, Biological Activities, and Applications of the Polysaccharides from *Zingiber officinale* Roscoe. (Ginger): A Review. Molecules 2023;28(9).
6. Zhang H, Ren Y, Vomiting C, Li X, Wang S, Categorized I. Antioxidant activity and total phenolic content of three varieties of Ginger (*Zingiber officinale*) in decoction and infusion extraction method Antioxidant activity and total phenolic content of three varieties of Ginger (*Zingiber officinale*) in decoct. :4–10.
7. Hidayat R, Patricia Wulandari. Methods of Extraction: Maceration, Percolation and Decoction. Eureka Herba Indones 2021;2(1):73–79.
8. Cha J, Kim CT, Cho YJ. Optimizing extraction conditions for functional compounds from ginger (*Zingiber officinale* Roscoe) using response surface methodology. Food Sci Biotechnol [homepage on the Internet] 2020;29(3):379–385. Available from: <https://doi.org/10.1007/s10068-019-00667-9>
9. Shahabi Mohammadabadi S, Goli M, Naji Tabasi S. Optimization of Bioactive Compound Extraction from Eggplant Peel by Response Surface Methodology: Ultrasound-Assisted Solvent Qualitative and Quantitative Effect. Foods 2022;11(20).
10. Yusof N, Munaim MSA, Veloo Kutty R. Optimization of total phenolic compounds extracted from propolis by ultrasound- assisted extraction. Chem Eng Commun [homepage on the Internet] 2021;208(4):564–572. Available from: <https://doi.org/10.1080/00986445.2020.1761799>
11. Beg S, Swain S, Rahman M, Hasnain MS, Imam SS. Application of Design of Experiments (DoE) in Pharmaceutical Product and Process Optimization [Homepage on the Internet]. Elsevier Inc., 2019; Available from: <http://dx.doi.org/10.1016/B978-0-12-815799-2.00003-4>
12. Spence C. Ginger: The pungent spice. Int J Gastron Food Sci [homepage on the Internet] 2023;33(August):100793. Available from: <https://doi.org/10.1016/j.ijgfs.2023.100793>
13. Nurhadi B, Suriati, Tensiska, Saputra RA, Sukri N. The role of encapsulant materials on the stability of bioactive compounds of red ginger (*Zingiber officinale* Roscoe. var. Rubrum) extract

- powder during storage. *Food Chem* 2020;333(June).
14. Fitri Yanti C, Nilawati Usman A, Ahmad M, Ilhammudin, Ariyandi A, Budiaman. Analysis of Ginger Varieties (*Zingiber Officinale*) in Indonesia, Nutritional Content and Potential for Health. *BIO Web Conf* 2024;96.
 15. Windarsih G, Utami DW, Yuriyah S. Morphological characteristics of zingiberaceae in serang district, banten, indonesia. *Biodiversitas* 2021;22(12):5507–5529.
 16. Zhang S, Kou X, Zhao H, Mak KK, Balijepalli MK, Pichika MR. *Zingiber officinale* var. *rubrum*: Red Ginger's Medicinal Uses. *Molecules* 2022;27(3).
 17. Rini Daud Supu, Ajeng Diantini JLk. Red Ginger (*Zingiber officinale* var. *rubrum*): Its Chemical Constituen, Pharmacological Activities and Safety. 2018;7(1):1–25.
 18. Yefrida Y, Refilda R, Hamidah N, Rosman W. Penentuan Kandungan Antioksidan Total pada Infusa Selada Hijau (*Lactuca sativa* L.) Hidroponik dan Konvensional secara Spektrofotometri dengan Modified Phenantroline Method (MPM). *J Ris Kim* 2022;13(1):122–129.
 19. Yefrida, Refilda, Detri Susanti S. Antioxidants And Total Phenolics From Plants Of The Malvaceae Family. 2023;21(1):1–07. Available from: <https://jurnal.kimia.fmipa.unmul.ac.id/index.php/JKM>
 20. Mendonça J da S, Guimarães R de CA, Zorgetto-Pinheiro VA, et al. Natural Antioxidant Evaluation: A Review of Detection Methods. *Molecules* 2022;27(11):1–37.
 21. Yıldız E, Çabuk H. Determination of the Synthetic Antioxidants Butylated Hydroxyanisole (BHA) and Butylated Hydroxytoluene (BHT) by Matrix Acidity-Induced Switchable Hydrophilicity Solvent-Based Homogeneous Liquid-Liquid Microextraction (MAL-SHS-HLLME) and High-Performance . *Anal Lett* [homepage on the Internet] 2022;55(3):480–494. Available from: <https://doi.org/10.1080/00032719.2021.1941072>
 22. Gulcin İ, Alwasel SH. DPPH Radical Scavenging Assay. *Processes* 2023;11(8).
 23. Menezes BB de, Frescura LM, Duarte R, Villetti MA, Rosa MB da. A critical examination of the DPPH method: Mistakes and inconsistencies in stoichiometry and IC50 determination by UV–Vis spectroscopy. *Anal Chim Acta* [homepage on the Internet] 2021;1157:338398. Available from: <https://doi.org/10.1016/j.aca.2021.338398>
 24. Dawidowicz AL, Wianowska D, Olszowy M. On practical problems in estimation of antioxidant activity of compounds by DPPH method (Problems in estimation of antioxidant activity). *Food Chem* 2012;131(3):1037–1043. Available from: <http://dx.doi.org/10.1016/j.foodchem.2011.09.067>
 25. Echegaray N, Pateiro M, Munekata PES, et al. Measurement of antioxidant capacity of meat and meat products: Methods and applications. *Molecules* 2021;26(13).
 26. Prado MAR and JM. *Natural Product Extraction*. London: The Royal Society of Chemistry, 2022;

27. Pourkhanali K, Khayati G, Mizani F, Raouf F. Isolation, identification and optimization of enhanced production of laccase from *Galactomyces geotrichum* under solid-state fermentation. *Prep Biochem Biotechnol* 2021;51(7):659–668. Available from: <https://doi.org/10.1080/10826068.2020.1848867>
28. Rahman MF, Khayat ME, EL-Mongy MA, Yakasai HM, Yasid NA, Shukor MY. Central Composite Design-based Optimization of *Staphylococcus* sp. strain Amr-15 Growth on Acrylamide as a Nitrogen Source. *J Environ Microbiol Toxicol* 2022;10(2):13–22.
29. Shokri R, Jalilzadeh Yengejeh R, Babaei AA, Derikvand E, Almasi A. UV activation of hydrogen peroxide for removal of azithromycin antibiotic from aqueous solution: determination of optimum conditions by response surface methodology. *Toxin Rev* 2020;39(3):284–291. Available from: <https://doi.org/10.1080/15569543.2018.1517803>
30. Aghababaie M, Beheshti M, Razmjou A, Bordbar AK. Enzymatic biodiesel production from crude *Eruca sativa* oil using *Candida rugosa* lipase in a solvent-free system using response surface methodology. *Biofuels* 2020;11(1):93–99. Available from: <http://dx.doi.org/10.1080/17597269.2017.1345359>
31. Li W, Zhao LC, Wang Z, Zheng YN, Liang J, Wang H. Response surface methodology to optimize enzymatic preparation of deapio-platycodin D and platycodin D from radix platycodi. *Int J Mol Sci* 2012;13(4):4089–4100.
32. Rahimi S, Mohammadi F, Yavari Z. Removal of rr198 dye by tio₂/fe₃o₄/persulfate nanoparticles under uv-led irradiation and comparison of ofat and ccd experimental design in rsm modelling. *Indian J Chem Technol* 2020;27(4):283–293.
33. Hwang SJ, Lee JH. Comparison of antioxidant activities expressed as equivalents of standard antioxidant. *Food Sci Technol* 2023;43:1–5.
34. Zargarchi S, Esatbeyoglu T. Assessing the impact of cold plasma rotational dynamics on ginger's total phenolic content, antioxidant activity, surface structure and color using response surface methodology. 2024;208(August):116682.
35. Alqahtani NK, Salih ZA, Asiri SA, et al. Optimizing physicochemical properties, antioxidant potential, and antibacterial activity of dry ginger extract using sonication treatment. *Heliyon* [homepage on the Internet] 2024;10(16):e36473. Available from: <https://doi.org/10.1016/j.heliyon.2024.e36473>
36. Rahim NFA, Muhammad N, Abdullah N, Talip BA, Poh KH. The interaction effect and optimal formulation of selected polyherbal extracts towards antioxidant activity. *Food Res* 2020;4(6):2042–2048.
37. Sumazian Y, Syahida A, Hakiman M, Maziah M. Antioxidant activities, flavonoids, ascorbic acid and phenolic contents of Malaysian vegetables. *J Med Plants Res* 2010;4(10):881–890.
38. Sanna D, Delogu G, Mulas M, Schirra M, Fadda A. Determination of Free Radical Scavenging Activity of Plant Extracts Through DPPH Assay: An EPR and UV-Vis Study. *Food Anal Methods*

2012;5(4):759–766.

39. Thi T, Truyen T, Thi N, Hue L. The Effects OF Extraction Conditions on Antioxidant Activity of Sesquiterpen Extract From *Cypers rotundus* L . 2024;24(3):14–22.
40. A AJC. Archive of SID . ir Application of Design of Experiment (DoE) for Optimization of Multiple Parameter Resource Constrain Process : Taguchi- Archive of SID . ir Archive of SID . ir Archive of SID . ir. 2023;6(4):391–400.
41. Cao Q, Yan J, Sun Z, et al. Simultaneous optimization of ultrasound-assisted extraction for total flavonoid content and antioxidant activity of the tender stem of *Triarrhena lutarioriparia* using response surface methodology. *Food Sci Biotechnol* 2021;30(1):37–45. Available from: <https://doi.org/10.1007/s10068-020-00851-2>
42. Mukherjee S, Mandal N, Dey A, Mondal B. An approach towards optimization of the extraction of polyphenolic antioxidants from ginger (*Zingiber officinale*). *J Food Sci Technol* 2014;51(11):3301–3308.
43. Antony A, Farid M. Effect of Temperatures on Polyphenols during Extraction. *Appl Sci* 2022;12(4).
44. Sobhani A, Noormohammadi N, Moradi K, Ebrahimi M, Khanahmadi M. Optimization of heat and ultrasound assisted extraction of bioactive compounds from *Echinacea purpurea* using response surface methodology. *J Appl Res Med Aromat Plants* 2022;31(April).
45. Robles-Apodaca SM, González-Vega RI, Ruíz-Cruz S, et al. Optimization of Extraction Process for Improving Polyphenols and Antioxidant Activity from Papaya Seeds (*Carica papaya* L.) Using Response Surface Methodology. *Processes* 2024;12(12).
46. Kourelatou A, Chatzimitakos T, Athanasiadis V, et al. Seeking Optimal Extraction Method for Augmenting *Hibiscus sabdariffa* Bioactive Compounds and Antioxidant Activity. *Processes* 2024;12(3):1–19.
47. Sanou A, Konaté K, Kabakdé K, et al. Modelling and optimisation of ultrasound-assisted extraction of roselle phenolic compounds using the surface response method. *Sci Rep* 2023;13(1):1–9. Available from: <https://doi.org/10.1038/s41598-023-27434-5>
48. Thanh NT. Optimization of Microwave-Assisted Extraction of Antioxidant Compounds from Roots of *Polygonum multiflorum* Thunb. at Vietnam using Response Surface Methodology. *Malaysian J Chem* 2023;25(4):21–29.
49. Kim I, Ha JH, Jeong Y. Optimization of extraction conditions for antioxidant activity of *acer tegmentosum* using response surface methodology. *Appl Sci* 2021;11(3):1–20.
50. Fernández-Escalante E, Ibañez R, San-Román MF. Opportunities of desalination concentrates for lithium recovery: Optimal separation by synergic solvents. *Sep Purif Technol* 2023;326(July).
51. Sayem ASM, Ahmed T, Mithun MUK, Rashid M, Rana MR. Optimising ultrasound-assisted extraction conditions for maximising phenolic, flavonoid content and antioxidant activity in hog

- plum peel and seed: A response surface methodology approach. *J Agric Food Res* 2024;18(July).
52. Makanjuola SA, Enujiugha VN, Omoba OS, Sanni DM. Optimization and prediction of antioxidant properties of a tea-ginger extract. *Food Sci Nutr* 2015;3(5):443–452.
53. Propolis G, Cavalaro RI, Felipe L, Fabricio DF. Ultrasound-Assisted Extraction of Antioxidants from *Melastoma malabathricum* Linn.: Modeling and Optimization Using Box–Behnken Design. 2020;(Figure 1).
54. Ghafoor K, Juhaimi F Al, Özcan MM, Uslu N, Babiker EE, Mohamed Ahmed IA. Total phenolics, total carotenoids, individual phenolics and antioxidant activity of ginger (*Zingiber officinale*) rhizome as affected by drying methods. 2020;126(December 2019).

