

DAFTAR PUSTAKA

1. Hutchinson A, Nesbitt A, Joshi A, Clubb A, Perera M. Overactive bladder syndrome: Management and treatment options. *Aust J Gen Pract.* 2020;49(9):593–8.
2. White N, Iglesia CB. Overactive Bladder. *Obstet Gynecol Clin North Am* [Internet]. 2016;43(1):59–68. Available from: <http://dx.doi.org/10.1016/j.ogc.2015.10.002>
3. Utama BI, Widayat W, Arwan B. The Incidence of Overactive Bladder in KIA Poly Patients at Pauh Health Center in Padang City Using Overactive Bladder Symptoms Score. *J Obgin Emas.* 2019;3(2):83–9.
4. Arnold J, McLeod N, Thani-Gasalam R, Rashid P. Overactive bladder syndrome: management and treatment options. *Aust Fam Physician.* 2012;41(11):878–83.
5. Ellsworth PI. Overactive bladder: etiology, diagnosis, and impact. *Medscape.* Medscape; 2021.
6. Tirtayasa PMW, Rahardjo HE. A survey on the management of overactive bladder by Indonesian urologists. *Med J Indones.* 2015;24(2):91–6.
7. Leron E, Weintraub AY, Mastrolia SA, Schwarzman P. Overactive bladder syndrome: evaluation and management. *Curr Urol.* 2017;11(3):117–25.
8. Hutchinson A, Nesbitt A, Joshi A, Clubb A, Perera M. *AJG P-09-2020-Clinical-Hutchinson-Overactive-Bladder-Syndrome-WEB.* 2020;49(9):593–8.
9. Malde S, Kelly S, Saad S, Sahai A. Case-finding tools for the diagnosis of OAB in women: A narrative review. *Neurourol Urodyn.* 2020;39(1):13–24.
10. Scarneciu I, Lupu S, Bratu OG, Teodorescu A, Maxim LS, Brinza A, et al. Overactive bladder: A review and update. *Exp Ther Med.* 2021;22(6):1–8.
11. Fang JJ, Wu MP, Yen YC, Wu JC, Chin HY. Overactive bladder syndrome is associated with detrusor overactivity and abnormal voiding pattern in nulliparous women. *J Chinese Med Assoc.* 2021;84(9):865–9.
12. Blomquist JL, Muñoz A, Carroll M, Handa VL. Association of Delivery Mode with Pelvic Floor Disorders after Childbirth. *JAMA - J Am Med*

Assoc. 2018;320(23):2438–47.

13. Fakhrizal E, Priyatini T, Santoso BI, Junizaf, Moegni F, Djusad S, et al. Prevalence and risk factors of persistent stress urinary incontinence at three months postpartum in Indonesian women. *Med J Indones*. 2016;25(3):163–70.
14. Dinas Pengendalian Penduduk, Keluarga Berencana, Kependudukan dan Pencatatan Sipil Provinsi Sumatera Barat. *Profil Perkembangan Kependudukan Provinsi Sumatera Barat Tahun 2018*. 2018.
15. Badan Pusat Statistika Padang. *Kecamatan Padang Utara dalam Angka 2018*. 2018.
16. Chuang FC, Hsiao SM, Kuo HC. The overactive bladder symptom score, international prostate symptom score-storage subscore, and urgency severity score in patients with overactive bladder and hypersensitive bladder: Which scoring system is best? *Int Neurourol J*. 2018;22(2):99–106.
17. Ismy J. *Penatalaksanaan Over Active Bladder (OAB)*. In: *Acch Surgery Update 2*. 2017.
18. Shawahna R, Hijaz H, Jallad K, Abushamma M, Sawafta M. Prevalence of overactive bladder symptoms and their impact on health-related quality of life of medical and dentistry students: a multicenter cross-sectional study. *BMC Urol*. 2021;21(1):142.
19. Eapen RS, Radomski SB. Review of the epidemiology of overactive bladder. *Res reports Urol*. 2016;8:71–6.
20. Chuang YC, Liu SP, Lee KS, Liao L, Wang J, Yoo TK, et al. Prevalence of overactive bladder in China, Taiwan and South Korea: Results from a cross-sectional, population-based study. *Low Urin Tract Symptoms*. 2019 Jan;11(1):48–55.
21. Robinson D, Cardozo L. Managing overactive bladder. *Climacteric* [Internet]. 2019;22(3):250–6. Available from: <https://doi.org/10.1080/13697137.2018.1552254>
22. Elbaset MA, Taha DE, Sharaf DE, Ashour R, El-Hefnawy AS. Obesity and Overactive Bladder: Is It a Matter of Body Weight, Fat Distribution or Function? A Preliminary Results. *Urology* [Internet]. 2020;143:91–6.

Available from: <https://doi.org/10.1016/j.urology.2020.04.115>

23. Hagovska M, Švihra J, Buková A, Horbacz A, Dračková D, Lupták J, et al. The Relationship between Overweight and Overactive Bladder Symptoms. *Obes Facts*. 2020;13(3):297–306.
24. Tähtinen RM, Cartwright R, Tsui JF, Aaltonen RL, Aoki Y, Cárdenas JL, et al. Long-term Impact of Mode of Delivery on Stress Urinary Incontinence and Urgency Urinary Incontinence: A Systematic Review and Meta-analysis. *Eur Urol*. 2016;70(1):148–58.
25. Handa VL, Pierce CB, Munoz A, Blomquist JL. Longitudinal changes in overactive bladder and stress incontinence among parous women. *Neurourol Urodyn*. 2015;34(4):356–61.
26. Gunawan IPGF, Suarjana IN, Wulandari KF. An overview of overactive bladder. *Bali Med J*. 2021;10(1):208–10.
27. Chu FM, Dmochowski R. Pathophysiology of overactive bladder. *Am J Med*. 2006;119(3 SUPPL. 1):3–8.
28. Wallace KM, Drake MJ. Overactive bladder. *F1000Research*. 2015;4.
29. Lightner DJ, Gomelsky A, Souter L, Vasavada SP. Diagnosis and treatment of overactive bladder (non-neurogenic) in adults: AUA/SUFU guideline amendment 2019. *J Urol*. 2019;202(3):558–63.
30. Pal M, Bandyopadhyay S, Roy A. Overactive bladder symptom score – translation and linguistic validation in Bengali. *J Fam Med Prim Care*. 2022;11(1).
31. Foon R, Drake MJ. The overactive bladder. *Ther Adv Urol*. 2010 Aug;2(4):147–55.
32. Fontaine C, Papworth E, Pascoe J, Hashim H. Update on the management of overactive bladder. *Ther Adv Urol*. 2021;13:17562872211039034.
33. Rhomadhon R. Hubungan Antara Nilai Overactive Bladder Symptom Score (OABSS) Dengan Stres, Depresi, Dan Kecemasan Pada Mahasiswa/Wi Preklinik Program Studi Kedokteran (PSKed) Yang Menderita Overactive Bladder (OAB). UIN Syarif Hidayatullah Jakarta-FK; 2018.
34. Homma Y, Fujimura T. Psychometric validation of the English version of the overactive bladder symptom score. *Urology*. 2014;84(1):46–50.

35. Peterson AC, Sehgal A, Crump RT, Baverstock R, Sutherland JM, Carlson K. Evaluating the 8-item overactive bladder questionnaire (OAB-v8) using item response theory. *Neurourol Urodyn*. 2018;37(3):1095–100.
36. Basra RK, Cortes E, Khullar V, Kelleher C. A comparison study of two lower urinary tract symptoms screening tools in clinical practice: The B-SAQ and OAB-V8 questionnaires. *J Obstet Gynaecol (Lahore)*. 2012;32(7):666–71.
37. Zhu J, Hu X, Dong X, Li L. Associations between risk factors and overactive bladder: a meta-analysis. *Female Pelvic Med Reconstr Surg*. 2019;25(3):238.
38. Hutton EK, Hannah ME, Willan AR, Ross S, Allen AC, Armson BA, et al. Urinary stress incontinence and other maternal outcomes 2 years after caesarean or vaginal birth for twin pregnancy: a multicentre randomised trial. *BJOG An Int J Obstet Gynaecol*. 2018;125(13):1682–90.
39. Fang JJ, Wu MP, Yen YC, Wu JC, Chin HY. Overactive bladder syndrome is associated with detrusor overactivity and abnormal voiding pattern in nulliparous women. *J Chinese Med Assoc*. 2021;84(9):865–9.
40. Blomquist JL, Carroll M, Muñoz A, Handa VL. Pelvic floor muscle strength and the incidence of pelvic floor disorders after vaginal and cesarean delivery. *Am J Obstet Gynecol* [Internet]. 2020;222(1):62.e1-62.e8. Available from: <https://doi.org/10.1016/j.ajog.2019.08.003>
41. Nygaard IE, Wolpern A, Bardsley T, Egger MJ, Shaw JM. Early postpartum physical activity and pelvic floor support and symptoms 1 year postpartum. *Am J Obstet Gynecol*. 2021;224(2):193.e1-193.e19.
42. Scarpa KP, Herrmann V, Palma PCR, Ricetto CLZ, Morais S. Sintomas do trato urinário inferior três anos após o parto: estudo prospectivo. *Rev Bras Ginecol e Obs*. 2008;30:355–9.
43. Martínez-Agulló E, Ruiz-Cerdá J, Arlandis S, Rebollo P, Pérez M, Chaves J. Analysis of overactive bladder and urinary incontinence in working women aged between 25 and 64 years. EPICC study. *Actas Urol españolas*. 2010 Jul;34:618–24.
44. Palma T, Raimondi M, Souto S, Fozzatti M, Palma P, Ricetto C. Prospective study of prevalence of overactive bladder symptoms and child-bearing in women of reproductive age. *J Obstet Gynaecol Res*. 2013 Jun;39:0.

45. Lukacz ES, Lawrence JM, Contreras R, Nager CW, Luber KM. Parity, mode of delivery, and pelvic floor disorders. *Obstet Gynecol.* 2006 Jun;107(6):1253–60.
46. Torkestani F, Zafarghandi N, Davati A, Hadavand SH, Garshasbi M. Case-controlled study of the relationship between delivery method and incidence of post-partum urinary incontinence. *J Int Med Res.* 2009;37(1):214–9.

