

DAFTAR PUSTAKA

- Agoston, D. V., Shutes-David, A. and Peskind, E.R. (2017) ‘Biofluid biomarkers of traumatic brain injury’, *Brain Injury*, 31(9), pp. 1195–1203. Available at: <https://doi.org/10.1080/02699052.2017.1357836>.
- Al-Koofee, D.A. and Mubarak, S.M. (2019) ‘Genetic polymorphisms.’, *American journal of therapeutics*, 9(4), p. 271. Available at: <https://doi.org/10.1097/00045391-200207000-00001>.
- Amalia, L. (2021) ‘Glial Fibrillary Acidic Protein (GFAP): Neuroinflammation Biomarker in Acute Ischemic Stroke’, *Journal of Inflammation Research*, 14, pp. 7501–7506. Available at: <https://doi.org/10.2147/JIR.S342097>.
- Anderson, K.J., Scheff, S.W., Miller, K.M., Roberts, K.N., Gilmer, L.K., Yang, C. and Shaw, G. (2008) ‘The phosphorylated axonal form of the neurofilament subunit NF-H (pNF-H) as a blood biomarker of traumatic brain injury’, *Journal of Neurotrauma*, 25(9), pp. 1079–1085. Available at: <https://doi.org/10.1089/neu.2007.0488>.
- Anderson, M.A., Ao, Y. and Sofroniew, M. V. (2014) ‘Heterogeneity of reactive astrocytes’, *Neuroscience Letters*, 565, pp. 23–29. Available at: <https://doi.org/10.1016/j.neulet.2013.12.030>.
- Apriawanti, V., Widodo, T.W., Handayani, S. and Mulyani, T. (2019) ‘Hubungan antara Glasgow Coma Scale dan lama perawatan pada pasien cedera otak traumatis dengan perdarahan subdural’, *Jurnal Kesehatan Khatulistiwa*, [online].
- Bazarian, J.J., Biberthaler, P., Welch, R.D., Lewis, L.M., Barzo, P., Bogner-Flatz, V. et al. (2018) ‘Serum GFAP and UCH-L1 for prediction of absence of intracranial injuries on head CT (ALERT-TBI): a multicentre observational study’, *The Lancet Neurology*, 17(9), pp. 782–789. Available at: [https://doi.org/10.1016/S1474-4422\(18\)30231-X](https://doi.org/10.1016/S1474-4422(18)30231-X).
- Berger, R.P. (2011) *Biochemical markers of head trauma in children*, *Child Abuse and Neglect*. Elsevier Inc. Available at: <https://doi.org/10.1016/B978-1-4160-6393-3.00046-4>.
- Blumbergs, P.C. (2011) ‘Neurophatology of traumatic brain injury’, in H.R. Winn (ed.) *Youmans Neurological Surgery*. Saunders, an imprint of Elsevier, pp. 3288–3299.
- Böhmer, A.E., Osés, J.P., Schmidt, A.P., Perón, C., Lara, D.R., Souza, D.O. et al. (2011) ‘Neuron-specific enolase, S100B, and glial fibrillary acidic protein levels as outcome predictors in patients with severe traumatic brain injury’, *Neurosurgery*, 68(6), pp. 1624–1630. Available at: <https://doi.org/10.1227/NEU.0b013e318214a81f>

- Borg, K., Holm, L., Undén, J., von Euler, M., & Ingebrigtsen, T. (2012) ‘Serum levels of biochemical markers of traumatic brain injury’, *ISRN Emergency Medicine*, 2012, pp. 1–7. Available at: <https://doi.org/10.5402/2012/417313>
- Brady, S., Colman, D.R. and Brophy, P. (2014) *Subcellular Organization of the Nervous System: Organelles and Their Functions*. Third Edit, *From Molecules to Networks: An Introduction to Cellular and Molecular Neuroscience: Third Edition*. Third Edit. Elsevier Inc. Available at: <https://doi.org/10.1016/B978-0-12-397179-1.00002-6>.
- Brazinova, A., Rehorčíková, V., Taylor, M.S., Buckova, V., Majdan, M., Psota, M. and Peeters, W. (2021) ‘Epidemiology of traumatic brain injury in Europe: A living systematic review’, *Journal of Neurotrauma*, 38(10), pp. 1411–1440. Available at: <https://doi.org/10.1089/neu.2015.4126>
- Brooks, G.A. and Martin, N.A. (2015) ‘Cerebral metabolism following traumatic brain injury: New discoveries with implications for treatment’, *Frontiers in Neuroscience*, 8(FEB), pp. 1–13. Available at: <https://doi.org/10.3389/fnins.2014.00408>.
- Casmiro, M., Pierpaoli, F., Tugnoli, V., Calbucci, F., Gallassi, R. and Zappaterra, S. (2005) ‘Cerebrospinal fluid and serum neuron-specific enolase concentrations in a normal population’, *European Journal of Neurology*, 12(5), pp. 369–374. Available at: <https://doi.org/10.1111/j.1468-1331.2004.01021.x>.
- Chabok, S.Y., Moghadamnia, A.A., Norouzi-Javidan, A., Ahmadi, A., Saneei, Z. and Aarabi, S. (2012) ‘Neuron-specific enolase and S100BB as outcome predictors in severe diffuse axonal injury’, *Journal of Trauma and Acute Care Surgery*, 72(6), pp. 1654–1657. Available at: <https://doi.org/10.1097/TA.0b013e318246887e>.
- Cheng, F., Yuan, Q., Yang, J., Wang, W., Liu, H. and Wang, J. (2014) ‘The prognostic value of serum neuron-specific enolase in traumatic brain injury: Systematic review and meta-analysis’, *PLoS ONE*, 9(9), e106680. Available at: <https://doi.org/10.1371/journal.pone.0106680>.
- Czeiter, E., Amrein, K., Gravesteijn, B.Y., Lecky, F., Menon, D.K. and Maas, A.I.R. (2020) ‘Blood biomarkers on admission in acute traumatic brain injury: Relations to severity, CT findings and care path in the CENTER-TBI study’, *EBioMedicine*, 56, 102785. Available at: <https://doi.org/10.1016/j.ebiom.2020.102785>.
- Demlie, T.A., Gashaw, A., Moges, T., Abebe, B., Tsegaye, A. and Birhanu, E. (2023) ‘Incidence and predictors of mortality among traumatic brain injury patients admitted to Amhara region Comprehensive Specialized Hospitals, northwest Ethiopia, 2022’, *BMC Emergency Medicine*, 23(1), p. 9. Available at: <https://doi.org/10.1186/s12873-023-00823-9>.
- Dewan, M.C., Rattani, A., Gupta, S., Baticulon, R.E., Hung, Y.C. and Punchak, M. (2019) ‘Estimating the global incidence of traumatic brain injury’, *Journal of*

- Neurosurgery*, 130(4), pp. 1080–1097. Available at: <https://doi.org/10.3171/2017.10.JNS17352>.
- Donkin, J.J. and Vink, R. (2010) ‘Mechanisms of cerebral edema in traumatic brain injury: Therapeutic developments’, *Current Opinion in Neurology*, 23(3), pp. 293–299. Available at: <https://doi.org/10.1097/WCO.0b013e328337f451>.
- Karesioglu, E., Kocamaz, D., Sahinturk, V., Arslan, B., Ozaydin, I. and Yilmaz, M. (2022) ‘Serum pNF-H levels in the first six hours after experimental mild traumatic brain injury in rats’, *European Review for Medical and Pharmacological Sciences*, 26, pp. 6928–6934.
- Eng, D.L. and Eng, L.F. (2011) ‘Glial Fibrillary Acidic Protein : The Intermediate Filament Protein of Astrocytes’, in R. Nixon and A. Yuan (eds) *Cytoskeleton of the Nervous System*. vol 3. Newyork: Springer, pp. 455–501.
- Faried, A., Arifin, M., Arifin, M.Z., Widjaya, F.K.G., Gani, A.R. and Sutanto, R. (2017) ‘Characteristics of Moderate and Severe Traumatic Brain Injury of Motorcycle Crashes in Bandung, Indonesia’, *World Neurosurgery*, 100, pp. 195–200. Available at: <https://doi.org/10.1016/J.WNEU.2016.12.133>.
- Farooqui, A.A. (2010) *Neurochemical aspects of neurotraumatic and neurodegenerative diseases, Neurochemical Aspects of Neurotraumatic and Neurodegenerative Diseases*. Available at: <https://doi.org/10.1007/978-1-4419-6652-0>.
- Feala, J.D., Abdulhameed, M.D., Okonkwo, D.O., Vodovotz, Y., and Androulakis, I.P. (2013) ‘Systems biology approaches for discovering biomarkers for traumatic brain injury’, *Journal of Neurotrauma*, 30(13), pp. 1101–1116. Available at: <https://doi.org/10.1089/neu.2012.2631>.
- Fraser, D.D., Close, T.E., Rose, K.L., Ward, R., Mehl, M., and Young, B. (2011) ‘Severe traumatic brain injury in children elevates glial fibrillary acidic protein in cerebrospinal fluid and serum’, *Pediatric Critical Care Medicine*, 12(3), pp. 319–324. Available at: <https://doi.org/10.1097/PCC.0b013e3181e8b32d>.
- Ganeshalingham, A. and Beca, J. (2021) ‘Serum biomarkers in severe paediatric traumatic brain injury—a narrative review’, *Translational Pediatrics*. AME Publishing Company, pp. 2720–2737. Available at: <https://doi.org/10.21037/tp-20-386>.
- Ghandour, H.Z., Naim, W., Samaha, A., Saade, M., Yaghi, S., and El Sayed, M. (2022) ‘Traumatic brain injury patient characteristics and outcomes in Lebanon: A multicenter retrospective cohort study’, *Journal of Global Health Reports*, 6. Available at: <https://doi.org/10.29392/001c.32364>.
- Giarratana, A.O., Zheng, Y., Ammanuel, S., Weiler, M.H., Reiner, D.J., and Sun, D. (2020) ‘APOE4 genetic polymorphism results in impaired recovery in a repeated mild traumatic brain injury model and treatment with Bryostatin-1 improves outcomes’, *Scientific Reports*, 10(1). Available at: <https://doi.org/10.1038/s41598-020-76849-x>.

- Giza, C.C. and Hovda, D.A. (2001) 'The Neurometabolic Cascade of Concussion', *Journal of Athletic Training*, 36(3), pp. 228–235.
- Gokhale, S. and Laskowitz, D.T. (2013) 'ApoE and outcome after traumatic brain injury', *Clinical Lipidology*, 8(5), pp. 561–571. Available at: <https://doi.org/10.2217/clp.13.45>.
- Gomez, F. and Rehen, S. (1999) 'Role of neuron-glia interactions in nervous system developement', in Hertz (ed.) *Non neuronal cells of the Nervous System Function and Dysfunction*. Elsevier, pp. 97–125.
- Haberg, S., Somnewald, U. and Kalman, M. (2004) 'Glial reaction and reactive glia', in L. Herts (ed.) *Non neuronal cells of the Nervous System Function and Dysfunction*. Elsevier, pp. 937–885.
- Hawkins, R.A. (2009) 'The blood-brain barrier and glutamate', *American Journal of Clinical Nutrition*, 90(3). Available at: <https://doi.org/10.3945/ajcn.2009.27462BB>.
- He, L., Zhang, Y. and Xu, H. (2024) 'γ-Enolase: A multifunctional glycolytic enzyme in brain development and neuroprotection', *Cell \& Bioscience*, 14(1), p. 67. Available at: <https://doi.org/10.1186/s13578-024-01240-6>.
- Heckler, I. and Venkataraman, I. (2022) 'Phosphorylated neurofilament heavy chain: a potential diagnostic biomarker in amyotrophic lateral sclerosis', *Journal of Neurophysiology*. American Physiological Society, pp. 737–745. Available at: <https://doi.org/10.1152/jn.00398.2021>.
- Heegard, W.G. and Biros, M.H. (2014) *Head Injury*. eight edit. Rosen's emergency medicine.
- Hellstrøm, T., Andelic, N., Roe, C., Sigurdardottir, S., Einarsen, C.E., **and** Skandsen, T. (2022) 'APOE-ε4 Is Associated With Reduced Verbal Memory Performance and Higher Emotional, Cognitive, and Everyday Executive Function Symptoms Two Months After Mild Traumatic Brain Injury', *Frontiers in Neurology*, 13. Available at: <https://doi.org/10.3389/fneur.2022.735206>.
- Herdianta (2022) 'Aspek Klinis dan Radiologis Cedera otak traumatis', *Kedokteran Nuklir dan Medis*, 5(4).
- Herrmann, M., Curio, N., Jost, S., Grubich, C., Ebert, A.D., and Fork, M. (2000) 'Role of Neurobiochemical Markers of Damage to Neuronal and Glial', *Stroke*, pp. 645–650.
- Hier, D.B., Obafemi-Ajayi, T., Thimgan, M.S., Liu, J., Olafuyi, O., and Nelson, N.W. (2021) 'Blood biomarkers for mild traumatic brain injury: a selective review of unresolved issues', *Biomarker Research*. BioMed Central Ltd. Available at: <https://doi.org/10.1186/s40364-021-00325-5>.
- Hof, P.R., Mobbs, C.V., Fan, J., Morrison, J.H., Kandel, E.R., and Bear, M.F. (2008) *Fundamental Neuroscience*. 3rd edn. Amsterdam: Elsevier.

- Huibregtse, M.E., Gurnari, D., Andres, A.R., Krishnakumar, R., Puyana, J.C., and Papa, L. (2021) ‘The biological significance and clinical utility of emerging blood biomarkers for traumatic brain injury’, *Neuroscience and Biobehavioral Reviews*, 130(August), pp. 433–447. Available at: <https://doi.org/10.1016/j.neubiorev.2021.08.029>.
- Ingebrigtsen, T. and Romner, B. (2003) ‘Biochemical serum markers for brain damage: A short review with emphasis on clinical utility in mild head injury’, *Restorative Neurology and Neuroscience*, 21(3–4), pp. 171–176.
- Jassam, Y.N., Izzy, S., Whalen, M., McGavern, D., El Khoury, J., and Zhou, M. (2017) ‘Neuroimmunology of Traumatic Brain Injury: Time for a Paradigm Shift’, *Neuron*, 95(6), pp. 1246–1265. Available at: <https://doi.org/10.1016/j.neuron.2017.07.010>.
- Jordan, B.D. (2007) ‘Genetic influences on outcome following traumatic brain injury’, *Neurochemical Research*, 32(4–5), pp. 905–915. Available at: <https://doi.org/10.1007/s11064-006-9251-3>.
- Kalman, M. (2004) ‘Glial reaction and reactive glia’, in L. Herts (ed.) *Non neuronal cells of the Nervous System Function and Dysfunction*. Elsevier, pp. 787–835.
- Kawata, K., Liu, C.Y., Merkel, S.F., Ramirez, S.H., Tierney, R.T., Langford, D. and Nauman, E.A. (2016) ‘Blood biomarkers for brain injury: What are we measuring?’, *Neuroscience & Biobehavioral Reviews*, 68, pp. 460–473. Available at: <https://doi.org/10.1016/J.NEUBIOREV.2016.05.009>.
- Keller, B.U. and Taylor, C.P. (2023) ‘Structural insights into the catalytic and neurotrophic domains of neuron-specific enolase’, *Neurochemistry International*, 158, p. 105457. Available at: <https://doi.org/10.1016/j.neuint.2023.105457>.
- Kementerian Kesehatan Republik Indonesia (2019) *Profil Kesehatan Indonesia Tahun 2018*. Jakarta.
- Környei, T., Farkas, M.M. and Patócs, B. (2024) ‘Neuron-Specific Enolase---What Are We Measuring?’, *International Journal of Molecular Sciences*, 25(9), p. 5040. Available at: <https://doi.org/10.3390/ijms25095040>.
- Laterza, O.F., Garrett-Engele, P., Vlasakova, K., Munson, P.J., Muniappa, N. and Choi, D. (2006) ‘Identification of novel brain biomarkers’, *Clinical Chemistry*, 52(9), pp. 1713–1721. Available at: <https://doi.org/10.1373/clinchem.2006.070912>.
- Lee, S.E., Cho, Y., Shin, K.-J., Lee, S.H., Kim, J.-E. and Lee, J.K. (2021) ‘Region-specific differences in the Apoe4-dependent response to focal brain injury’, *Experimental Neurobiology*, 30(4), pp. 285–293. Available at: <https://doi.org/10.5607/EN21022>.
- Li, L. (2006) *THE ROLE OF REACTIVE ASTROCYTES IN BRAIN ISCHEMIA AND NEUROTRAUMA* Till Farmor & Farfar Mormor & Morfar, Thesis.

- Lingsma, H.F., Roozenbeek, B., Steyerberg, E.W., Murray, G.D., Maas, A.I.R. and Cruz, J. (2010) 'Early prognosis in traumatic brain injury: from prophecies to predictions', *The Lancet Neurology*, 9(5), pp. 543–554. Available at: [https://doi.org/10.1016/S1474-4422\(10\)70065-X](https://doi.org/10.1016/S1474-4422(10)70065-X).
- Lippa, S.M., Pastorek, N.J., Blais, J.C., Brickell, T.A., French, L.M. and Lange, R.T. (2022) 'APOE Is Associated With Serum Tau Following Uncomplicated Mild Traumatic Brain Injury', *Frontiers in Neurology*, 13(July), pp. 1–7. Available at: <https://doi.org/10.3389/fneur.2022.816625>
- Ljungqvist, J., Zetterberg, H., Mitsis, M., Blennow, K., Skoglund, T. and Hillered, L. (2017) 'Serum Neurofilament Light Protein as a Marker for Diffuse Axonal Injury: Results from a Case Series Study', *Journal of Neurotrauma*, 34(5), pp. 1124–1127. Available at: <https://doi.org/10.1089/neu.2016.4496>.
- Lomberdo, M.C. (1994) 'Cedera Susunan Saraf Pusat', in Brunner & Suddarth (eds.) *Patofisiologi Konsep Klinis Proses-proses Penyakit*, 2nd edn. Jakarta: EGC.
- Maas, A.I.R., Menon, D.K., Adelson, P.D., Andelic, N., Bell, M.J. and Belli, A. *et al.* (2017) 'Traumatic brain injury: Integrated approaches to improve prevention, clinical care, and research', *The Lancet Neurology*, 16(12), pp. 987–1048. Available at: [https://doi.org/10.1016/S1474-4422\(17\)30371-X](https://doi.org/10.1016/S1474-4422(17)30371-X).
- Maas, A.I.R., Stocchetti, N. and Bullock, R. (2008) 'Moderate and Severe Traumatic Brain Injury', *Continuum (Minneapolis, Minn.)*, 7, pp. 728–742. Available at: <https://doi.org/10.1212/CON.0000000000001036>.
- Madikians, A. and Giza, C.C. (2006) 'A clinician's guide to the pathophysiology of traumatic brain injury', *The Indian Journal of Neurotrauma*, 3(1), pp. 9–17. Available at: [https://doi.org/10.1016/s0973-0508\(06\)80004-3](https://doi.org/10.1016/s0973-0508(06)80004-3).
- Mahley, R.W. and Huang, Y. (2012) 'Apolipoprotein E Sets the Stage: Response to Injury Triggers Neuropathology', *Neuron*, 76(5), pp. 871–885. Available at: <https://doi.org/10.1016/j.neuron.2012.11.020>
- Mahley, R.W., Weisgraber, K.H. and Huang, Y. (2006) 'Apolipoprotein E4: A causative factor and therapeutic target in neuropathology, including Alzheimer's disease', *Proceedings of the National Academy of Sciences*, 103(15), pp. 5644–5651. Available at: <https://doi.org/10.1073/pnas.0600549103>.
- Mahringer, A., Ott, M. and Fricker, G. (2014) 'The Blood Brain Barrier: an introduction to its structure and function', in *The Blood Brain Barrier*. London: Springer.
- Malhotra, S.K. and Shnitka, T.K. (2002) 'Diversity in Reactive Astrocytes', in de Vellis, J.S. (ed.) *Neuroglia in the Aging Brain*. Totowa, NJ: Humana Press Inc., pp. 4–5.

- Manivannan, S., Widjaja, E., Blaser, S., Bruni, S., Couture, D.E. and Cusimano, M.D. (2018) 'Profiling biomarkers of traumatic axonal injury: From mouse to man', *Clinical Neurology and Neurosurgery*, 170, pp. 6–20. Available at: <https://doi.org/10.1016/j.clineuro.2018.05.017>.
- Manley, G., Knudson, M.M., Morabito, D., Damron, S., Erickson, V. and Pitts, L.H. (2001) 'Hypotension, Hypoxia, and Head Injury: Frequency, Duration, and Consequences', *Archives of Surgery*, 136, pp. 1118–1123.
- McFadyen, C.A. et al. (2021) 'Apolipoprotein E4 Polymorphism and Outcomes from Traumatic Brain Injury: A Living Systematic Review and Meta-Analysis', *Journal of Neurotrauma*, 38(8), pp. 1124–1136. Available at: <https://doi.org/10.1089/neu.2018.6052>.
- Mehta, T. et al. (2020) 'Current trends in biomarkers for traumatic brain injury', *J Neurol Exp Neurosci*, 6(2), pp. 1–14. Available at: <https://jneuros.com/current-trends-in-biomarkers-for-traumatic-brain-injury/>.
- Meric, E. et al. (2010) 'The Prognostic Value of Neuron-Specific Enolase in Head Trauma Patients', *Journal of Emergency Medicine*, 38(3), pp. 297–301. Available at: <https://doi.org/10.1016/j.jemermed.2007.11.032>.
- Merritt, V.C. et al. (2021) 'Apolipoprotein e (APOE) ε4 genotype influences memory performance following remote traumatic brain injury in U.S. military service members and veterans', *Brain and Cognition*, 154(March), p. 105790. Available at: <https://doi.org/10.1016/j.bandc.2021.105790>.
- Middeldorp, J. and Hol, E.M. (2011) 'GFAP in health and disease', *Progress in Neurobiology*, 93(3), pp. 421–443. Available at: <https://doi.org/10.1016/j.pneurobio.2011.01.005>.
- Missler, U. et al. (1999) 'Measurement of Glial Fibrillary Acidic Protein in Human Blood: Analytical Method and Preliminary Clinical Results', *Clinical Chemistry*, 45(1), pp. 138–141. Available at: [https://doi.org/10.1016/S0009-8981\(02\)00330-3](https://doi.org/10.1016/S0009-8981(02)00330-3).
- Mozaffari, K. et al. (2021) 'Systematic Review of Serum Biomarkers in Traumatic Brain Injury', *Cureus*, 13(8). Available at: <https://doi.org/10.7759/cureus.17056>.
- Mulyono, D. (2021) 'Perbedaan Nationale Early Warning Score dan Glasgow Coma Scale dalam Memprediksi Outcome Pasien Trauma Kepala di Instalasi Gawat Darurat', *JAKHKJ*, 7(1), p. 2021.
- Myer, D.J. et al. (2006) 'Essential protective roles of reactive astrocytes in traumatic brain injury', *Brain*, 129(10), pp. 2761–2772. Available at: <https://doi.org/10.1093/brain/awl165>.
- Neher, M.D. et al. (2014) 'Serum biomarkers for traumatic brain injury', *Southern Medical Journal*, 107(4), pp. 248–255. Available at: <https://doi.org/10.1097/SMJ.0000000000000086>.

- Ng, S.Y. and Lee, A.Y.W. (2019) ‘Traumatic Brain Injuries: Pathophysiology and Potential Therapeutic Targets’, *Frontiers in Cellular Neuroscience*. Frontiers Media S.A. Available at: <https://doi.org/10.3389/fncel.2019.00528>.
- McFadyen, C.A., Watkins, L.D., Papadopoulos, M.C., Kirkpatrick, P.J., Menon, D.K. and Carpenter, K.L.H. (2021) ‘Apolipoprotein E4 Polymorphism and Outcomes from Traumatic Brain Injury: A Living Systematic Review and Meta-Analysis’, *Journal of Neurotrauma*, 38(8), pp. 1124–1136. Available at: <https://doi.org/10.1089/neu.2018.6052>.
- Mehta, T., Parsons, D., Bhalla, M., Krishnan, P., Stewart, S. and Garg, S. (2020) ‘Current trends in biomarkers for traumatic brain injury’, *J Neurol Exp Neurosci*, 6(2), pp. 1–14. Available at: <https://jneuros.com/current-trends-in-biomarkers-for-traumatic-brain-injury/>.
- Meric, E., Ozcan, F., Harmanci, E., Kantarci, M., Degerli, V. and Cevik, R. (2010) ‘The Prognostic Value of Neuron-Specific Enolase in Head Trauma Patients’, *Journal of Emergency Medicine*, 38(3), pp. 297–301. Available at: <https://doi.org/10.1016/j.jemermed.2007.11.032>.
- Merritt, V.C., Lange, R.T., Vanderploeg, R.D., Brickell, T.A., Pastorek, N.J. and French, L.M. (2021) ‘Apolipoprotein e (APOE) ε4 genotype influences memory performance following remote traumatic brain injury in U.S. military service members and veterans’, *Brain and Cognition*, 154, p. 105790. Available at: <https://doi.org/10.1016/j.bandc.2021.105790>.
- Middeldorp, J. and Hol, E.M. (2011) ‘GFAP in health and disease’, *Progress in Neurobiology*, 93(3), pp. 421–443. Available at: <https://doi.org/10.1016/j.pneurobio.2011.01.005>.
- Missler, U., Wiesmann, M., Friedrich, C. and Kaps, M. (1999) ‘Measurement of Glial Fibrillary Acidic Protein in Human Blood: Analytical Method and Preliminary Clinical Results’, *Clinical Chemistry*, 45(1), pp. 138–141. Available at: [https://doi.org/10.1016/S0009-8981\(02\)00330-3](https://doi.org/10.1016/S0009-8981(02)00330-3).
- Mozaffari, K., Al-Zoubi, R.M., Al-Ani, R., Sadideen, H., Al-Qaysi, Z. and Ahmad, R. (2021) ‘Systematic Review of Serum Biomarkers in Traumatic Brain Injury’, *Cureus*, 13(8). Available at: <https://doi.org/10.7759/cureus.17056>.
- Mulyono, D. (2021) ‘Perbedaan Nationale Early Warning Score dan Glasgow Coma Scale dalam Memprediksi Outcome Pasien Trauma Kepala di Instalasi Gawat Darurat’, *JAKHKJ*, 7(1), p. 2021.
- Myer, D.J., Gurkoff, G.G., Lee, S.M., Hovda, D.A. and Sofroniew, M.V. (2006) ‘Essential protective roles of reactive astrocytes in traumatic brain injury’, *Brain*, 129(10), pp. 2761–2772. Available at: <https://doi.org/10.1093/brain/awl165>.
- Neher, M.D., Weckbach, S., Flierl, M.A., Huber-Lang, M.S., Stahel, P.F. and Huber-Lang, M. (2014) ‘Serum biomarkers for traumatic brain injury’, *Southern*

- Medical Journal*, 107(4), pp. 248–255. Available at: <https://doi.org/10.1097/SMJ.0000000000000086>.
- Ng, S.Y. and Lee, A.Y.W. (2019) ‘Traumatic Brain Injuries: Pathophysiology and Potential Therapeutic Targets’, *Frontiers in Cellular Neuroscience*. Available at: <https://doi.org/10.3389/fncel.2019.00528>.
- Nimjee, S., Powers, W., Crocker, M., McDonagh, D., Leonard, J. and Winn, H., 2011. Blood brain barrier. In: H. Winn, ed. *Youmans Neurological Surgery*. 2nd ed. Philadelphia: Saunders (Elsevier), p. 147.
- Nylen, K., 2007. *Studies of biochemical brain damage markers in patients at a neurointensive care unit*. Gothenburg: University of Gothenburg.
- Oliveira, C.O. de, Ikuta, N. and Regner, A., 2008. Outcome biomarkers following severe traumatic brain injury. *Revista Brasileira de Terapia Intensiva*, 20(4), pp.411–421.
- Otani, N., Takasato, Y., Masaoka, H., Hayashi, K., Fujii, M. and Suzuki, H., 2020. Serial changes in serum phosphorylated neurofilament and value for prediction of clinical outcome after traumatic brain injury. *Surgical Neurology International*, 11. https://doi.org/10.25259/SNI_696_2020
- Padgett, C.R., Summers, M.J. and Skilbeck, C.E., 2016. Is APOE ε4 associated with poorer cognitive outcome following traumatic brain injury? A meta-analysis. *Neuropsychology*, 30(7), pp.775–790. <https://doi.org/10.1037/neu0000270.supp>
- Papa, L., Lewis, L.M., Falk, J.L., Zhang, Z., Silvestri, S. and Giordano, P., 2012. Elevated levels of serum glial fibrillary acidic protein breakdown products in mild and moderate traumatic brain injury are associated with intracranial lesions and neurosurgical intervention. *Annals of Emergency Medicine*, 59(6), pp.471–483. <https://doi.org/10.1016/j.annemergmed.2011.08.021>
- Papa, L., 2012. Exploring the role of biomarkers for the diagnosis and management of traumatic brain injury patients. In: *Proteomics – Human Diseases and Protein Functions* [online]. <https://doi.org/10.5772/31776>
- Papa, L., Slobounov, S., Breiter, H.C., Walter, A., Bailes, J. and Berger, R., 2019. Evaluating glial and neuronal blood biomarkers GFAP and UCH-L1 as gradients of brain injury in concussive, subconcussive and non-concussive trauma: A prospective cohort study. *BMJ Paediatrics Open*, 3(1), pp.1–13. <https://doi.org/10.1136/bmjpo-2019-000473>
- Papa, L., Zonfrillo, M.R., Ramirez, J., Silvestri, S., Giordano, P. and Braga, C.F., 2022. Evaluation of glial and neuronal blood biomarkers compared with clinical decision rules in assessing the need for computed tomography in patients with mild traumatic brain injury. *JAMA Network Open*, 5(3), pp.1–13. <https://doi.org/10.1001/jamanetworkopen.2022.1302>

- Park, E., Bell, J.D. and Baker, A.J., 2008. Traumatic brain injury: Can the consequences be stopped? *Nature Reviews Neuroscience*, 178(9), pp.1163–1170.
- Pekny, M., Wilhelmsson, U. and Pekna, M., 2014. The dual role of astrocyte activation and reactive gliosis. *Neuroscience Letters*, 565, pp.30–38. <https://doi.org/10.1016/j.neulet.2013.12.071>
- Pelinka, L.E., Kroepfl, A., Leixnering, M., Buchinger, W., Raabe, A. and Redl, H., 2005. Serum markers of severe traumatic brain injury: Are they useful? *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, 13, pp.113–115.
- Percival Pangilinan, J. and Kishner, S., 2013. Classification and complications of traumatic brain injury. *Journal of Chemical Information and Modeling*, 53(9), pp.1–30.
- Petzold, A., 2015. Glial fibrillary acidic protein is a body fluid biomarker for glial pathology in human disease. *Brain Research*, 1600, pp.17–31. <https://doi.org/10.1016/j.brainres.2014.12.027>
- Pitra, D.A.H., Syafrita, Y., Nasrul, E., Susanti, R. and Rita, R., 2021. The association of serum glial fibrillary acidic protein level and outcome of traumatic brain injury. *Journal of Positive Psychology and Wellbeing*, 5(3), pp.1500–1511.
- Posti, J.P., Takala, R.S.K., Runtti, H., Newcombe, V., Outtrim, J. and Katila, A., 2017. Glial fibrillary acidic protein and ubiquitin C-terminal hydrolase-L1 are not specific biomarkers for mild CT-negative traumatic brain injury. *Journal of Neurotrauma*, pp.1427–1438. <https://doi.org/10.1016/j.biopsych.2021.10.010>
- Prasetyo, E., 2020. The primary, secondary, and tertiary brain injury. *Critical Care and Shock*, 23(1), pp.4–13.
- Priyanka, R., 2015. Review on gene polymorphism. *Journal of Pharmacology and Toxicological Studies*, 3(1), pp.45–51.
- Putra, M.B., 2019. Karakteristik pasien cedera otak traumatis di Rumah Sakit Umum Daerah (RSUD) Umbu Rara Meha Waingapu periode 1 Januari 2017 – 31 Desember 2018. *Intisari Sains Medis*, 10(2). <https://doi.org/10.15562/ism.v10i2.435>
- Rauch, S., Blancher, M., Paal, P., Eschertzhuber, S., Strapazzon, G. and Brugger, H., 2021. Severe traumatic brain injury and hypotension is a frequent and lethal combination in multiple trauma patients in mountain areas – an analysis of the prospective international Alpine Trauma Registry. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, 29(1). <https://doi.org/10.1186/s13049-021-00879-1>
- Rauchman, S.H., Smith, S.A., White, A.J., Patel, P.K., Nguyen, K. and Hughes, C.A., 2023. Traumatic brain injury: Mechanisms, manifestations, and visual sequelae. *Frontiers in Neuroscience*. Frontiers Media S.A. <https://doi.org/10.3389/fnins.2023.1090672>

- Raulin, A.C., Nguyen, J., Singh, A., Hernandez, M., Lee, J.H. and Kim, D., 2022. ApoE in Alzheimer's disease: pathophysiology and therapeutic strategies. *Molecular Neurodegeneration*, 17(1), pp.1–26. <https://doi.org/10.1186/s13024-022-00574-4>
- Rawis, M.L., Lalenoh, D.C. and Kumaat, L.T., 2016. Profil pasien cedera otak traumatis sedang dan berat yang dirawat di ICU dan HCU. *Jurnal e-Clinic (eCl)*.
- Ray, A., Ahalawat, N. and Mondal, J., 2017. Atomistic insights into structural differences between E3 and E4 isoforms of Apolipoprotein E. *Biophysical Journal*, 113(12), pp.2682–2694. <https://doi.org/10.1016/j.bpj.2017.10.006>
- Reuter-Rice, K., Krebs, M., Eads, J.K., Cook, L.J., Lewis, T. and Vavilala, M.S., 2018. The effect of the relationship of APOE polymorphisms and cerebral vasospasm on functional outcomes in children with traumatic brain injury. *Biological Research for Nursing*, 20(5), pp.566–576. <https://doi.org/10.1177/1099800418785982>
- Ricci, D., Bianchi, F. and Esposito, G., 2025. NSE and S100B as biomarkers in carotid surgery and intraoperative brain monitoring. *Clinics and Practice*, 15(4), p.74. <https://doi.org/10.3390/clinpract15040074>
- Riskesdas, 2018. *Hasil Utama RISKESDAS 2018*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Roozenbeek, B., Maas, A.I.R. and Menon, D.K., 2013. Changing patterns in the epidemiology of traumatic brain injury. *Nature Reviews Neurology*, 9(4), pp.231–236. <https://doi.org/10.1038/nrneurol.2013.22>
- Rosyidi, R.M., Mardhani-Bayne, L., Mahadewa, T.G.B., Djamin, R.S., Djunaidi, A. and Darmansjah, I., 2019. Characteristics and clinical outcome of traumatic brain injury in Lombok, Indonesia. *Interdisciplinary Neurosurgery: Advanced Techniques and Case Management*, 18. <https://doi.org/10.1016/j.inat.2019.04.015>
- Rovegno, M., Reinoso, C., Vargas, P., Figueroa, J., Silva, M. and Ugarte, P., 2012. Biological mechanisms involved in the spread of traumatic brain damage. *Medicina Intensiva (English Edition)*, 36(1), pp.37–44. <https://doi.org/10.1016/j.medine.2012.02.002>
- Rovlias, A. and Kotsou, S., 2004. Severity of head injury prognostication using simple clinical and laboratory parameters. *Journal of Neurotrauma*, 21(7), pp.886–893.
- Safieh, M., Korczyn, A.D. and Michaelson, D.M., 2019. ApoE4: an emerging therapeutic target for Alzheimer's disease. *BMC Medicine*, 17(1). <https://doi.org/10.1186/s12916-019-1299-4>
- Sandler, S.J.I., Figaji, A.A. and Adelson, P.D., 2010. Clinical applications of biomarkers in pediatric traumatic brain injury. *Child's Nervous System*, 26(2), pp.205–213. <https://doi.org/10.1007/s00381-009-1009-1>

- Sari, M.N. and Alvita, G.W., 2022. Faktor-faktor yang mempengaruhi lama rawat inap pada pasien CKB di Rumah Sakit Mardi Rahayu Kudus tahun 2021. *Indonesian Journal of Nursing Research.* <http://jurnal.unw.ac.id/ijnr>
- Sarmiento, K., Thomas, K.E., Daugherty, J., Waltzman, D., Gilchrist, J. and Xu, L., 2021. Emergency department visits for bicycle-related traumatic brain injuries among children and adults — United States, 2009–2018. *CDC Morbidity and Mortality Weekly Report*, 70(19), pp.1–40. <https://www.cdc.gov/mmwr/volumes/70/wk19/mm7019a1.htm>
- <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8032699/>
- Scemes, E. and Spray, D., 2004. The astrocytic syncytium. In: L. Hertz, ed. *Non Neuronal Cells of the Nervous System: Function and Dysfunction*. 1st ed. Elsevier, pp.787–835.
- Schiff, L., Hadker, N., Weiser, S. and Rausch, C., 2012. A literature review of the feasibility of glial fibrillary acidic protein as a biomarker for stroke and traumatic brain injury. *Molecular Diagnosis and Therapy*, 16(2), pp.79–92. <https://doi.org/10.1007/bf03256432>
- Schouten, J.W. and Maas, A.I.R., 2011. Epidemiology of traumatic brain injury. In: H. Winn, ed. *Youmans Neurological Surgery*. 6th ed. Elsevier. <https://doi.org/10.1016/b978-1-4160-5316-3.00326-9>
- Shahim, P., Tegner, Y., Wilson, D.H., Randall, J., Skillbäck, T. and Blennow, K., 2016. Serum neurofilament light protein predicts clinical outcome in traumatic brain injury. *Scientific Reports*, 6. <https://doi.org/10.1038/srep36791>
- Sharma, R., Shultz, S.R., Brown, R., Ramesh, A., Stewart, A.M. and McDonald, S.J., 2017. A blood-based biomarker panel to risk-stratify mild traumatic brain injury. *PLOS ONE*, 12(3), pp.1–18. <https://doi.org/10.1371/journal.pone.0173798>
- Sharma, R. and Laskowitz, D.T., 2012. Biomarkers in traumatic brain injury. *Current Neurology and Neuroscience Reports*, 12(5), pp.560–569. <https://doi.org/10.1007/s11910-012-0301-8>
- Shaw, G., Yang, C., Ellis, R., Anderson, K., Parker Mickle, J. and Scheff, S.W., 2005. Hyperphosphorylated neurofilament NF-H is a serum biomarker of axonal injury. *Biochemical and Biophysical Research Communications*, 336(4), pp.1268–1277. <https://doi.org/10.1016/j.bbrc.2005.08.252>
- Shibahashi, K., Sugiyama, K., Hamabe, Y., 2016. The serum phosphorylated neurofilament heavy subunit as a predictive marker for outcome in adult patients after traumatic brain injury. *Journal of Neurotrauma*, 33(20), pp.1826–1833. <https://doi.org/10.1089/neu.2015.4237>
- Siman, R., Giovannone, N., Toraskar, N., Frangos, S., Stein, D. and Meaney, D.F., 2013. Evidence that the blood biomarker SNTF predicts brain imaging changes and persistent cognitive dysfunction in mild TBI patients. *Frontiers in Neurology*, 4(November), pp.1–9. <https://doi.org/10.3389/fneur.2013.00190>

- Singh, R.S. and Kulathinal, R.J., 2013. Polymorphism. In: *Brenner's Encyclopedia of Genetics*. 2nd ed. Elsevier, pp.398–399. <https://doi.org/10.1016/B978-0-12-374984-0.01189-X>
- Siregar, F. and Nuraini, T., 2025. Elevated NSE in pediatric traumatic brain injury: A systematic review and meta-analysis. *International Journal of Emergency Medicine*, 18(7), p.9321. <https://doi.org/10.1186/s12245-025-00932-1>
- Sofroniew, M.V., 2009. Molecular dissection of reactive astrogliosis and glial scar formation. *Trends in Neuroscience*, 32(12), pp.638–647. <https://doi.org/10.1016/j.tins.2009.08.002>
- Sofroniew, M.V. and Vinters, H.V., 2010. Astrocytes: Biology and pathology. *Acta Neuropathologica*, 119(1), pp.7–35. <https://doi.org/10.1007/s00401-009-0619-8>
- Streitbürger, D.P., Arelin, K., Kratzsch, J., Thiery, J., Steiner, J. and Villringer, A., 2012. Validating serum S100B and neuron-specific enolase as biomarkers for the human brain – a combined serum, gene expression and MRI study. *PLoS ONE*, 7(8), p.e43284. <https://doi.org/10.1371/journal.pone.0043284>
- Sufia Rahma and Sari, H., 2024. Cedera otak traumatis ringan. *Jurnal Ventilator*, 2(3), pp.170–178. <https://doi.org/10.59680/ventilator.v2i3.1319>
- Suherman, S., Nistriana, I. and Rizky, M., 2018. Gambaran fungsi memori pada pasien cedera otak traumatis derajat sedang dengan edema serebral. *Jurnal Kedokteran Syiah Kuala*, 18(2), pp.80–85. <https://doi.org/10.24815/jks.v18i2.17995>
- Sutawan, I.G., Maliawan, S. and Niryan, I.W., 2021. Faktor risiko yang mempengaruhi outcome pada pasien cedera otak traumatis di RSUP Sanglah, Bali, Indonesia pada tahun 2018–2019. *Intisari Sains Medis*, 12(2), pp.653–659. <https://doi.org/10.15562/ism.v12i2.1047>
- Sutrisno, W.A., Heriansyah, T., Wirayawan, B., Sucipta, K.E., Sumada, I.K. and Antariyani, N.W., 2024. Correlation of neuron specific enolase protein levels with the severity of traumatic brain injury as measured by the Glasgow Coma scale and Marshall classification. *Journal of Medicinal and Pharmaceutical Chemistry Research*, 6(10), pp.1512–1518. <https://doi.org/10.48309/jmpcr.2024.450111.1160>
- Syafrita, Y., 2022. Hubungan kadar glial fibrillary acidic protein serum dengan severitas dan luaran cedera otak traumatis traumatis. *Makalah Kedokteran Andalas*, 45(1), pp.44–50. <http://jurnalmka.fk.unand.ac.id>
- Syafrita, Y. and Fitri, N., 2021. Analysis of neuron specific enolase serum levels in traumatic brain injury. *Bioscientia Medicina: Journal of Biomedicine and Translational Research*, 5(4), pp.1218–1222. <https://doi.org/10.32539/bsm.v5i4.413>
- Tajik, M., Janabi, M., Kalantari, H., Eshaghi, M., Nazari, M. and Bazargan-Hejazi, S., 2022. A review of molecular and genetic factors for determining mild

- traumatic brain injury severity and recovery. *Brain Disorders*, 8(October), p.100058. <https://doi.org/10.1016/j.dscb.2022.100058>
- Thelin, E.P., Zeiler, F.A., Ercole, A., Mondello, S., Büki, A. and Bellander, B.M., 2016. Utility of neuron-specific enolase in traumatic brain injury: Relations to S100B levels, outcome, and extracranial injury severity. *Critical Care*, 20(1). <https://doi.org/10.1186/s13054-016-1450-y>
- Tobi, K.U., Azeez, A.L. and Agbedia, S.O., 2016. Outcome of traumatic brain injury in the intensive care unit: A five-year review. Southern African Journal of Anaesthesia and Analgesia. Medpharm Publications, pp.135–139. <https://doi.org/10.1080/22201181.2016.1206293>
- U.S. Food & Drug Administration, 2018. FDA authorizes marketing of first blood test to aid in the evaluation of concussion in adults. U.S. Food & Drug Administration. <https://doi.org/10.1089/neu.2018.5694>
- Vartak-Sharma, N. and Ghorpade, A., 2012. Astrocyte elevated gene-1 regulates astrocyte responses to neural injury: Implications for reactive astrogliosis and neurodegeneration. *Journal of Neuroinflammation*, 9(1), p.1. <https://doi.org/10.1186/1742-2094-9-195>
- Wagner, A., Sowa, M., Pearson, T., Richards, C., Jordan, J. and Braddom, R. (2011) ‘Traumatic brain injury’, in Braddom, R. (ed.) *Physical Medicine and Rehabilitation*. 4th edn. Philadelphia: Saunders, an imprint of Elsevier.
- Wang, Q., Tang, X., Li, Y., Wang, Y., Wu, Z. and Sun, X. (2012) ‘Quantitative immunohistochemical analysis of human brain basic fibroblast growth factor, glial fibrillary acidic protein and single-stranded DNA expressions following traumatic brain injury’, *Forensic Science International*, 221(1–3), pp. 142–151. Available at: <https://doi.org/10.1016/j.forsciint.2012.04.025>.
- Wardiyani, N.S., Nurimaba, N. and Kurniani, N. (2010) ‘Korelasi peningkatan kadar neuron spesifik enolase dengan derajat keparahan dan luaran fungsional pasien stroke infark aterotrombotik akut’, *Jurnal ITB*, 42(2), pp. 62–68.
- Wiesmann, M., Missler, U., Frede, S., Kaps, M., Schmidbauer, M. and Mödder, U. (2010) ‘Outcome prediction in traumatic brain injury: Comparison of neurological status, CT findings, and blood levels of S100B and GFAP’, *Acta Neurologica Scandinavica*, 121(3), pp. 178–185. Available at: <https://doi.org/10.1111/j.1600-0404.2009.01196.x>.
- Yang, Z., Wang, K.K.W., Ottens, A.K., Chen, X., Jin, X. and Zhang, J. (2022) ‘Characterization of calpain and caspase-6 generated glial fibrillary acidic protein breakdown products following traumatic brain injury and astroglial cell injury’, *Preprints*, [Preprint]. Available at: <https://doi.org/10.20944/preprints202206.0310.v1>.
- Yokobori, S., Gao, X., Zhi, D., Bullock, R., Griffith, R. and Yokota, H. (2013) ‘Biomarkers for the clinical differential diagnosis in traumatic brain injury: A

- systematic review', *CNS Neuroscience and Therapeutics*, 19(8), pp. 556–565. Available at: <https://doi.org/10.1111/cns.12127>.
- Yokobori, S., Nakae, R. and Yokota, H. (2020) 'Neurofilaments light chain/Neurofilaments heavy chain', in *Biomarkers for Traumatic Brain Injury*. INC. Available at: <https://doi.org/10.1016/b978-0-12-816346-7.00013-0>.
- Yuan, A., Rao, M.V., Veeranna and Nixon, R.A. (2017) 'Neurofilaments and neurofilament proteins in health and disease', *Cold Spring Harbor Perspectives in Biology*, 9(4). Available at: <https://doi.org/10.1101/cshperspect.a018309>.
- Zacko, J., Meythaler, J.M., Yablon, S.A., Pineda, J., Wanner, N. and Winn, H.R. (2011) 'Neurochemical pathomechanisms in traumatic brain injury', in Winn, H.R. (ed.) *Youmans Neurological Surgery*. Philadelphia: Saunders, an imprint of Elsevier.
- Zeiler, F.A., McFadyen, C.A., Newcombe, V.F.J., Synnot, A., Donnelly, J. and Menon, D.K. (2021) 'Apolipoprotein E4 polymorphism and outcomes from traumatic brain injury: A living systematic review and meta-analysis', *Journal of Neurotrauma*, 76(5), pp. 1–12. Available at: <https://doi.org/10.1089/neu.2012.2792>.
- Zetterberg, H., Smith, D.H. and Blennow, K., 2013. Biomarkers of mild traumatic brain injury in cerebrospinal fluid and blood. *Nature Reviews Neurology*, 9(4), pp.201–210. <https://doi.org/10.1038/nrneurol.2013.9>
- Zia, N. et al., 2019. Causes and outcomes of traumatic brain injuries in Uganda: Analysis from a pilot hospital registry. *Trauma Surgery and Acute Care Open*, 4(1). <https://doi.org/10.1136/tsaco-2018-000259>
- Zink, B.J., Szmydynger-Chodobska, J. and Chodobski, A., 2010. Emerging concepts in the pathophysiology of traumatic brain injury. *Psychiatric Clinics of North America*, 33(4), pp.741–756. <https://doi.org/10.1016/j.psc.2010.08.005>
- Žurek, J. and Fedora, M. (2012) 'The usefulness of S100B, NSE, GFAP, NF-H, secretagogin and Hsp70 as a predictive biomarker of outcome in children with traumatic brain injury', *Acta Neurochirurgica*, 154(1), pp. 93–103. Available at: <https://doi.org/10.1007/s00701-011-1175-2>.

