

REFERENCES

- Abenavoli, M. R., Lupini, A., Oliva, S., & Sorgonà, A. (2010). Allelochemical effects on net nitrate uptake and plasma membrane H⁺-ATPase activity in maize seedlings. *Biologia Plantarum*, 54(1), 149–153. <https://doi.org/10.1007/s10535-010-0024-0>
- Ahmad, S., Belwal, V., Punia, S. S., Ram, M., Dalip, Rajput, S. S., Kunwar, R., Meena, M. K., Gupta, D., Kumawat, G. L., Hussain, T., & Mohamed, H. I. (2023). Role of Plant Secondary Metabolites and Phytohormones in Drought Tolerance: A Review. *Gesunde Pflanzen*, 75(4), 729–746. <https://doi.org/10.1007/s10343-022-00795-z>
- Aimar, D., Calafat, M., Andrade, A. M., Carassay, L., Abdala, G. I., & Molas, M. L. (2011). Drought tolerance and stress hormones: From model organisms to forage crops. *Plants and Environment*, 10(6), 137–164.
- Ain, Q., Mushtaq, W., Shadab, M., & Siddiqui, M. B. (2023). Allelopathy: an alternative tool for sustainable agriculture. *Physiology and Molecular Biology of Plants*, 29(4), 495–511. <https://doi.org/10.1007/s12298-023-01305-9>
- Al-Khayri, J. M., Rashmi, R., Toppo, V., Chole, P. B., Banadka, A., Sudheer, W. N., Nagella, P., Shehata, W. F., Al-Mssallem, M. Q., Alessa, F. M., Almaghasla, M. I., & Rezk, A. A. S. (2023). Plant Secondary Metabolites: The Weapons for Biotic Stress Management. *Metabolites*, 13(6), 1–37. <https://doi.org/10.3390/metabo13060716>
- Alluri, G., & Saha, D. (2024). Emerging Perspectives on Non-Chemical Weed Management Tactics in Container Ornamental Production in the United States. *Horticulturae*, 10(3). <https://doi.org/10.3390/horticulturae10030281>
- Amini, S., Azizi, M., Joharchi, M. R., & Moradinezhad, F. (2016). Evaluation of allelopathic activity of 68 medicinal and wild plant species of Iran by Sandwich method. *International Journal of Horticultural Science and Technology*, 3(2), 243–253.
- Araújo, C. A., Morgado, C. S. A., Gomes, A. K. C., Gomes, A. C. C., & Simas, N. K. (2021). Asteraceae family: A review of its allelopathic potential and the case of *Acmella oleracea* and *Sphagneticola trilobata*. *Rodriguesia*, 72. <https://doi.org/10.1590/2175-7860202172137>
- Arif, Y., Singh, P., Siddiqui, H., Bajguz, A., & Hayat, S. (2020). Salinity induced physiological and biochemical changes in plants: An omic approach towards salt stress tolerance. *Plant Physiology and Biochemistry*, 156, 64–77.
- Arora, S., Husain, T., & Prasad, S. M. (2024). Allelochemicals as biocontrol agents: Promising aspects, challenges and opportunities. *South African Journal of Botany*, 166, 503–511. <https://doi.org/10.1016/j.sajb.2024.01.029>
- Arroussi, J., Ouerfelli, M., Smaoui, A., Ahmed, H. Ben, Kaâb, S. Ben, & Kaâb, L. B. Ben. (2022). Antioxidant activity of seven plant extracts collected from Tunisia and their allelopathic potential on *Lactuca sativa* L. and *Phalaris*

- minor* L. *South African Journal of Botany*, 148, 135–143. <https://doi.org/10.1016/j.sajb.2022.04.029>
- Ashraf, M. A., Iqbal, M., Rasheed, R., Hussain, I., Riaz, M., & Arif, M. S. (2018). Environmental stress and secondary metabolites in plants: an overview. In *Plant metabolites and regulation under environmental stress* (pp. 153–167). <https://doi.org/10.1016/B978-0-12-812689-9.00008-X>
- Awal, M. A., Ahsan, A. K. M. A., & Pramanik, M. H. R. (2020). Effect of Aqueous Extract of Grass Weeds on Seed Germination and Seedling Growth of Vegetable Crops. *Asian Journal of Research in Botany*, 4(1), 1–10.
- Awuchi, C. G., & Amagwula, I. O. (2020). The Biochemistry, Toxicology, and Uses of the Pharmacologically Active Phytochemicals: Alkaloids, Terpenes, Polyphenols, and Glycosides. *Merit Research Journals*, 5(1), 6–21. <https://doi.org/10.22146/jfps.666>
- Babaahmadi, H., Ghanbari, A., Asadi, G., & Emami, M. K. (2013). Allelopathic effect from some medicinal plants on germination of *Alyssum hirsutum* and *Amaranthus retroflexus*. *International Journal of Agronomy and Plant Production*, 4(12).
- Babu, M., Abirami, T., Velu, K., Janaki, A., & Fahd, A. (2025). Larvicidal activity and phytochemical composition of *Kyllinga brevifolia* (Poales: Cyperaceae) against mosquito vectors. *Journal of the Kansas Entomological Society*, 98(2), 62–74.
- Bajwa, A. A. (2014). Sustainable weed management in conservation agriculture. *Crop Protection*, 65, 105–113. <https://doi.org/10.1016/j.cropro.2014.07.014>
- Batish, D. R., Singh, H. P., Kaur, S., Kohli, R. K., & Yadav, S. S. (2008). Caffeic acid affects early growth, and morphogenetic response of hypocotyl cuttings of mung bean (*Phaseolus aureus*). *Journal of Plant Physiology*, 165(3), 297–305. <https://doi.org/10.1016/j.jplph.2007.05.003>
- Belgeri, A., & Adkins, S. W. (2015). Allelopathic potential of invasive parthenium weed (*Parthenium hysterophorus* L.) seedlings on grassland species in Australia. *Allelopathy Journal*, 36(1).
- Belz, R. G., Reinhardt, C. F., Foxcroft, L. C., & Hurle, K. (2007). Residue allelopathy in *Parthenium hysterophorus* L.-Does parthenin play a leading role? *Crop Protection*, 26(3), 237–245. <https://doi.org/10.1016/j.cropro.2005.06.009>
- Berendji, S., Asghari, J. B., & Matin, A. A. (2008). Allelopathic potential of rice (*Oryza sativa*) varieties on seedling growth of barnyardgrass (*Echinochloa crus-galli*). *Journal of Plant Interactions*, 3(3), 175–180. <https://doi.org/10.1080/17429140802032855>
- Bergmark, C. L., Jackson, W. A., Volk, R. J., & Blum, U. (1992). Differential inhibition by ferulic acid of nitrate and ammonium uptake in *Zea mays* L. *Plant Physiology*, 98(2), 639–645. <https://doi.org/10.1104/pp.98.2.639>
- Bhadoria, P. (2010). Allelopathy: A Natural Way towards Weed Management. *American Journal of Experimental Agriculture*, 1(1), 7–20.

<https://doi.org/10.9734/ajea/2011/002>

- Bhatla, S. C., & Lal, M. A. (2023). Secondary Metabolites. In *Plant Physiology, Development and Metabolism* (pp. 1099–1166). Springer Nature. https://doi.org/10.1007/978-981-99-5736-1_33
- Bocso, N. S., & Butnariu, M. (2022). The biological role of primary and secondary plants metabolites. *Journal of Nutrition and Food Processing*, 5(3), 1–7.
- Boeckler, G. A., Gershenzon, J., & Unsicker, S. B. (2011). Phenolic glycosides of the Salicaceae and their role as anti-herbivore defenses. *Phytochemistry*, 72(13), 1497–1509. <https://doi.org/10.1016/j.phytochem.2011.01.038>
- Bonanomi, G., Zotti, M., Idbella, M., Mazzoleni, S., & Abd-ElGawad, A. M. (2021). Microbiota modulation of allelopathy depends on litter chemistry: Mitigation or exacerbation? *Science of the Total Environment*, 776, 145942. <https://doi.org/10.1016/j.scitotenv.2021.145942>
- Boncan, D. A. T., Tsang, S. S. K., Li, C., Lee, I. H. T., Lam, H. M., Chan, T. F., & Hui, J. H. L. (2020). Terpenes and terpenoids in plants: Interactions with environment and insects. *International Journal of Molecular Sciences*, 21(19), 1–19. <https://doi.org/10.3390/ijms21197382>
- Bond, W., & Grundy, A. C. (2001). Non-chemical weed management in organic farming systems. *Weed Research*, 41(5), 383–405. <https://doi.org/10.1046/j.1365-3180.2001.00246.x>
- Bryson, C. T., Carter, R., McCarty, L. B., & Yelverton, F. H. (1997). Kyllinga, a Genus of Neglected Weeds in the Continental United States. *Weed Technology*, 11(4), 838–842. <https://doi.org/https://www.jstor.org/stable/3988785>
- Chagas, F. O., Pessotti, R. de C., Caraballo-Rodríguez, A. M., & Pupo, M. T. (2018). Chemical signaling involved in plant–microbe interactions. *Chemical Society Reviews*, 47(5), 1652–1704.
- Chandra, M., Kushwaha, S., Mishra, B., & Sangwan, N. (2022). Molecular and structural insights for the regulation of terpenoids in *Ocimum basilicum* and *Ocimum tenuiflorum*. *Plant Growth Regulation*, 97(1), 61–75. <https://doi.org/10.1007/s10725-022-00796-y>
- Chaniago, I. (2004). *Mechanisms and modes of action of weed interference in soybean at the physiological, biochemical, and cellular levels*. University of New England, Armidale, Australia.
- Cheema, Z. A., & Khaliq, A. (2000). Use of sorghum allelopathic properties to control weeds in irrigated wheat in a semi arid region of Punjab. *Agriculture, Ecosystems and Environment*, 79(2–3), 105–112. [https://doi.org/10.1016/S0167-8809\(99\)00140-1](https://doi.org/10.1016/S0167-8809(99)00140-1)
- Cheng, F., & Cheng, Z. (2015). Research progress on the use of plant allelopathy in agriculture and the physiological and ecological mechanisms of allelopathy. *Frontiers in Plant Science*, 6, 1020. <https://doi.org/10.3389/fpls.2015.01020>

- Chon, S. U. (2013). Total polyphenols and bioactivity of seeds and sprouts in several legumes. *Current Pharmaceutical Design*, 19(34), 6112–6124.
- Chopra, N., Tewari, G., Tewari, L. M., Upreti, B., & Pandey, N. (2017). Allelopathic Effect of *Echinochloa colona* L. And *Cyperus iria* L. Weed Extracts on the Seed Germination and Seedling Growth of Rice and Soyabean. *Advances in Agriculture*, 2017. <https://doi.org/10.1155/2017/5748524>
- Chou, C. H., & Leu, L. L. (1992). Allelopathic substances and interactions of *Delonix regia* (Boj) Raf. *Journal of Chemical Ecology*, 18(12), 2285–2303. <https://doi.org/10.1007/BF00984951>
- Cipollini, M. L. (2000). Secondary metabolites of vertebrate-dispersed fruits: evidence for adaptive functions. *Revista Chilena de Historia Natural*, 73(3), 421–440.
- Deng, W., Yang, Q., Chen, Y., Yang, M., Xia, Z., Zhu, J., Chen, Y., Cai, J., & Yuan, S. (2020). Cyhalofop-butyl and Glyphosate Multiple-Herbicide Resistance Evolved in an *Eleusine indica* Population Collected in Chinese Direct-Seeding Rice. *Journal of Agricultural and Food Chemistry*, 68(9), 2623–2630. <https://doi.org/10.1021/acs.jafc.9b07342>
- Devi, S. R., & Prasad, M. N. V. (1992). Effect of ferulic acid on growth and hydrolytic enzyme activities of germinating maize seeds. *Journal of Chemical Ecology*, 18(11), 1981–1990. <https://doi.org/10.1007/BF00981921>
- Ding, H., Ahmad, I., Ali, A., Khan, A. R., & Cheng, Z. (2018). Bioassay evaluation of the potential allelopathic effects of garlic (*Allium sativum* L.) root exudates on lettuce and cucumber. *Pakistan Journal of Botany*, 50(1), 371–380.
- Divekar, P. A., Narayana, S., Divekar, B. A., Kumar, R., Gadratagi, B. G., Ray, A., Singh, A. K., Rani, V., Singh, V., Singh, A. K., Kumar, A., Singh, R. P., Meena, R. S., & Behera, T. K. (2022). Plant Secondary Metabolites as Defense Tools against Herbivores for Sustainable Crop Protection. *International Journal of Molecular Sciences*, 23(5). <https://doi.org/10.3390/ijms23052690>
- Dubey, R., Gupta, D. K., & Sharma, G. K. (2020). Chemical stress on plants. In *New frontiers in stress management for durable agriculture* (pp. 101–128). Singapore: Springer Singapore. https://doi.org/10.1007/978-981-15-1322-0_7
- Duke, S. O., Blair, A. C., Dayan, F. E., Johnson, R. D., Meepagala, K. M., Cook, D., & Bajsa, J. (2009). Is (-)-catechin a novel weapon of spotted Knapweed (*Centaurea stoebe*)? *Journal of Chemical Ecology*, 35(2), 141–153. <https://doi.org/10.1007/s10886-008-9587-z>
- Ehrenfeld, J. G. (2010). Ecosystem consequences of biological invasions. *Annual Review of Ecology, Evolution, and Systematics*, 41(1), 59–80. <https://doi.org/10.1146/annurev-ecolsys-102209-144650>
- Einhellig, F. A. (1994). Allelopathy: Current Status and Future Goals. *American*

- Chemical Society*, 1–24. <https://doi.org/10.1021/bk-1995-0582.ch001>
- El-Masry, R. R., Ahmed, S. A. A., El-Rokiek, K. G., Messiha, N. K., & Mohamed, S. A. (2019). Allelopathic activity of the leaf powder of *Ficus nitida* on the growth and yield of *Vicia faba* and associated weeds. *Bulletin of the National Research Centre*, 43, 1–7. <https://doi.org/10.1186/s42269-019-0114-x>
- Elsayed, G. (2011). Plant secondary substances and insects behaviour. *Archives of Phytopathology and Plant Protection*, 44(16), 1534–1549. <https://doi.org/10.1080/03235408.2010.507957>
- Frerot, B., Leppik, E., Groot, A. T., Unbehend, M., & Holopainen, J. K. (2017). Chemical signatures in plant–insect interactions. In *Advances in botanical research* (pp. 139–177). Academic Press. <https://doi.org/10.1016/bs.abr.2016.10.003>
- Fritz, D., Bernardi, A. P., Haas, J. S., Ascoli, B. M., De Bordignon, S. A. L., & Von Poser, G. (2007). Germination and growth inhibitory effects of *Hypericum myrianthum* and *H. polyanthemum* extracts on *Lactuca sativa* L. *Revista Brasileira de Farmacognosia*, 17(1), 44–48. <https://doi.org/10.1590/S0102-695X2007000100010>
- Fu, F., Luo, J., Zhao, L., Yang, F., & Wang, N. (2024). Impact of cellulose and lignin on restoration of vegetation and soil chemical properties for saline-alkali soil of songnen plain. *Plos One*, 19(1), 1–12. <https://doi.org/10.1371/journal.pone.0296366>
- Fujii, Y. (1990). Survey of Japanese weeds and crops for the detection of water-extractable allelopathic chemicals using Richards' function fitted to lettuce germination test. *Weed Res. Jpn*, 35, 362–370.
- Fujii, Y., Parvez, S. S., Parvez, M. M., Ohmae, Y., & Iida, O. (2003). Screening of 239 medicinal plant species for allelopathic activity using the sandwich method. *Weed Biology and Management*, 3(4), 233–241.
- Fujii, Y., Shibuya, T., Nakatani, K., Itani, T., Hiradate, S., & Parvez, M. M. (2004). Assessment method for allelopathic effect from leaf litter leachates. *Weed Biology and Management*, 4(1), 19–23. <https://doi.org/10.1111/j.1445-6664.2003.00113.x>
- Gholamhoseini, M., AghaAlikhani, M., Sanavy, S. A. M. M., Mirlatifi, S. M., & Zakikhani, H. (2013). Response of corn and redroot pigweed to nitrogen fertilizer in different irrigation regimes. *Agronomy Journal*, 105(4), 1107–1118. <https://doi.org/10.2134/agronj2012.0499>
- Greer, M. J., Wilson, G. W. T., Hickman, K. R., & Wilson, S. M. (2014). Experimental evidence that invasive grasses use allelopathic biochemicals as a potential mechanism for invasion: chemical warfare in nature. *Plant and Soil*, 385(1–2), 165–179. <https://doi.org/10.1007/s11104-014-2209-3>
- Handayani, A., Junaedi, D. I., & Zuhud, E. A. M. (2021). Ecological risk assessment of potentially invasive alien plant species in Cibodas Biosphere Reserve, West Java, Indonesia. *IOP Conference Series: Earth and Environmental Science*, 914(1), 012035. <https://doi.org/10.1088/1755->

1315/914/1/012035

- Harborne, J. B. (2001). Secondary Metabolites: Attracting Pollinators. *Encyclopedia of Life Sciences*, 1–5. <https://doi.org/10.1038/npg.els.0000909>
- Harborne, J. B., & Mabry, T. J. (2013). *The flavonoids: advances in research*.
- Hejl, A. M., & Koster, K. L. (2004). The allelochemical sorgoleone inhibits root H⁺-ATPase and water uptake. *Journal of Chemical Ecology*, 30(11), 2181–2191. <https://doi.org/10.1023/B:JOEC.0000048782.87862.7f>
- Hernández-Aro, M., Hernández-Pérez, R., Guillén-Sánchez, D., & Torres-Garcia, S. (2016). A influência dos resíduos de *Sphagneticola trilobata* em plantas daninhas e culturas. *Planta Daninha*, 34(1), 81–90. <https://doi.org/10.1590/S0100-83582016340100008>
- Horvath, D. P., Clay, S. A., Swanton, C. J., Anderson, J. V., & Chao, W. S. (2023). Weed-induced crop yield loss: a new paradigm and new challenges. *Trends in Plant Science*, 28(5), 567–582. <https://doi.org/10.1016/j.tplants.2022.12.014>
- Hosni, K., Hassen, I., Sebei, H., & Casabianca, H. (2013). Secondary metabolites from *Chrysanthemum coronarium* (Garland) flowerheads: Chemical composition and biological activities. *Industrial Crops and Products*, 44(2013), 263–271. <https://doi.org/10.1016/j.indcrop.2012.11.033>
- Hussain, F., Rasool, A., Aziz, K., Raisham, S., Aziz, S., Badshah, L., & Hussain, W. (2020). Allelopathic inhibition of germination, seedling growth and cell division of selected plant species by *Calotropis procera* (Ait.) Ait. *Plant Science Today*, 7(1), 1–8. <https://doi.org/10.14719/PST.2020.7.1.606>
- Hussain, M. I., El-Sheikh, M. A., & Reigosa, M. J. (2020). Allelopathic potential of Aqueous extract from *Acacia melanoxylon* r. Br. on *Lactuca sativa*. *Plants*, 9(9), 1–13. <https://doi.org/10.3390/plants9091228>
- Hussain, M. I., & Reigosa, M. J. (2011). Allelochemical stress inhibits growth, leaf water relations, PSII photochemistry, non-photochemical fluorescence quenching, and heat energy dissipation in three C₃ perennial species. *Journal of Experimental Botany*, 62(13), 4533–4545. <https://doi.org/10.1093/jxb/err161>
- Iannucci, A., Fragasso, M., Platani, C., & Papa, R. (2013). Plant growth and phenolic compounds in the rhizosphere soil of wild oat (*Avena fatua* L.). *Frontiers in Plant Science*, 4(DEC), 1–7. <https://doi.org/10.3389/fpls.2013.00509>
- Ilori, O. J., Otusanya, O. O., Adelusi, A. A., & Sanni, R. O. (2010). Allelopathic activities of some weeds in the Asteraceae family. *International Journal of Botany*, 6(2), 161–163.
- Imad, M., Idrees, M., Hadi, F., Memon, N. H., & Zhang, Z. (2021). Allelopathic Effect of *Parthenium hysterophorus* Extract on Seed Germination and Seedling Growth of Selected Plants. *Pakistan Journal of Botany*, 53(6), 2187–2197. [https://doi.org/10.30848/PJB2021-6\(9\)](https://doi.org/10.30848/PJB2021-6(9))
- Inderjit. (2001). Soil: environmental effects on allelochemical activity. *Agronomy Journal*, 93(1), 79–84.

- Inderjit, S., & Gross, E. (2000). Plant phenolics: potential role in aquatic and terrestrial ecosystems. *Proceedings of the Polyphenols*.
- Iqbal, A., & Fry, S. C. (2012). Potent endogenous allelopathic compounds in *Lepidium sativum* seed exudate: Effects on epidermal cell growth in *Amaranthus caudatus* seedlings. *Journal of Experimental Botany*, 63(7), 2595–2604. <https://doi.org/10.1093/jxb/err436>
- Ishak, M. S., Ain, M. N., Sahid, I., & Mardiana-Jansar, K. (2021). Allelopathic screening of Malaysian noxious weeds and several medicinal plants as potential alleloherbicides. *Journal of Environmental Biology*, 42, 762–774.
- Jabran, K., & Farooq, M. (2012). Implications of potential allelopathic crops in agricultural systems. In *Allelopathy: Current Trends and Future Applications* (pp. 349–385). Berlin, Heidelberg: Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-642-30595-5>
- Jabran, K., Mahajan, G., Sardana, V., & Chauhan, B. S. (2015). Allelopathy for weed control in agricultural systems. *Crop Protection*, 72, 57–65. <https://doi.org/10.1016/j.cropro.2015.03.004>
- Jose, A. M., & Shaji, A. G. and F. (2020). Allelopathic Effects Of Aqueous Leaf Extracts Of Two Invasive Plants (*Chromolaena odorata* (L.) R. M. King & H. Rob. And *Sphagneticola trilobata* (L.) Pruski) On Seed Germination Of *Amaranthus cruentus* L Antu. *Journal of Advanced Scientific Research*, 11(4), 198–201. http://www.sciensage.info/journal/1359303580JASR_3006121.pdf
- Judžentienė, A., Būdienė, J., Stancelytė, D., & Nedveckytė, I. (2024). Phytochemistry and Allelopathic Effects of *Tanacetum vulgare* L. (Tansy) Extracts on *Lepidium sativum* L. (Garden Pepper Cress) and *Lactuca sativa* L. (Lettuce). *Horticulturae*, 10(6), 538. <https://doi.org/10.3390/horticulturae10060538>
- Kashyap, P., Shikha, D., Gautam, S., & Rani, U. (2023). *Eleusine indica*. In *Harvesting Food from Weeds* (pp. 113–141).
- Kato-Noguchi, H., & Kato, M. (2023). Evolution of the Secondary Metabolites in Invasive Plant Species *Chromolaena odorata* for the Defense and Allelopathic Functions. *Plants*, 12(3), 6–12. <https://doi.org/10.3390/plants12030521>
- Kato-Noguchi, H., & Peters, R. J. (2013). The Role of Momilactones in Rice Allelopathy. *Journal of Chemical Ecology*, 39(2), 175–185. <https://doi.org/10.1007/s10886-013-0236-9>
- Kawabata, O., Nishimoto, R. K., & Tang Chung Shih. (1994). Interference of two *Kyllinga* species (*Kyllinga nemoralis* and *Kyllinga brevifolia*) on bermudagrass (*Cynodon dactylon*) growth. *Weed Technology*, 8(1), 83–86. <https://doi.org/10.1017/s0890037x00039245>
- Kelton, J. A., & Price, A. J. (2011). Weed science and management. *Soils, Plant Growth and Crop Production, Encyclopedia of Life Support Systems (EOLSS)*, 76–101.
- Khan, A., Kanwal, F., Ullah, S., Fahad, M., Tariq, L., Altaf, M. T., Riaz, A., &

- Zhang, G. (2025). Plant Secondary Metabolites—Central Regulators Against Abiotic and Biotic Stresses. *Metabolites*, 15(4), 1–27. <https://doi.org/10.3390/metabo15040276>
- Khang, D. T., Anh, L. H., Thu Ha, P. T., Tuyen, P. T., Quan, N. Van, Minh, L. T., Quan, N. T., Minh, T. N., Xuan, T. D., Khanh, T. D., & Trung, K. H. (2016). Allelopathic Activity of Dehulled Rice and its Allelochemicals on Weed Germination. *International Letters of Natural Sciences*, 58, 1–10. <https://doi.org/10.56431/p-61d2tq>
- Khatri, K., Negi, B., Bargali, K., & Bargali, S. S. (2024). Toxicological assessment of invasive *Ageratina adenophora* on germination and growth efficiency of native tree and crop species of Kumaun Himalaya. *Ecotoxicology*. <https://doi.org/10.1007/s10646-024-02768-6>
- Klingman, G. C., Ashton, F. M., & Noordhoff, L. J. (1975). *Weed science: principles and practice*.
- Komai, K., & Tang, C. S. (1989). Chemical constituents and inhibitory activities of essential oils from *Cyperus brevifolius* and *C. kyllingia*. *Journal of Chemical Ecology*, 15(8), 2171–2176. <https://doi.org/10.1007/BF01014107>
- Koodkaew, I., & Rottasa, R. (2017). Allelopathic effects of giant sensitive plant (*Mimosa pigra*) leaf powder on germination and growth of popping pod and purslane. *International Journal of Agriculture and Biology*, 19(5), 1113–1118.
- Kostina-Bednarz, M., Płonka, J., & Barchanska, H. (2023). Allelopathy as a source of bioherbicides: challenges and prospects for sustainable agriculture. *Reviews in Environmental Science and Biotechnology*, 22(2), 471–504. <https://doi.org/10.1007/s11157-023-09656-1>
- Křístková, E., Doležalová, I., Lebeda, A., Vinter, V., & Novotná, A. (2008). Description of morphological characters of lettuce (*Lactuca sativa* L.) genetic resources. *Horticultural Science*, 35(3), 113–129. <https://doi.org/10.17221/4/2008-hortsci>
- Kubiak, A., Wolna-Maruwka, A., Niewiadomska, A., & Pilarska, A. A. (2022). The problem of weed infestation of agricultural plantations vs. the assumptions of the European biodiversity strategy. *Agronomy*, 12(8), 1808.
- Kumar, S., Abedin, M. M., Singh, A. K., & Das, S. (2020). Role of phenolic compounds in plant-defensive mechanisms. In *Plant phenolics in sustainable agriculture: volume 1* (pp. 517–532). Springer Singapore. https://doi.org/10.1007/978-981-15-4890-1_22
- Kumar, S., Kumari, S., Rana, S. S., Rana, R. S., Anwar, T., Qureshi, H., Saleh, M. A., Alamer, K. H., Attia, H., Ercisli, S., & Aghayeva, S. (2024). Weed management challenges in modern agriculture: The role of environmental factors and fertilization strategies. *Crop Protection*, 185, 106903. <https://doi.org/10.1016/j.cropro.2024.106903>
- Kumar, S., & Pandey, A. K. (2013). Chemistry and biological activities of flavonoids: An overview. *The Scientific World Journal*, 2013. <https://doi.org/10.1155/2013/162750>

- Kupcinskiene, E., Budreviciute, R., Jasionyte, V., Simanaviciute, L., Jociene, L., Krokaite-kudakiene, E., Rekasius, T., & Marozas, V. (2024). Allelopathic Properties of the Species Comprising Communities of Invasive *Impatiens* spp. and Antioxidant System of Invaders' Populations. *Diversity*, 17(1), 1–25.
- Kurek-Górecka, A., Rzepecka-Stojko, A., Górecki, M., Stojko, J., Sosada, M., & Swierczek-Zieba, G. (2014). Structure and antioxidant activity of polyphenols derived from propolis. *Molecules*, 19(1), 78–101. <https://doi.org/10.3390/molecules19010078>
- Kurniadie, D., Widiyanto, R., Umiyati, U., Widayat, D., Nasahi, C., & Budiawan, A. (2023). Management of *Eleusine indica* (L.) Gaertn Resistance to Glyphosate Herbicide in Indonesia. *Agronomy*, 13(6), 1–10. <https://doi.org/10.3390/agronomy13061649>
- Kurze, E., Wüst, M., Liao, J., McGraphery, K., Hoffmann, T., Song, C., & Schwab, W. (2022). Structure–function relationship of terpenoid glycosyltransferases from plants. *Natural Product Reports*, 39(2), 389–409.
- Lambers, H., & Oliveira, R. S. (2019). Biotic influences: ecological biochemistry: allelopathy and defense against herbivores. In *Plant physiological ecology* (pp. 541–581). Cham: Springer International Publishing.
- Li, J., Chen, L., Chen, Q., Miao, Y., Peng, Z., Huang, B., Guo, L., Liu, D., & Du, H. (2021). Allelopathic effect of *Artemisia argyi* on the germination and growth of various weeds. *Scientific Reports*, 11(1), 4303. <https://doi.org/10.1038/s41598-021-83752-6>
- Li, Y., Kong, D., Fu, Y., Sussman, M. R., & Wu, H. (2020). The effect of developmental and environmental factors on secondary metabolites in medicinal plants. *Plant Physiology and Biochemistry*, 148(January), 80–89. <https://doi.org/10.1016/j.plaphy.2020.01.006>
- Li, Z. H., Wang, Q., Ruan, X., Pan, C. De, & Jiang, D. A. (2010). Phenolics and plant allelopathy. *Molecules*, 15(12), 8933–8952. <https://doi.org/10.3390/molecules15128933>
- Lisar, S. Y., Rahman, I. M., Hossain, M. M., & Motafakkerazad, R. (2012). Causes, effects and responses. In *Water stress*. INTECH Open Access.
- Loffredo, E., Monaci, L., & Senesi, N. (2005). Humic substances can modulate the allelopathic potential of caffeic, ferulic, and salicylic acids for seedlings of lettuce (*Lactuca sativa* L.) and tomato (*Lycopersicon esculentum* Mill.). *Journal of Agricultural and Food Chemistry*, 53(24), 9424–9430. <https://doi.org/10.1021/jf050924b>
- Lotina-Hennsen, B., King-Diaz, B., Aguilar, M. I., & Hernandez Terrones, M. G. (2006). Plant secondary metabolites. Targets and mechanisms of allelopathy. In *Allelopathy: A physiological process with Ecological implications* (pp. 229–265). Dordrecht: Springer Netherlands. https://doi.org/10.1007/1-4020-4280-9_11
- M'barek, K., Zribi, I., & Haouala, R. (2018). Allelopathic effects of *Tetralinis*

- articulata on barley, lettuce, radish and tomato. *Allelopathy Journal*, 43(2), 187–202. <https://doi.org/10.26651/allelo.j./2018-43-2-1140>
- Ma, J., Xie, Y., Yang, Y., Jing, C., You, X., Yang, J., Sun, C., Qin, S., Chen, J., Cao, K., Huang, J., & Li, Y. (2022). AMF colonization affects allelopathic effects of *Zea mays* L. root exudates and community structure of rhizosphere bacteria. *Frontiers in Plant Science*, 13(November), 1–15. <https://doi.org/10.3389/fpls.2022.1050104>
- McCutcheon, S. C., & Jorgensen, S. E. (2008). Phytoremediation. In *Encyclopedia of Ecology* (pp. 2751–2766). Academic Press. <https://doi.org/https://doi.org/10.1016/B978-008045405-4.00069-0>
- Mén, T. T., Cường, N. Q., Thư, N. T. A., Quyên, P. L. T., Phương, P. C., Lê, C. T. N., Yên, N. Đ. H., & Khang, Đ. T. (2019). Nghiên cứu khả năng ức chế nảy mầm hạt của cao chiết xuất từ cây sài đất ba thùy (*Wedelia trilobata* (L.) Hitchc). *Can Tho University Journal of Science*, 55(1), 85–90. <https://doi.org/10.22144/ctu.jsi.2019.011>
- Mirmostafae, S., Azizi, M., & Fujii, Y. (2020). Study of allelopathic interaction of essential oils from medicinal and aromatic plants on seed germination and seedling growth of lettuce. *Agronomy*, 10(2), 163. <https://doi.org/10.3390/agronomy10020163>
- Misra, D., Dutta, W., Jha, G., & Ray, P. (2023). Interactions and Regulatory Functions of Phenolics in Soil-Plant-Climate Nexus. *Agronomy*, 13(2), 1–18. <https://doi.org/10.3390/agronomy13020280>
- Molisch, H. (1938). Der Einfluss einer Pflanze auf die Andere, Allelopathie. *Nature*, 141(3568), 493–493.
- Monteiro, A., & Santos, S. (2022). Sustainable Approach to Weed Management: The Role of Precision Weed Management. *Agronomy*, 12(1), 118. <https://doi.org/10.3390/agronomy12010118>
- Morais, C. D., Santos, L. D., & Rossetto, C. V. (2014). Oil radish development agronomic affected by sunflower plants residues. *Bioscience Journal*, 30(1), 117–128.
- Morikawa, C. I. O., Miyaura, R., de Lourdes Tapia y Figueroa, M., Rengifo Salgado, E. L., & Fujii, Y. (2012). Screening of 170 Peruvian plant species for allelopathic activity by using the Sandwich Method. *Weed Biology and Management*, 12(1), 1–11. <https://doi.org/10.1111/j.1445-6664.2011.00429.x>
- Mukaromah, A. S., Purwestri, Y. A., & Fujii, Y. (2017). Determination of Allelopathic Potential in Mahogany (*Swietenia macrophylla* King) Leaf Litter Using Sandwich Method. *Indonesian Journal of Biotechnology*, 21(2), 93. <https://doi.org/10.22146/ijbiotech.16456>
- Mushtaq, W., Siddiqui, M. B., Hakeem, K. R., Mushtaq, W., Siddiqui, M. B., & Hakeem, K. R. (2020). Allelopathy potential of weeds belonging to the family. In *Allelopathy: Potential for Green Agriculture* (pp. 37–43). Springer. <https://doi.org/https://doi.org/10.1007/978-3-030-40807-7>
- Nie, C., Luo, S., Zeng, R., Mo, M., Li, H., & Lin, C. (2005). Allelopathic potential

- of *Wedelia trilobata* L.: effects on germination, growth and physiological parameters of rice. In *Proceedings of the 4th World Congress on Allelopathy, "Establishing the Scientific Base", Wagga Wagga, New South Wales, Australia, 21-26 August 2005* (pp. 469–472). Centre for Rural Social Research, Charles Sturt University.
- Nouri, H., Talab, Z. A., & Tavassoli, A. (2013). Effect of weed allelopathic of sorghum (*Sorghum halepense*) on germination and seedling growth of wheat, Alvand cultivar. *Annals of Biological Research*, 3(3), 1283–1293. <http://scholarsresearchlibrary.com/archive.html>
- Noushahi, H. A., Khan, A. H., Noushahi, U. F., Hussain, M., Javed, T., Zafar, M., Batool, M., Ahmed, U., Liu, K., Harrison, M. T., Saud, S., Fahad, S., & Shu, S. (2022). Biosynthetic pathways of triterpenoids and strategies to improve their Biosynthetic Efficiency. *Plant Growth Regulation*, 97(3), 439–454. <https://doi.org/10.1007/s10725-022-00818-9>
- Nuaman, R. S. (2023). Detection of Active Chemicals in *Zea mays* Waste and Separation Allelopathic Substances Utilizing Hplc Technology. *World Journal of Pharmaceutical Science and Research*, 2(6), 252–257.
- Oerke, E. C. (2006). Crop losses to pests. *The Journal of Agricultural Science*, 144(1), 31–43.
- Ogata, T., Hamachi, M., & Nishi, K. (2008). Organic Herbicide for Paddy Field. *Japan Patent*, 2008050329.
- Okon, O. G. (2019). Effect of salinity on physiological processes in plants. In *Microorganisms in saline environments: strategies and functions* (pp. 237–262). Cham: Springer International Publishing.
- Oliveira, A., & Romão, N. (2015). Growth Inhibition and Pro-apoptotic Action of *Eleusine indica* (L) Gaertn Extracts in *Allium test*. *European Journal of Medicinal Plants*, 8(3), 121–127. <https://doi.org/10.9734/ejmp/2015/18099>
- Othman, M. R., Leong, S. T., Bakar, B., Awang, K., & Annuar, M. S. M. (2012). Allelopathic Potentials of *Cuscuta campestris* Yuncker Extracts on Germination and Growth of Radish (*Raphanus sativus* L.) and Lettuce (*Lactuca sativa* L.). *Journal of Agricultural Science*, 4(9), 57–63. <https://doi.org/10.5539/jas.v4n9p57>
- Pannacci, E., Masi, M., Farneselli, M., & Tei, F. (2020). Evaluation of mugwort (*Artemisia vulgaris* L.) aqueous extract as a potential bioherbicide to control *Amaranthus retroflexus* L. in Maize. *Agriculture*, 10(12), 642. <https://doi.org/10.3390/agriculture10120642>
- Patterson, D. T. (1981). Effects of allelopathic chemicals on growth and physiological responses of soybean (*Glycine max*). *Weed Science*, 29(1), 53–59.
- Peerzada, A. M. (2017). Biology, agricultural impact, and management of *Cyperus rotundus* L.: the world's most tenacious weed. *Acta Physiologiae Plantarum*, 39(12), 270. <https://doi.org/10.1007/s11738-017-2574-7>
- Perera, K. R. S., Ratnayake, R. M. C. S., & Epa, U. P. K. (2023). Allelopathic

- effects of the invasive plant *Wedelia* (*Sphagneticola trilobata* L.) aqueous extract on common beans (*Phaseolus vulgaris* L.). *Journal of Experimental Biology and Agricultural Sciences*, 11(3), 542–549. [https://doi.org/10.18006/2023.11\(3\).542.549](https://doi.org/10.18006/2023.11(3).542.549)
- Phuong, N. N. T., Độ, N. Đ., Tài, N. H. T., Phương, P. T., & Đức, T. H. (2017). Khảo sát hàm lượng flavonoid, alkaloid và khả năng kháng khuẩn của cao chiết cỏ Mần Trầu (*Eleusine indica*). *Can Tho University, Journal of Science*, 53, 54. <https://doi.org/10.22144/ctu.jvn.2017.157>
- Pollastri, S., & Tattini, M. (2011). Flavonols: Old compounds for old roles. *Annals of Botany*, 108(7), 1225–1233. <https://doi.org/10.1093/aob/mcr234>
- Purvis, C. E. (1990). Allelopathy: a new direction in weed control. *Plant Protection Quarterly*, 5(2), 55–59.
- Qi, S. S., Dai, Z. C., Zhai, D. L., Chen, S. C., Si, C. C., Huang, P., Wang, R. P., Zhong, Q. X., & Du, D. L. (2014). Curvilinear effects of invasive plants on plant diversity: Plant community invaded by *Sphagneticola trilobata*. *PLoS ONE*, 9(11), 1–16. <https://doi.org/10.1371/journal.pone.0113964>
- Qin, C., Nagai, M., Hagins, W., & Hobbs, R. (2011). The allelopathic effects of juglone containing nuts. *The Journal of Experimental Secondary Science*, 1. www.jes2s.com/pdfs/qin_article.com
- Rahardiyan, D., Poluakan, M., Moko, E. M., & Ngangi, J. (2019). The fractions of phenolic and flavonoid compounds of the leaves of north Sulawesi's bashful plant (*Mimosa pudica*). *Journal of Physics: Conference Series*, 1317(1). <https://doi.org/10.1088/1742-6596/1317/1/012066>
- Ramírez-Gómez, X. S., Jiménez-García, S. N., Campos, V. B. C., & García, M. L. (2019). Plant metabolites in plant defense against pathogens. In *Plant diseases-current threats and management trends*. IntechOpen.
- Randall, J. M. (1997). Defining weeds of natural areas. In *Assessment and management of plant invasions* (pp. 18–25). NY: Springer New York.
- Rao, A. N. (2021). *Echinochloa colona* and *Echinochloa crus-galli*. In *Biology and Management of Problematic Crop Weed Species* (pp. 197–239). Academic Press.
- Rashid, M. H., Asaeda, T., & Uddin, M. N. (2010). The Allelopathic Potential of Kudzu (*Pueraria montana*). *Weed Science*, 58(1), 47–55. <https://doi.org/10.1614/ws-09-106.1>
- Rasul, R. A., Ahmed, O. H., & Tofiq, S. I. (2024). Effect of Herbicides Application on Weed Growth and Subsequent Important on Yield of Three Wheat Varieties. *Diyala Agricultural Sciences Journal*, 16(1), 8–25.
- Razak, A. M., Tan, J. K., Mohd Said, M., & Makpol, S. (2023). Modulating Effects of Zingiberaceae Phenolic Compounds on Neurotrophic Factors and Their Potential as Neuroprotectants in Brain Disorders and Age-Associated Neurodegenerative Disorders: A Review. *Nutrients*, 15(11). <https://doi.org/10.3390/nu15112564>
- Rehman, S., Shahzad, B., Bajwa, A. A., Hussain, S., Rehman, A., Cheema, S. A.,

- Abbas, T., Ali, A., Shah, L., Adkins, S., & Li, P. (2019). Utilizing the Allelopathic Potential of Brassica Species for Sustainable Crop Production: A Review. *Journal of Plant Growth Regulation*, 38(1), 343–356. <https://doi.org/10.1007/s00344-018-9798-7>
- Rezende, F. M., Ferreira, M. J. P., Clausen, M. H., Rossi, M., & Furlan, C. M. (2019). Acylated flavonoid glycosides are the main pigments that determine the flower colour of the brazilian native tree *Tibouchina pulchra* (Cham.) Cogn. *Molecules*, 24(4). <https://doi.org/10.3390/molecules24040718>
- Rice, E. L. (1974). *Allelopathy*. Academic Press.
- Rice, E. L. (1984). *Allelopathy*, 2nd ed. Academic Press.
- Rietveld, W. J. (1983). Allelopathic effects of juglone on germination and growth of several herbaceous and woody species. *Journal of Chemical Ecology*, 9, 295–308.
- Rodiyati, A., & Nakagoshi, N. (2003). Annual growth and phenology of *Kyllinga brevifolia* (Rottb.) Hassk. in temperate and tropical regions. *Journal of International Development and Cooperation*, 1.
- Rojas-Sandoval, J., & Acevedo-Rodríguez, P. (2013). *Sphagnetica trilobata* (wedelia). *CABI Compendium*. <https://doi.org/https://doi.org/10.1079/cabicompendium.56714>
- Rojas-Sandoval, J., & Acevedo-Rodríguez, P. (2014). *Eleusine indica* (goose grass). *CABI Compendium*. <https://doi.org/https://doi.org/10.1079/cabicompendium.20675>
- Russell, A. E. (2014). Unexpected Effects of Chitin, Cellulose, and Lignin Addition on Soil Dynamics in a Wet Tropical Forest. *Ecosystems*, 17, 918–930. <https://doi.org/10.1007/s10021-014-9769-1>
- Saberi, M., Davari, A., Tarnian, F., Shahreki, M., & Shahreki, E. (2013). Allelopathic Effects of *Eucalyptus camaldulensis* on Seed Germination and Initial Growth of four range species. *Scholars Research Library Annals of Biological Research*, 4(1), 152–159.
- Sachdev, S., Ansari, S. A., Ansari, M. I., Fujita, M., & Hasanuzzaman, M. (2021). Abiotic stress and reactive oxygen species: Generation, signaling, and defense mechanisms. *Antioxidants*, 10(2), 1–37. <https://doi.org/10.3390/antiox10020277>
- Saleem, S., Mushtaq, N. U., Shah, W. H., Rasool, A., & Hakeem, Khalid RehmanRehman, R. U. (2022). Beneficial role of phytochemicals in oxidative stress mitigation in plants. In *Antioxidant defense in plants: molecular basis of regulation* (pp. 435–451). Singapore: Springer Nature Singapore. https://doi.org/10.1007/978-981-16-7981-0_20
- Scavo, A., Abbate, C., & Mauromicale, G. (2019). Plant allelochemicals: agronomic, nutritional and ecological relevance in the soil system. *Plant and Soil*, 442, 23–48. <https://doi.org/10.1007/s11104-019-04190-y>
- Schandry, N., & Becker, C. (2020). Allelopathic Plants: Models for Studying Plant–Interkingdom Interactions. *Trends in Plant Science*, 25(2), 176–185.

<https://doi.org/10.1016/j.tplants.2019.11.004>

- Semmar, N. (2024). Overview on Different Biotic Interactions Involving Phytochemical Signals. In *Secondary Metabolites in Plant Stress Adaptation: Analytic Space of Secondary Metabolites* (pp. 311–315). Cham: Springer International Publishing.
- Slavković, F., & Bendahmane, A. (2023). Floral Phytochemistry: Impact of Volatile Organic Compounds and Nectar Secondary Metabolites on Pollinator Behavior and Health. *Chemistry and Biodiversity*, 20(4). <https://doi.org/10.1002/cbdv.202201139>
- Song, T., Chen, J. X., Shan, L. M., Qian, Y. C., Chen, M. X., Han, J. G., & Zhu, F. Y. (2024). Allelopathy research on the continuous cropping problem of poplar (populus). *Phytochemistry Reviews*, 6. <https://doi.org/10.1007/s11101-024-09948-6>
- Sothearith, Y., Appiah, K. S., Motobayashi, T., Watanabe, I., Somaly, C., Sugiyama, A., & Fujii, Y. (2021). Evaluation of allelopathic potentials from medicinal plant species in Phnom Kulen National Park, Cambodia by the sandwich method. *Sustainability*, 13(1), 264. <https://doi.org/10.3390/su13010264>
- Srivastava, P., Andersen, P. C., Marois, J. J., Wright, D. L., Srivastava, M., & Harmon, P. F. (2013). Effect of phenolic compounds on growth and ligninolytic enzyme production in *Botryosphaeria* isolates. *Crop Protection*, 43, 146–156. <https://doi.org/https://doi.org/10.1016/j.cropro.2012.09.015>
- Starý, P., & González, D. (1991). The *Chenopodium aphid*, *Hayhurstia atriplicis* (L.)(Horn., Aphididae), a parasitoid reservoir and a source of biocontrol agents in pest management. *Journal of Applied Entomology*, 111(1–5), 243–248.
- Surendran, A., Siddiqui, Y., Saud, H. M., Ali, N. S., & Manickam, S. (2017). The antagonistic effect of phenolic compounds on ligninolytic and cellulolytic enzymes of *Ganoderma boninense*, causing basal stem rot in oil palm. *International Journal of Agriculture and Biology*, 19(6), 1437–1446. <https://doi.org/10.17957/IJAB/15.0439>
- Sutherland, S. (2004). What makes a weed a weed: Life history traits of native and exotic plants in the USA. *Oecologia*, 141(1), 24–39. <https://doi.org/10.1007/s00442-004-1628-x>
- Takács, A., Cserpes, M., & Pásztor, G. (2018). The role of *Solanum nigrum* in the spread of potato viruses. *IX International Scientific Agriculture Symposium “AGROSYM 2018”, Jahorina, Bosnia and Herzegovina, 4-7 October 2018. Book of Proceedings*, 895–900.
- Talhi, F., Kalla, A., Gherraf, N., & Zellagui, A. (2020). Allelopathic effect of the aqueous extract of *Lantana camara* L. on the germination and development of four vegetable species. *International Journal of Chemical and Biochemical Sciences*, 18, 116–121.
- Tantiado, R. G., & Saylo, M. C. (2012). Allelopathic potential of selected grasses

- (Family Poaceae) on the germination of lettuce seeds (*Lactuca sativa*). *International Journal of Bio-Science and Bio-Technology*, 4(2), 27–34.
- Taylor, L. P., & Grotewold, E. (2005). Flavonoids as developmental regulators. *Current Opinion in Plant Biology*, 8(3), 317–323. <https://doi.org/10.1016/j.pbi.2005.03.005>
- Teasdale, J. R., & Mohler, C. L. (2000). The quantitative relationship between weed emergence and the physical properties of mulches. *Weed Science*, 48(3), 385–392. [https://doi.org/10.1614/0043-1745\(2000\)048\[0385:tqrbwe\]2.0.co;2](https://doi.org/10.1614/0043-1745(2000)048[0385:tqrbwe]2.0.co;2)
- Tian, M., Li, Q., Zhao, W., Qiao, B., Shi, S., Yu, M., Li, X., Li, C., & Zhao, C. (2022). Potential Allelopathic Interference of *Abutilon theophrasti* Medik. Powder/Extract on Seed Germination, Seedling Growth and Root System Activity of Maize, Wheat and Soybean. *Agronomy*, 12(4). <https://doi.org/10.3390/agronomy12040844>
- Tong, W. Y. (2013). Biotransformation of terpenoids and steroids. In *Natural products* (Issue May 2013, pp. 2733–2759). Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-22144-6>
- Treutter, D. (2005). Significance of flavonoids in plant resistance and enhancement of their biosynthesis. *Plant Biology*, 7(6), 581–591. <https://doi.org/10.1055/s-2005-873009>
- Valentino, M. J. G., Dulay, R. M. R., & Lindain, J. M. G. (2018). Phytochemicals, anti-oxidant potential and total phenolics of *Eleusine indica* leaf extracts. *International Journal of Biosciences*, 13(5), 317–321. <https://doi.org/10.12692/ijb/13.5.317-321>
- Valentovič, P., Luxová, M., Kolarovič, L., & Gašparíková, O. (2006). Effect of osmotic stress on compatible solutes content, membrane stability and water relations in two maize cultivars. *Plant, Soil and Environment*, 52(4), 186–191. <https://doi.org/10.17221/3364-pse>
- Vazquez, R. I., Stinner, B. R., & McCartney, D. A. (2003). Corn and weed residue decomposition in northeast Ohio organic and conventional dairy farms. *Agriculture, Ecosystems and Environment*, 95(2–3), 559–565. [https://doi.org/10.1016/S0167-8809\(02\)00176-7](https://doi.org/10.1016/S0167-8809(02)00176-7)
- Vilà, M., Beaury, E. M., Blumenthal, D. M., Bradley, B. A., Early, R., Laginhas, B. B., Trillo, A., Dukes, J. S., Sorte, C. J. B., & Ibáñez, I. (2021). Understanding the combined impacts of weeds and climate change on crops. *Environmental Research Letters*, 16(3), 034043.
- Wang, C., Wu, B., & Jiang, K. (2019). Allelopathic effects of Canada goldenrod leaf extracts on the seed germination and seedling growth of lettuce reinforced under salt stress. *Ecotoxicology*, 28(1), 103–116. <https://doi.org/10.1007/s10646-018-2004-7>
- Wang, C., Xiao, H., Zhao, L., Liu, J., Wang, L., Zhang, F., Shi, Y., & Du, D. (2016). The allelopathic effects of invasive plant *Solidago canadensis* on seed germination and growth of *Lactuca sativa* enhanced by different types of acid deposition. *Ecotoxicology*, 25(3), 555–562.

<https://doi.org/10.1007/s10646-016-1614-1>

- Wang, R., Liu, Y., Sun, S., Si, Y., Liu, X., Liu, X., Zhang, S., & Wang, W. (2020). Capsaicinoids from hot pepper (*Capsicum annuum* L.) and their phytotoxic effect on seedling growth of lettuce (*Lactuca sativa* L.). *Natural Product Research*, 34(11), 1597–1601. <https://doi.org/10.1080/14786419.2018.1519821>
- War, A. R., Buhroo, A. A., Hussain, B., Ahmad, T., Nair, R. M., & Sharma, H. C. (2019). Plant Defense and Insect Adaptation with Reference to Secondary Metabolites. In *Co-evolution of secondary metabolites* (pp. 1–28). Springer Nature Switzerland. https://doi.org/10.1007/978-3-319-76887-8_60-1
- Waterman, P. G. (1992). Roles for secondary metabolites in plants. In *Secondary metabolites: their function and evolution* (pp. 255–275). <https://doi.org/https://doi.org/10.1002/9780470514344.ch15>
- Weston, L. A., & Mathesius, U. (2013). Flavonoids: Their Structure, Biosynthesis and Role in the Rhizosphere, Including Allelopathy. *Journal of Chemical Ecology*, 39(2), 283–297. <https://doi.org/10.1007/s10886-013-0248-5>
- Woyessa, D. (2022). Weed Control Methods Used in Agriculture. *American Journal of Life Science and Innovation*, 1(1), 19–26. <https://doi.org/10.54536/ajlsi.v1i1.413>
- Xuan, T. D., Khanh, T. D., Chung, M. I., & Tawata, S. (2005). Cropping and Forage Systems / Crop Ecology / Organic Farming The Exploitation of Crop Allelopathy in Sustainable Agricultural Production. *Journal of Agronomy and Crop Science*, 191, 162–171. <http://www3.interscience.wiley.com/cgi-bin/fulltext/118738615/PDFSTART>
- Yang, W., Guo, Y., Li, Y., Zheng, Y., Dong, K., & Dong, Y. (2022). Cinnamic Acid Toxicity on the Structural Resistance and Photosynthetic Physiology of Faba Bean Promoted the Occurrence of Fusarium Wilt of Faba Bean, Which Was Alleviated Through Wheat and Faba Bean Intercropping. *Frontiers in Plant Science*, 13(June), 1–16. <https://doi.org/10.3389/fpls.2022.857780>
- Yasmin, M., Akonda, M. M. R., Rahman, M. M., & Seal, H. P. (2016). Assessment of allelopathic potential of different weeds on germination and early growth of aromatic rice. *Journal of Biodiversity and Environmental Sciences (JBES)*, 8(1), 236–241. <http://www.innspub.net/jbes/assessment-of-allelopathic-potential-of-different-weeds-on-germination-and-early-growth-of-aromatic-rice/>
- Yazaki, K., Arimura, G. I., & Ohnishi, T. (2017). “Hidden” terpenoids in plants: Their biosynthesis, localization and ecological roles. *Plant and Cell Physiology*, 58(10), 1615–1621. <https://doi.org/10.1093/pcp/pcx123>
- Yu, J. Q., & Matsui, Y. (1997). Effects of root exudates of cucumber (*Cucumis sativus*) and allelochemicals on ion uptake by cucumber seedlings. *Journal of Chemical Ecology*, 23(3), 817–827.
- Zhao, J., Yang, Z., Zou, J., & Li, Q. (2022). Allelopathic effects of sesame extracts on seed germination of moso bamboo and identification of potential

allelochemicals. *Scientific Reports*, 12(1), 6661.
<https://doi.org/10.1038/s41598-022-10695-x>

Ziegler, J., & Facchini, P. J. (2008). Alkaloid biosynthesis: Metabolism and trafficking. *Annual Review of Plant Biology*, 59, 735–769.
<https://doi.org/10.1146/annurev.arplant.59.032607.092730>

Zimdahl, R. L., & Basinger, N. T. (2024). *Fundamentals of weed science*. Elsevier.

Zohaib, A., Abbas, T., & Tabassum, T. (2016). Weeds Cause Losses in Field Crops through Allelopathy. *Notulae Scientia Biologicae*, 8(1), 47–56.
<https://doi.org/10.15835/nsb.8.1.9752>

Zohra, Z. F., Benamar, H., Bennaceur, M., & Abderrazak, M. (2019). Allelopathic effect of *Tetralinis articulata* (vahl) mast (from Algeria) on germination and growth of *Lactuca sativa* L. *Bioscience Research*, 16(1), 493–499.

