

DAFTAR PUSTAKA

1. Amarya S, Singh K, Sabharwal M. Ageing process and physiological changes. In: D'Onofrio G, Greco A, Sancar D, editors. *Gerontology*. InTech; 2018:1–12.
2. Larsson L, Degens H, Li M, Salvati L, Lee YI, Thompson W, et al. Sarcopenia: aging-related loss of muscle mass and function. *Physiol Rev*. 2019;99(1):427–511.
3. European Commission Statistical Office of the European Union. *Ageing europe: looking at the lives of older people in the EU : 2020 edition*. LU: Publications Office; 2020:15–32.
4. Kementerian Hukum dan Hak Asasi Manusia Republik Indonesia. *Undang-Undang Republik Indonesia Nomor 13 Tahun 1998 Tentang Kesejahteraan Lanjut Usia*. Badan Pembinaan Hukum Nasional; 1998.
5. United Nations Department of Economic and Social Affairs. *World Social Report 2023: Leaving no one behind in an ageing world*. United Nations; 2023:17–29.
6. *World population prospects 2022: summary of results*. New York: United Nations; 2022:3–39.
7. Direktorat Statistik Kesejahteraan Rakyat. *Statistik Penduduk Lanjut Usia 2023*. Jakarta: Badan Pusat Statistik; 2023.
8. Maresova P, Javanmardi E, Barakovic S, Barakovic Husic J, Tomsone S, Krejcar O, et al. Consequences of chronic diseases and other limitations associated with old age – a scoping review. *BMC Public Health*. 2019;19(1):1431–42.
9. Laporan Tahunan Dinas Kesehatan Kota Padang 2023. Edisi 2024. Diakses https://dinkes.padang.go.id/uploads/audios/dinkes_66d004fc297f0.pdf.
10. Lin X, Li H. Obesity: epidemiology, pathophysiology, and therapeutics. *Frontiers in Endocrinology*. Frontiers Media S.A.; 2021; 12:1-9.
11. Kementerian Kesehatan Republik Indonesia. *Laporan nasional Riskesdas 2018*. 2018. 57–578 p.

12. Kemenkes. Survei Kesehatan Indonesia (SKI) 2023 [Internet]. 2023. Available from: https://drive.google.com/file/d/1rjNDG_f8xG6-Y9wmhJUnXhJ-vUFevVJC/view
13. Choi K, Jang HY, Ahn MJ, Hwang SH, Chung JW, et al. The association of the serum levels of myostatin, follistatin, and interleukin-6 with sarcopenia, and their impacts on survival in patients with hepatocellular carcinoma. National Center for Biotechnology Information. Clin Mol Hepatol. 2020;26(4):492–505. doi: [10.3350/cmh.2020.0005](https://doi.org/10.3350/cmh.2020.0005)
14. Curtis M, Swan L, Fox R, Warters A, O’Sullivan M. Associations between body mass index and probable aarcopenia in community-dwelling older adults. National Center for Biotechnology Information. Nutrients. 2023;15(6):1505. doi: [10.3390/nu15061505](https://doi.org/10.3390/nu15061505)
15. Aras Ş, Üstünsoy S, Armutçu F. Indices of central and peripheral obesity: anthropometric measurements and laboratory parameters of metabolic syndrome and thyroid function. Balkan Med J. 2015;32:414–20.
16. Álvarez R, Torres J, Artola G, Epelde G, Arranz S, Marrugat G. Obinter: A holistic approach to catalyse the self-management of chronic obesity. Sensors (Switzerland). 2020;20(18):1–27.
17. Han DR, Chang KV, Li CM, Lin YH, Kao TW, et al. Skeletal muscle mass adjusted by height correlated better with muscular functions than that adjusted by body weight in defining sarcopenia. Scientific Report. 2016; (6):19457. doi: [10.1038/srep19457](https://doi.org/10.1038/srep19457)
18. Sayer AA, Cruz-Jentoft A. Sarcopenia definition, diagnosis and treatment: consensus is growing. Age Ageing. 2022;51(10):220–35.
19. Shafiee G, Keshtkar A, Soltani A, Ahadi Z, Larijani B, Heshmat R. Prevalence of sarcopenia in the world: a systematic review and meta- analysis of general population studies. J Diabetes Metab Disord. 2017;16(1):21–32.
20. Mayhew AJ, Amog K, Phillips S, Parise G, McNicholas PD, De Souza RJ, et al. The prevalence of sarcopenia in community-dwelling older adults, an exploration of differences between studies and within definitions: a systematic review and meta-analyses. Age Ageing. 2019;48(1):48–56.

21. Kim M, Won CW. Sarcopenia in korean community-dwelling adults aged 70 years and older: Application of screening and diagnostic tools from the Asian Working Group for Sarcopenia 2019 update. *J Am Med Dir Assoc.* 2020;21(6):752–8.
22. Makizako H, Nakai Y, Tomioka K, Taniguchi Y. Prevalence of sarcopenia defined using the Asia Working Group for Sarcopenia criteria in japanese community-dwelling older adults: A systematic review and meta-analysis. *Phys Ther Res.* 2019;22(2):53–7.
23. Liu X, Hou L, Xia X, Liu Y, Zuo Z, Zhang Y, et al. Prevalence of sarcopenia in multi ethnics adults and the association with cognitive impairment: findings from West-China health and aging trend study. *BMC Geriatr.* 2020;20(1):63–74.
24. Harimurti K, Setiati S, Soejono CH, Aryana IS, Sunarti S, Budiningsih F, et al. Sarcopenia in a multiethnic state: a cross-sectional data analysis of multicentre Indonesia longitudinal aging study. *Acta Med Indones.* 2023;55(1):1–11.
25. McKay BR, Ogborn DI, Bellamy LM, Tarnopolsky MA, Parise G. Myostatin is associated with age-related human muscle stem cell dysfunction. *The FASEB Journal.* 2012; 26: 2509–21
26. Peng LN, Lee WJ, Lin MH, Chen LK. Healthy community-living older men differ from women in association between myostatin levels and skeletal muscle mass. *Journal of Cachexia, Sarcopenia and Muscle.* 2018; 10: 1–8
27. Yarasheski KE, Bhasin SH, Hikim IS, Loduca JP, Cadavid NFG. Serum Myostatin-immunoreactive protein is increased in 69-92 year old women and men with muscle wasting. *The Journal of Nutrition, Health & Aging.* 2002; 6: 1–6
28. Hamasaki H, Kawashima Y, Katsuyama H, Sako A, Goto A. Association of handgrip strength with hospitalization, cardiovascular events, and mortality in Japanese patients with type 2 diabetes. *Sci Rep.* 2017;7:1–9
29. Beavers KM, Beavers DP, Houston DK, Harris TB, Hue TF, Koster A, et al. Associations between body composition and gait-speed decline: results from the health, aging and body composition study. *Am J Clin Nutr.* 2013;97(6):552–60.
30. Moon JJ, Park S, Ryu SM, Park C, Park S. New skeletal muscle mass index in

diagnosis of sarcopenia. *J Bone Metab.* 2018;25:15–21.

31. Franzago M, Di Nicola M, Fraticelli F, Marchioni M, Stuppia L, Vitacolonna E. Nutrigenetic variants and response to diet/lifestyle intervention in obese subjects: a pilot study. *Acta Diabetol.* 2022 Jan 1;59(1):69–81.
32. Lin X, Li H. Obesity: epidemiology, pathophysiology, and therapeutics. *Frontiers in Endocrinology.* Frontiers Media S.A.; 2021; 12:1-9
33. PERKENI. Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Dewasa di Indonesia.
34. Hubbard RE, Lang IA, Llewellyn DJ, Rockwood K. Frailty, body mass index, and abdominal obesity in older people. *J Gerontol.* 2009;65(4):1–5.
35. International Working Group on Sarcopenia. Sarcopenia: an undiagnosed condition in older adults - current consensus definition: prevalence, etiology, and consequences. *J Am Med Dir Assoc.* 2012;12(4):249–56.
36. Baumgartner RN. Body composition in healthy aging. *Ann New York Acad Sci.* 2000;904:437–48.
37. Dhillon RJ. Pathogenesis and management of sarcopenia. *Clin Geriatr Med.* 2017;33(3):17–26.
38. Donini LM, Busetto L, Bauer JM, Bischoff S, Boirie Y, Cederholm T, et al. Critical appraisal of definitions and diagnostic criteria for sarcopenic obesity based on a systematic review. *Clin Nutr.* 2019;39(8):1–21.
39. Pruner L, Cararo S, Mortean L, Rodrigues A. Quality of life associated with handgrip strength and sarcopenia: EpiFloripa Aging Study. *Arch Gerontol Geriatr.* 2019;81:234–9.
40. Carvalho DHT de, Santos JLF, Oliveira C de, Alexandre T da S. Does abdominal obesity accelerate muscle strength decline in older adults? Evidence from the English longitudinal study of ageing. *Journals Gerontol Med Sci.* 2019;74(7):1105–11.
41. Hassinen M, Komulainen P, Lakka TA, Väisänen SB, Rauramaa R. Associations of body composition and physical activity with balance and walking ability in the elderly. *J Phys Act Heal.* 2005;3:298–306.
42. Pruner L, Cararo S, Mortean L, Rodrigues A. Quality of life associated with handgrip strength and sarcopenia: EpiFloripa Aging Study. *Arch Gerontol*

Geriatr. 2019;81:234–9.

43. Santos CM, Amorim G, Samora R, Guerra RO. Prevalence of obesity, sarcopenic obesity and associated factors: A FIBRA Network study. *Fisioter.* 2017;30:S161- 9.
44. Lee Y, Chen C, Lee Y, Chen C, Lee C. Body anthropometric measurements of Singaporean adult and elderly population. *Measurement.* 2019;148:1–9.
45. Aryana IGPS. Sarkopenia pada lansia: Problem diagnosis dan tatalaksana. Baswara Press; 2021:1–148.
46. Widajanti N, Ichwani J, Dharmanta RS, Firdausi H, Haryono Y, Yulianti E, et al. Sarcopenia and frailty profile in the elderly community of surabaya: A descriptive study. *Acta Med Indones.* 2020;52(1):1–12.
47. Laksmi PW, Setiati S, Tamin TZ, Soewondo P. Effect of metformin on handgrip strength, gait speed, myostatin serum level, and health-related quality of life: a double blind randomized controlled trial among non-diabetic pre-frail elderly patients. *Acta Medica Indones - Indones J Intern Med.* 2017;49(2):118–27.
48. Cannataro R, Carbone L, Petro JL, Cione E, Vargas S, Angulo H, et al. Sarcopenia: etiology, nutritional approaches, and miRNAs. *Int J Mol Sci.* 2021;22(18):9724.
49. Saadeddine D, Itani L, Kreidieh D, El Masri D, Tannir H, El Ghoch M. Association between levels of physical activity, sarcopenia, type 2 diabetes and the quality of life of elderly people in community dwellings in Lebanon. *Geriatrics.* 2021;6(1):28.
50. Hastuti J, Kagawa M, Byrne NM. Determination of new anthropometric cut-off values for obesity screening in Indonesian adults. *Asia Pac J Clin Nutr.* 2017;26:650–6.
51. Stenholm S, Sallinen J, Koster A, Rantanen T, Sainio P, Heliövaara M, et al. Association between obesity history and hand grip strength in older adults — exploring the roles of inflammation and insulin resistance as mediating factors. *J Gerontol.* 2011;66A(3):341–8.
52. Zeghibi V, Borba C, Costa TL, Moreira CA. Sarcopenia in endocrine and non-endocrine disorders. *Eur Soc Endocrinol.* 2019;180:R185–99.

53. Rolland Y, Czerwinski S, Kan GA Van, Morley JE, Cesari M, Onder G, et al. Sarcopenia: its assessment, etiology, pathogenesis, consequences, and future perspectives. *J Nutr Heal Aging*. 2014;12(7):433–50.
54. Dhillon RJ. Pathogenesis and management of sarcopenia. *Clin Geriatr Med*. 2017;33(3):17–26.
55. Cho MR, Lee S, Song SK. A review of sarcopenia pathophysiology, diagnosis, treatment and future direction. *J Korean Med Sci*. 2022;37(18):146–61.
56. Dhar M, Kapoor N, Suastika K, Khamseh ME, Selim S, Kumar V, et al. South Asian Working Action Group on SARCOpenia (SWAG-SARCO) – A consensus document. *Osteoporos Sarcopenia*. 2022;8(2):35–57.
57. Chen L, Woo J, Assantachai P, Auyeung T, Chou M, Iijima K, et al. Asian working group for sarcopenia: 2019 consensus update on sarcopenia diagnosis and treatment. *J Am Med Dir Assoc*. 2020;1–8.
58. Kwak JY, Hwang H, Kim SK, Choi JY, Lee SM, Bang H et al. Prediction of sarcopenia using a combination of multiple serum biomarkers. *Scientific Reports Journal*. 2018; 8: 1–7
59. Edwin NN, I Gusti PSA. Sarkopenia pada usia lanjut: patogenesis, diagnosis, dan tatalaksana. *Jurnal Penyakit Dalam Indonesia*. 2023. 10(3):163-73.
60. Jan B, Marta S, Zyta BW. Myostatin as a biomarker of muscle wasting and other pathologies-state of the art and knowledge gaps. *Nutrient*. 2020. 16(6).1-22
61. World Health Organization. Obesity and Overweight [Internet]. 2021. Diakses : <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>
62. International Diabetes Institute WHOCC for the E of DM and HP for ND. The Asia-Pacific perspective: redefining obesity and its treatment. 2000; p. 8–24.
63. Wharton S, Lau DCW, Vallis M, Sharma AM, Biertho L, Campbell-Scherer D, et al. Obesity in adults: A clinical practice guideline. *CMAJ*. 2020;192(31):E875–91.
64. Gjermani E, Kirstein AS, Kolbig F, Kirchhof M, Bundalian L, Katzmann JL, et al. Obesity—an update on the basic pathophysiology and review of recent therapeutic advances. *Biomolecules*. MDPI. 2021;11:1-28
65. Oussaada SM, van Galen KA, Cooman MI, Kleinendorst L, Hazebroek EJ, van

- Haelst MM, et al. The pathogenesis of obesity. *Metabolism*. 2019;92:26–36.
66. Batsis JA, Villareal DT. Sarcopenic obesity in older adults: aetiology, epidemiology and treatment strategies. *Nat Rev Endocrinol*. 2019;14(9):513–37.
 67. Bohn B, Simic-schleicher G, Kiess W, Siegfried W, Oelert M, Tuschy S, et al. BMI or BIA: is body mass index or body fat mass a better predictor of cardiovascular risk in overweight or obese children and adolescents? *Eur J Obes*. 2015;156–65.
 68. Yanai H, Hirowatari Y. Correlations of body mass index, Japan-defined and IDF- defined waist circumference, visceral and subcutaneous fat area with metabolic parameters in Japanese patients with type 2 diabetes. *J Diabetes Mellit*. 2015;5:290–4.
 69. American Association of Clinical Endocrinologist. What is the disease of obesity? Anthropometric evaluation. AACE/ACE Algorithm Med Care Patients with Obes. 2016;1–17.
 70. International Diabetes Federation. The IDF consensus worldwide definition of the metabolic syndrome. 2006;11.
 71. Prado, C.M., Batsis, J.A., Donini LM et al. Sarcopenic obesity in older adults: A clinical overview. *Nat Rev Endocrinol*. 2024;20:261–77.
 72. Prado CM, Cushen SJ, Orsso CE, Ryan AM. Sarcopenia and cachexia in the era of obesity: clinical and nutritional impact. *Proc Nutr Soc*. 2016;75:188– 98.
 73. Lee D, Drenowatz C, Blair SN. Physical activity and sarcopenic obesity: definition, assessment, prevalence and mechanism. *Futur Sci*. 2016;2(3):1–19.
 74. Kalinkovich A, Livshits G. Sarcopenic obesity or obese sarcopenia: A cross talk between age-associated adipose tissue and skeletal muscle inflammation as a main mechanism of the pathogenesis. *Ageing Res Rev*. 2019;35(2017):200–21.
 75. Zhang X, Xie X, Dou Q, Liu C, Zhang W, Yang Y, et al. Association of sarcopenic obesity with the risk of all-cause mortality among adults over a broad range of different settings: a updated meta-analysis. *BMC Geriatr*. 2019;19(183):1–14.
 76. Nascimento D da C, Oliveira S da C, Vieira DCL, Funghetto SS, Silva AO,

- Vadulga R, et al. The impact of sarcopenic obesity on inflammation, lean body mass, and muscle strength in elderly women. *Int J Gen Med*. 2018;11:443–9.
77. Nasimi N, Hossein M, Sohrabi Z. Nutritional status and body fat mass: Determinants of sarcopenia in community-dwelling older adults. *Exp Gerontol*. 2019;122(2019):67–73.
 78. Tabue-Teguo M, Perès K, Simo N, Goff M Le, Zepeda MUP, Feart C, et al. Gait speed and body mass index: results from the AMI study. *PLoS One*. 2020;15(3):1–8.
 79. Firdaus MR, Cahyawati WASN, Yasmina A, Noor Z, Wahyuni. Hubungan indeks massa tubuh dengan kecepatan berjalan pada lansia di PPRSLU Budi Sejahtera Banjarbaru. *Homeostasis*. 2024; 7(1): 139-46.
 80. Chen H, Hu H. The relationship and mechanism between education and functional health status transition among older persons in China. *BMC Geriatr*. 2018;18(1):89.
 81. Zhang CQ, Chung PK, Zhang R, Schüz B. Socioeconomic Inequalities in Older Adults' Health: The Roles of Neighborhood and Individual-Level Psychosocial and Behavioral Resources. *Front Public Health*. 2019;7:318.
 82. Zhu X, Ning B, Xia F, Wang B, Li Y, Zhang P, et al. The impact of widowhood on the quality of life of older adults: the mediating role of intergenerational support from children. *BMC Geriatr*. 2024;24(1):658.
 83. Wang X, Miao H, Jin Q. The structural characteristics of the lifestyle among older adults and its impact on the health in China. *Front Public Health*. 2023;11:1286530.
 84. Tan You Mei C, Seah Si Ying S, Yanshan DL, Koh SV, Karthikeyan G, Xia Jiawen O, et al. Prevalence and factors associated with sarcopenia among older adults in a post-acute hospital in Singapore. *PLoS One*. 2024;19(1):e0291702.
 85. Xiao H, Huang S, Xiao H, Zhang W, Cai S. Longitudinal association between body mass index and handgrip strength in community-dwelling older adults: a population-based nationwide cohort study. *BMC Geriatr*. 2025;25(1):711.
 86. Sallinen J, Stenholm S, Rantanen T, Heliövaara M, Sainio P, Koskinen S. Hand-grip strength cut points to screen older persons at risk for mobility limitation. *J Am Geriatr Soc*. 2010;58(9):1721-6.

87. Beavers KM, Beavers DP, Houston DK, Harris TB, Hue TF, Koster A, et al. Associations between body composition and gait-speed decline: results from the Health, Aging, and Body Composition study. *Am J Clin Nutr*. 2013;97(3):552-60.
88. Raza A, Mahmood I, Sultana T, Sultana S. Differences in gait dynamic stability of healthy subjects due to age, body mass, height, BMI, and walking speed. *Medicine in Novel Technology and Devices*. 2025; 26.
89. Correia de Faria Santarém G, de Cleve R, Santo MA, Bernhard AB, Gadducci AV, Greve JMD, et al. Correlation between Body Composition and Walking Capacity in Severe Obesity. *PLoS ONE*. 2015; 10(6): e0130268.
90. Alley DE, Ferrucci L, Barbagallo M, Studenski SA, Harris TB. A research agenda: the changing relationship between body weight and health in aging. *J Gerontol A Biol Sci Med Sci*. 2008;63(11):1257-9.
91. Ponti F, Santoro A, Mercatelli D, Gasperini C, Conte M, Martucci M, et al. Aging and Imaging Assessment of Body Composition: From Fat to Facts. *Front Endocrinol (Lausanne)*. 2020;10:861.
92. Jeong SM, Lee DH, Rezende LFM, Giovannucci EL. Different correlation of body mass index with body fatness and obesity-related biomarker according to age, sex and race-ethnicity. *Sci Rep*. 2023;13(1):3472.
93. Tey SL, Chew STH, How CH, Yalawar M, Baggs G, Chow WL, et al. Factors associated with muscle mass in community-dwelling older people in Singapore: Findings from the SHIELD study. *PLoS One*. 2019;14(10):e0223222.
94. Cheng Q et al. A cross-sectional study of loss of muscle mass corresponding to sarcopenia in healthy Chinese men and women: reference values, prevalence, and association with bone mass. *J Bone Miner Metab*. 2014;32(1):78–85
95. Marulanda-Díaz N, Estrada-Restrepo A, Milán AF, Narvaez-Sanchez R, Calderón JC, Giraldo-Giraldo NA. Relationship between myostatin, musclin, nutritional status, and functionality in older Colombian community-dwelling adults: A cross-sectional study. *Nutrition*. 2025;135:112767.

96. Bojar I, Mlak R, Homa-Mlak I, Prendecka M, Owoc A, Małecka-Massalska T. Association between myostatin serum concentration and body fat level in peri- and postmenopausal women. *Arch Med Sci.* 2021;18(2):365-375.
97. Perry CA, Van Guilder GP, Butterick TA. Decreased myostatin in response to a controlled DASH diet is associated with improved body composition and cardiometabolic biomarkers in older adults: results from a controlled-feeding diet intervention study. *BMC Nutr.* 2022;8(1):24.
98. Szulc P, Schoppet M, Goettsch C, Rauner M, Dschietzig T, Chapurlat R, et al. Endocrine and clinical correlates of myostatin serum concentration in men--the STRAMBO study. *J Clin Endocrinol Metab.* 2012;97(10):3700-8.
99. Fife E, Kostka J, Kroc Ł, Guligowska A, Pięłowska M, Sołtysik B, et al. Relationship of muscle function to circulating myostatin, follistatin and GDF11 in older women and men. *BMC Geriatr.* 2018;18(1):200.
100. Takao N, Kurose S, Miyauchi T, Onishi K, Tamanoi A, Tsuyuguchi R, et al. The relationship between changes in serum myostatin and adiponectin levels in patients with obesity undergoing a weight loss program. *BMC Endocr Disord.* 2021;21(1):147.
101. Xu HQ, Shi JP, Shen C, Liu Y, Liu JM, Zheng XY. Sarcopenia-related features and factors associated with low muscle mass, weak muscle strength, and reduced function in Chinese rural residents: a cross-sectional study. *Arch Osteoporos.* 2018;14(1):2.
102. Njoto EN, Suka Aryana IGP. Sarkopenia pada lanjut usia: patogenesis, diagnosis dan tata laksana. *J Penyakit Dalam Indones.* 2023;10(3):164-73.