

## DAFTAR PUSTAKA

- Adelman C, Freeman S, Paz Z, Sohmer H(2008). Salicylic acid injection before noise exposure reduces permanent threshold shift. *Audiol Neurotol*, 13, pp.266-72.
- Ahmadi (2009). *Psikologi Sosial*, Jakarta: Rineka Cipta. Cetakan Ketiga.
- Ahmed S, Fallah S, Garrido B, Gross A, King M, Morrish T, Pereira D, et al (2007). Use of portable audio devices. *Journal from Department of Psychology Universitas Toronto* [Online], 35: 35-52.
- Alberti PW (1997). Noise and the ear. In: Stephens D, ed 6th. *ScottBrown's Adult Audiology*. Great Britain: Butterworth-Heinemann, pp.1-34.
- American Hearing Research Foundation(2012).Noise Induced Hearing Loss. <http://American-Hearing.org/disorders/noise-induced-hearing-liss/#Whatis> Diakses Pada Tanggal 6 Oktober 2015 pukul 9.11 WIB.
- American Speech Language Hearing Association(2006).Survey of Teens and Adults About The Use of Personal Electronic Devices and Head Phones. <http://www.nsslha.org/NR/rdonlyres/10B67FA1-002C-4C7BBAOB-1C0A3AF98A63/0/zogbysurvey2006>.Diakses Pada Tanggal 11 Agustus 2015 pukul 21.00 WIB.
- Attias J, Sapir S, Bresloff I, Reshef-Haran I, Ising H (2004).Reduction in noise-induced temporary threshold shift in humans following oral magnesium intake.*Clin Otolaryngol*, 29, pp.635-41.
- Azwar, Saifuddin (2007). *Sikap Manusia : Teori dan Pengukurannya*. Yogyakarta : Pustaka Pelajar Offset.
- Bashiruddin J, Soetirto I (2012). *Gangguan Pendengaran Akibat Bising*.Jakarta : Badan Penerbit FK UI.
- Centers for Disease Control and Prevention (2011).Noise induced hearing loss. [www.cdc.gov/healthyyouth/noise](http://www.cdc.gov/healthyyouth/noise). Diakses pada Tanggal 28 Januari 2016 Pukul 23.15 WIB
- Dayakisni, Tri dan Hudaniah (2003).*Psikologi Sosial*. Malang: UMM Press.
- Fligor B. J. 2011, October 31, 2011. Recreational NIHL: A Summary of The Evidence. *Audiology Online Cyber Media*.<http://www.audiologyonline.com/articles/recreational-nihl-summary-evidence-800>. Diakses pada Tanggal 28 Januari 2016 Pukul 23.20.
- Gacek RR (2009). *Anatomy of the Auditory and Vestibular System in : Snow jr JB and Wackym PA Ballenger's. Otorhinolaryngology Head and Neck*

Surgery 17,Centennial edition. Philadhelpia. People's Medical Publishing House. p. 1- 157.

- Goldberg LR dan Richburg CM (2004). Minimal hearing impairment: Major myths with more than minimal implications. *Communication Disorders Quarterly*, 25:152–160.
- Guitierrez B, Moledero L (2007). Listening to music with headphoe: An assessment of noise exposure and hearing Damage. In *Headphone Sound Exposure and Hearing*: 59-146.
- Hadinoto SO (2014).Gambaran pengetahuan, sikap dan perilaku remaja tentang gangguan pendengaran akibat penggunaan Piranti dengar. Surabaya, Universitas Katolik Widya Mandala. Skripsi.
- Harrianto R 2013.Buku Ajar Kesehatan Kerja. Jakarta: EGC.
- Harrison RV(2008). Noise-induced hearing loss in children: A less than silent" enviromental danger. *Paediatrics and Child Health*, 13(5): 377-382.
- Hodgetts WE, Rieger JM, Szarko RA (2007). The Effects of Listening Environment and Earphone Style on Preferred Listening Levels of Normal Hearing Adults Using An MP3 Player. *Ear Hear*,28: 290-297.
- Irawati L (2012). Fisika Medik Proses Pendengaran. *Majalah Kedokteran Andalas*, 36(2): 155-162.
- Irmayanti M(2007). *Pengetahuan Dalam Kesehatan*. Jakarta: Lembaga Penerbitan FEUI.
- Isaacson J, dan Vora N (2003). American family physician review: Differential diagnosis and treatment of hearing loss. *Am Fam Phys*, 68(6): 1125-1134.
- Kim MG (2009). Hearing treshold of Korean adolescents associated with the use of personal music players. *Yonsei Medical Jurnal*, 50(6): 771-776.
- Laoh Alvin (2015). Hubungan penggunaan headset terhadap fungus pendengaran pada mahasiswa angkatan 2012 Fakultas Kedokteran Universitas Sam Ratulangi. *Jurnal Kedokteran Komunitas dan Tropiki*, 3(3):142-147.
- Levey S, Levey T, dan Fligor BJ (2012).The Effects of Noise-Induced Hearing Loss on Children and Young Adults.Contemporary Issues in Communication Science and Disorder, 39: 76-83.
- Liston SL. dan Duvalu AJ (1997).Embriologi, anatomi dan fisiologi telinga.Dalam: Adams, GL, Boeis, LR & Highler, PA. Boeis Buku Ajar Penyakit THT.Jakarta : EGC, 27-45.
- Martini FH, Nath JL, Bartholomew EF(2012). *Fundamentals of Anatomy & Physiology*. 9th ed. US: Benjamin Cummings.

- McCaffree MA (2008). Portable listening devices and noise-induced hearing loss. American Medical Association. [www.ama-assn.org/resources/doc/csaph/csaph6a08](http://www.ama-assn.org/resources/doc/csaph/csaph6a08). Diakses pada Tanggal 26 Januari 2016 Pukul 10.17.
- Moller AR (2006). Hearing Anatomy, Physiology, and Disorders of the Auditory System 2nd ed. Texas. Elsevier. Pp 41- 56.
- National Institute on Deafness and Other Communication Disorders. 2013. Noise-Induced Hearing Loss. <http://www.nidcd.nih.gov/health/hearing/pages/noise.aspx>. Diakses pada Tanggal 26 Januari 2016 Pukul 10.15 WIB.
- Nelson TJ (2009). Noise-induced hearing loss. Science notes. <http://www.randombio.com/noise.html>. Diakses pada Tanggal 29 Januari 2016 Pukul 1.47 WIB.
- Notoatmodjo S (2003). Pengantar Pendidikan Kesehatan Dan Ilmu Perilaku Kesehatan, Yogyakarta: Andi Offset. Edisi I.
- Notoatmodjo S (2012). Promosi kesehatan dan perilaku kesehatan. Jakarta: Rineka Cipta.
- Oghalai JS dan Brownell WE (2008). Anatomy and physiology of the ea. In Lalwani , A.K. Current Diagnosis & Treatment in Otolaryngology-Head and Neck Surgery. New York : McGraw-Hill Company 577-95.
- Purba, Juniarti (2013). Gambaran Faktor - Faktor Yang Memengaruhi Tindakan Ibu Terhadap Pemberian Imunisasi Dasar Lengkap Dan Tidak Lengkap Pada Balita (12 Bulan) Di Desa Secanggih Kecamatan Secanggih Kabupaten langkat Tahun 2013. Medan. Universitas Sumatera Utara. Skripsi.
- Portnuff CDF, Fligor BJ, dan Arehart KH (2011). Teenage Use of Portable Listening Devices: a Hazard to Hearing?. Journal of the American Academy of Audiology, 22: 663-677.
- Rabinowitz PM. Hearing Loss and Personal Music Players. BMJ (2010). <http://www.bmj.com/content/340/bmj.c1261.full>. Diakses pada Tanggal 29 Januari 2016 Pukul 1.17.
- Rahadian J, Prastowo NA, Haryono R (2010). Pengaruh penggunaan earphone terhadap fungsi pendengaran remaja. Majalah Kedokteran Indonesia 60 : 468-473.
- Robert AD (2002). Noise induced hearing loss. Current Diagnosis and Treatment in Otolaryngology-Head and Neck Surgery. New York: McGraw Hill.

- Robert AD (2006). Noise induced hearing loss. In Bailey BJ, ed IV. Head and Neck Surgery-Otolaryngology. Baltimore: Lippincott Williams & Wilkins: 2190-5.
- Sarwono S(2012).*Sosiologi Kesehatan Beberapa Konsep dan Aplikasinya* (edisi ke 4). Yogyakarta: Gadjah Mada University Press.
- Scientific Committee on Emerging and Newly Identified Health Risks (2008). Potential health risks of exposure to noise from personal music players and mobile phones including a music playing function. [http://ec.europa.eu/health/ph\\_risk/committees/04\\_scenihr/docs/scenihr\\_o\\_018](http://ec.europa.eu/health/ph_risk/committees/04_scenihr/docs/scenihr_o_018).Diakses pada Tanggal 29 Januari 2016 Pukul 1.13 WIB.
- Vogel I, Brug J, Van der Ploeg CP, Raat H (2011). Adolescents Risky MP3-Player Listening and Its Psychosocial Correlates. *Health Aeduc Res*, 26: 254-264
- Wittman R dan Scott K (2006).Survey of adults in Mexico and Puerto Rico about the use of personal electronic devices with headphone. *Zogby Internasional*, 1-26.
- World Health Organization(1997). Prevention of noise-induced hearing loss: Report of an informal consultation. Makalah disampaikan dalam *Seminar Report of an Informal Consultation*, WHO-PDH Geneva, 28-30 Oktober.
- World Health Organization (2007). State of hearing & ear care in south east asiaregion. WHO Regional Office SEARO.[http://who.int/pbd/deafness/activities/hearing\\_care/advance](http://who.int/pbd/deafness/activities/hearing_care/advance).Diakses pada Tanggal 18 Januari 2016 Pukul 22.00.

